

Math 322 Fall 2014 (Lior Silberman): Problem Set 1, due 11/9/2014

Practice and supplementary problems, and any problems specifically marked “OPT” (optional), “SUPP” (supplementary) or “PRAC” (practice) are *not for submission*. It is possible that the grader will not mark all problems.

Practice problems

P1 Find all solutions to the congruence $5x \equiv 1 \pmod{7}$.

The following problem is a review of the axioms for a vector space.

P2 Let X be a set. Carefully show that pointwise addition and scalar multiplication endow the set \mathbb{R}^X of functions from X to \mathbb{R} with the structure of an \mathbb{R} -vector space. Meditate on the case $X = [n] = \{0, 1, \dots, n-1\}$.

P3 (Euclid’s Lemma) Let a, b, q, r be four integers with $b = aq + r$. Show that the pairs $\{a, b\}$ and $\{a, r\}$ have the same sets of common divisors, hence the same greatest common divisor.

The integers

1. Show that for any integer k , one of the integers $k, k+2, k+4$ is divisible by 3.
2. Consider the equation $13x + 5y = 1$ for unknowns $x, y \in \mathbb{Z}$.
PRAC Exhibit a solution.
(a) Exhibit infinitely many solutions.
(*b) Show that you found *all* the solutions.
3. Let a, n be positive integers and let $d = \gcd(a, n)$. Show that the equation $ax \equiv 1 \pmod{n}$ has a solution iff $d = 1$.
4. Let $f: \mathbb{Z} \rightarrow \mathbb{Z}$ be *additive*, in that for all $x, y \in \mathbb{Z}$ we have $f(x+y) = f(x) + f(y)$.
PRAC Check that for any $a \in \mathbb{Z}$, $f_a(x) = ax$ is additive.
(a) Show that $f(0) = 0$ (hint: $0+0=0$).
(b) Show that $f(-x) = -f(x)$ for all $x \in \mathbb{Z}$.
(c) Show by induction on n that for all $n \geq 1$, $f(n) = f(1) \cdot n$.
(d) Show that every additive map is of the form f_a for some $a \in \mathbb{Z}$.
RMK Let H be the set of additive maps $\mathbb{Z} \rightarrow \mathbb{Z}$. We showed that the function $\varphi: H \rightarrow \mathbb{Z}$ given by $\varphi(f) = f(1)$ is a bijection (with inverse $\psi(a) = f_a$).
SUPP Show that the bijections φ, ψ are themselves additive maps (addition in H is defined pointwise).

Supplementary problems I: Functions, sets and relations

The following problem will be used in the upcoming discussion of permutations.

- A. Let X, Y, Z, W be sets and let $f: X \rightarrow Y$, $g: Y \rightarrow Z$ and $h: Z \rightarrow W$ be functions. Recall that the *composition* $g \circ f$ is the function $g \circ f: X \rightarrow Z$ such that $(g \circ f)(x) = g(f(x))$ for all $x \in X$.
- Show that composition is *associative*: that $h \circ (g \circ f) = (h \circ g) \circ f$ (recall that functions are equal if they have the same value at every x).
 - f is called *injective* or *one-to-one* (1:1) if $x \neq x'$ implies $f(x) \neq f(x')$. Show that if $g \circ f$ is injective then so is f .
 - g is called *surjective* or *onto* if for every $z \in Z$ there is $y \in Y$ such that $g(y) = z$. Show that if $g \circ f$ is surjective then so is g .
 - Suppose that f, g are both surjective or both injective. In either case show that the same holds for $g \circ f$.
 - Give an example of a set X and $f, g: X \rightarrow X$ such that $f \circ g \neq g \circ f$.

The following problem is intended to help clarify the construction of $\mathbb{Z}/n\mathbb{Z}$. See also section 0.1 of Dummit and Foote.

- B. A relation R between elements of a set X is (1) *reflexive* if xRx for every $x \in X$; (2) *symmetric* if for any x, y , xRy implies yRx (3) *transitive* if for any x, y, z , if xRy and yRz then xRz . Example: $<$ on \mathbb{R} isn't reflexive ($3 < 3$ doesn't hold), isn't symmetric ($3 < 5$ holds but $5 < 3$ doesn't) by is transitive (if $x < y$ and $y < z$ then $x < z$).
- Decide which of the properties holds for the relations $=, \leq$ on \mathbb{R} .
 - Ditto where R is the relation on \mathbb{R} given by $xRy \leftrightarrow |x - y| \leq 1$ (so $\frac{1}{2}R\frac{3}{4}$ holds but $0R2$ doesn't).
 - Ditto where $aRb \leftrightarrow ab > 0$ on \mathbb{Z} and where $aRb \leftrightarrow ab \geq 0$ on \mathbb{Z} .
- DEF An *equivalence relation* is one satisfying all of (1),(2),(3).
- On the set of integers \mathbb{Z} show that the relation “ a, b are both even or both odd” is an equivalence relation (in detail: you need to show that for any a “ a, a are both even or both odd” that “if a, b are both even or both odd then also b, a are both even or both odd” and that “if a, b are both even or both odd, and if b, c are both even or both odd” then “ a, c are both even or both odd”

Supplementary Problems II: Subsemigroups of $(\mathbb{Z}_{\geq 0}, +)$

- C. The Kingdom of Ruritania mints coins in the denominations d_1, \dots, d_r Marks (d_i are positive integers, of course). Let $d = \gcd(d_1, \dots, d_r)$.
- Show that every payable sum (total value of a combination of coins) is a multiple of d Marks.
 - Show that there exists $N \geq 1$ such that any multiple of d Marks exceeding N can be expressed using the given coins.
 - Let $H \subset \mathbb{Z}_{\geq 0}$ be the set of sums that can be paid using the coins. Show that H is closed under addition.
- DEF H is called the *subsemigroup of $\mathbb{Z}_{\geq 0}$ generated by $\{d_1, \dots, d_r\}$.*

- D. (partial classification of subsemigroups of $\mathbb{Z}_{\geq 0}$) Let $H \subset \mathbb{Z}_{\geq 0}$ be closed under addition.
- (a) Show that either $H = \{0\}$ or there are $N, d \geq 1$ such that d divides every element of h , and such that H contains all multiples of d exceeding N .
Hint: Enumerate the elements of H in increasing order as $\{h_i\}_{i=1}^{\infty}$ and consider the sequence $\{\gcd(h_1, \dots, h_m)\}_{m=1}^{\infty}$.
- (b) Conclude that H is *finitely generated*: there are $\{d_1, \dots, d_r\} \subset H$ such that H is obtained as in problem C.

Supplementary Problems III: Additive groups in \mathbb{R} .

- E. (just linear algebra)
- (a) Show that the usual addition and multiplication by rational numbers endow \mathbb{R} with the structure of a vector space over the field \mathbb{Q} .
- (b) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be additive ($f(x+y) = f(x) + f(y)$). Show that f is \mathbb{Z} -linear: that $f(nx) = nf(x)$ for all $x \in \mathbb{R}, n \in \mathbb{Z}$.
- (c) Show that f is \mathbb{Q} -linear: $f(rx) = rf(x)$ for all $r \in \mathbb{Q}$.
- (d) Let $B \subset \mathbb{R}$ be a basis for \mathbb{R} as a \mathbb{Q} -vector space (this is called a *Hamel basis*). Use B to construct a \mathbb{Q} -linear map $\mathbb{R} \rightarrow \mathbb{R}$ which is not of the form $x \mapsto ax$.
- F. (add topology ...)
- Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be additive.
- (a) Suppose that f is *continuous*. Show that $f(x) = ax$ where $a = f(1)$.
- (b) (If you have taken Math 422) Suppose that f is *Lebesgue (or Borel) measurable*. Show that there is $a \in \mathbb{R}$ such that $f(x) = ax$ a.e.
- (c) (“ \mathbb{R} has no field automorphisms”) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfy $f(x+y) = f(x) + f(y)$ and $f(xy) = f(x)f(y)$. Show that either $f(x) = 0$ for all x or $f(x) = x$ for all x .