

**MATH 100 – WORKSHEET 16**  
**MINIMA AND MAXIMA**

1. ABSOLUTE MINIMA AND MAXIMA BY HAND

**Theorem.** *If  $f$  is continuous on  $[a, b]$  it has an absolute maximum and minimum there.*

(1) Find the absolute maximum and minimum values of  $f(x) = |x|$  on the interval  $[-3, 5]$ .

(2) Find the absolute maximum and minimum of  $f(x) = \sqrt{x}$  on  $[0, 5]$ .

2. LOCAL MINIMA AND DERIVATIVES

**Theorem (Fermat).** *If, in addition,  $f$  is defined and differentiable near  $c$  (on both sides!) and has a local extremum at  $c$  then  $f'(c) = 0$ .*

**Procedure**

- Call  $c$  a *critical number* if  $f'(c) = 0$ , a *singularity* if  $f'(c)$  does not exist.
  - To find absolute maximum/minimum of a continuous function  $f$  defined on  $[a, b]$ :
    - Evaluate  $f(c)$  at all critical numbers.
    - Evaluate  $f(a), f(b)$
    - Choose largest, smallest value
- (1) (Final, 2011) Let  $f(x) = 6x^{1/5} + x^{6/5}$ .
- (a) Find the critical numbers and singularities of  $f$ .

(b) Find its absolute maximum and minimum on the interval  $[-32, 32]$ .

(2) (caution)

(a) Show that  $f(x) = (x - 1)^4 + 7$  attains its absolute minimum at  $x = 1$ .

(b) Show that  $f(x) = (x - 1)^3 + 7$  has  $f'(1) = 0$  but has no local minimum or maximum there.

(3) (Midterm, 2010) Find the maximum value of  $x\sqrt{1 - \frac{3}{4}x^2}$  on the interval  $[0, 1]$ .

(4) (Final, 2007) Let  $f(x) = x\sqrt{3 - x}$ .

(a) Find the domain of  $f$ .

(b) Determine the  $x$ -coordinates of any local maxima or minima of  $f$ .