

Math 412: Advanced Linear Algebra
Lecture Notes

Lior Silberman

These are rough notes for the Fall 2016 course, compiled September 7, 2017.
Copyright Lior Silberman. These notes are available for traditional academic reuse (with attribution), and are specifically excluded from the terms of UBC Policy 81.

Contents

Introduction	4
0.1. Goals and course plan (Lecture 1)	4
0.2. Review	5
0.3. Euler's Theorem	6
Chapter 1. Constructions	7
1.1. Direct sum, direct product (Lectures 2-4)	7
1.2. Quotients (Lecture 5)	8
1.3. Hom spaces and duality (Lectures 6-8)	9
1.4. Multilinear algebra and tensor products (Lectures 9-14)	14
Chapter 2. Structure Theory: The Jordan Canonical Form	18
2.1. Introduction (Lecture 15)	18
2.2. The minimal polynomial (Lecture 16)	19
2.3. Generalized eigenspaces and Cayley–Hamilton (Lectures 17-18)	21
2.4. Nilpotent maps and Jordan blocks (Lectures 19-20)	24
2.5. The Jordan canonical form (Lecture 21)	26
Chapter 3. Vector and matrix norms	30
3.1. Norms on vector spaces (Lecture 22)	30
3.2. Norms on matrices (Lectures 23-24)	32
3.3. Example: eigenvalues and the power method (Lecture 25)	33
3.4. Sequences and series of vectors and matrices (Lectures 26-27)	34
3.5. The exponential series (Lecture 28)	36
3.6. Invertibility and the resolvent (Lecture 29)	39
3.7. Holomorphic calculus	40
Chapter 4. Vignettes	41
4.1. The exponential map and structure theory for $GL_n(\mathbb{R})$	41
4.2. Representation Theory of Groups	41
Bibliography	42

Introduction

Lior Silberman, lior@Math.UBC.CA, <http://www.math.ubc.ca/~lior/>
 Office: Math Building 229B
 Phone: 604-827-3031

For administrative details see the syllabus.

0.1. Goals and course plan (Lecture 1)

Math	Metamath	Skills
Reinforce basics <ul style="list-style-type: none"> • Vector space, subspace • Linear independence, basis • Linear map • Eigenvalue, eigenvector 	Abstraction	Hard problems Multiple ideas
New ideas <ul style="list-style-type: none"> • Direct sum, product • $\text{Hom}(U, V)$ and duality • Quotients • Tensor products 	Constructions Universal properties	Abstract examples
Structure theory for linear maps <ul style="list-style-type: none"> • Matrix decompositions • LU, LL^\dagger and Computation • Minimal poly, Cayley–Hamilton • Jordan canonical form 	Canonical forms Algorithms	
Analysis <ul style="list-style-type: none"> • Norms • Holomorphic calculus (e^{tX}) 	Combining fields	

0.2. Review

0.2.1. Basic definitions. We want to give ourselves the *freedom* to have scalars other than real or complex.

DEFINITION 1 (Fields). A *field* is a quintuple $(F, \underline{0}, 1, +, \cdot)$ such that $(F, \underline{0}, +)$ and $(F \setminus \{0\}, 1, \cdot)$ are abelian groups, and the *distributive law* $\forall x, y, z \in F : x(y + z) = xy + xz$ holds.

EXAMPLE 2. $\mathbb{R}, \mathbb{C}, \mathbb{Q}, \mathbb{F}_2$ (via addition and multiplication tables; ex: show this is a field), \mathbb{F}_p .

EXERCISE 3. Every finite field has p^r elements for some prime p and some integer $r \geq 1$. Fact: there is one such field for every prime power.

DEFINITION 4. A *vector space* over a field F is a quadruple $(V, \underline{0}, +, \cdot)$ where $(V, \underline{0}, +)$ is an abelian group, and $\cdot : F \times V \rightarrow V$ is a map such that:

- (1) $1_F \underline{v} = \underline{v}$.
- (2) $\alpha(\beta \underline{v}) = (\alpha\beta) \underline{v}$.
- (3) $(\alpha + \beta)(\underline{v} + \underline{w}) = \alpha \underline{v} + \beta \underline{v} + \alpha \underline{w} + \beta \underline{w}$.

LEMMA 5. $0_F \cdot \underline{v} = \underline{0}$ for all \underline{v} .

PROOF. $0 \underline{v} = (0 + 0) \underline{v} = 0 \underline{v} + 0 \underline{v}$. Now subtract $0 \underline{v}$ from both sides. □

0.2.2. Bases and dimension. Fix a vector space V .

DEFINITION 6. Let $S \subset V$.

- $\underline{v} \in V$ *depends on* S if there are $\{\underline{v}_i\}_{i=1}^r \subset S$ and $\{a_i\}_{i=1}^r \subset F$ such that $\underline{v} = \sum_{i=1}^r a_i \underline{v}_i$ [empty sum is $\underline{0}$]
- Write $\text{Span}_F(S) \subset V$ for the set of vectors that depend on S .
- Call S *linearly dependent* if some $\underline{v} \in S$ depends on $S \setminus \{\underline{v}\}$, equivalently if there are distinct $\{\underline{v}_i\}_{i=1}^r \subset S$ and $\{a_i\}_{i=1}^r \subset F$ not all zero such that $\sum_{i=1}^r a_i \underline{v}_i = \underline{0}$.
- Call S *linearly independent* if it is not linearly dependent.

AXIOM 7 (Axiom of choice). *Every vector space has a basis.*

0.2.3. Examples. $\{\underline{0}\}, \mathbb{R}^n, F^X$.

0.3. Euler's Theorem

Let $G = (V, E)$ be a connected planar graph. A *face* of G is a finite connected component of $\mathbb{R}^2 \setminus G$.

THEOREM 8 (Euler). $v - e + f = 1$.

PROOF. Arbitrarily orient the edges. Let $\partial_E: \mathbb{R}^E \rightarrow \mathbb{R}^V$ be defined by $f((u, v)) = 1_v - 1_u$, $\partial_F: \mathbb{R}^F \rightarrow \mathbb{R}^E$ be given by the sum of edges around the face.

LEMMA 9. ∂_F is injective.

PROOF. Faces containing boundary edges are independent. Remove them and repeat. \square

LEMMA 10. $\text{Ker } \partial_E = \text{Im } \partial_F$.

PROOF. Suppose a combo of edges is in the kernel. Following a sequence with non-zero coefficients gives a closed loop, which can be expressed as a sum of faces. Now subtract a multiple to reduce the number of edges with non-zero coefficients. \square

LEMMA 11. $\text{Im}(\partial_E)$ is the set of functions with total weight zero.

PROOF. Clearly the image is contained there. Conversely, given f of total weight zero move the weight to a single vertex using elements of the image. [remark: quotient vector spaces] \square

Now $\dim \mathbb{R}^E = \dim \text{Ker } \partial_E + \dim \text{Im } \partial_E = \dim \text{Im } \partial_F + \dim \text{Im } \partial_E$ so

$$e = f + (v - 1).$$

\square

REMARK 12. Using \mathbb{F}_2 coefficients is even simpler.

CHAPTER 1

Constructions

Fix a field F .

1.1. Direct sum, direct product (Lectures 2-4)

1.1.1. Simplest case (Lecture 2).

CONSTRUCTION 13 (External direct sum). *Let U, V be vector spaces. Their direct sum, denoted $U \oplus V$, is the vector space whose underlying set is $U \times V$, with coordinate-wise addition and scalar multiplication.*

LEMMA 14. *This really is a vector space.*

REMARK 15. The Lemma serves to review the definition of vector space.

PROOF. Every property follows from the respective properties of U, V . □

REMARK 16. Direct products of groups are discussed in 322.

LEMMA 17. $\dim_F (U \oplus V) = \dim_F U + \dim_F V$.

REMARK 18. This Lemma serves to review the notion of basis.

PROOF. Let B_U, B_V be bases of U, V respectively. Then $\{(\underline{u}, \underline{0}_V)\}_{\underline{u} \in B_U} \sqcup \{(\underline{0}_U, \underline{v})\}_{\underline{v} \in B_V}$ is a basis of $U \oplus V$. □

EXAMPLE 19. $\mathbb{R}^n \oplus \mathbb{R}^m \simeq \mathbb{R}^{n+m}$.

1.1.2. Internal sum and direct sum (Lecture 3). A key situation is when U, V are subspaces of an “ambient” vector space W .

LEMMA 20. *Let W be a vector space, $U, V \subset W$. Then $\text{Span}_F (U \cup V) = \{\underline{u} + \underline{v} \mid \underline{u} \in U, \underline{v} \in V\}$.*

PROOF. RHS contained in the span by definition. It is a subspace (non-empty, closed under addition and scalar multiplication) which contains U, V hence contains the span. □

DEFINITION 21. The space in the previous lemma is called the *sum* of U, V and denoted $U + V$.

LEMMA 22. *Let $U, V \subset W$. There is a unique homomorphism $U \oplus V \rightarrow U + V$ which is the identity on U, V .*

PROOF. Define $f((\underline{u}, \underline{v})) = \underline{u} + \underline{v}$. Check that this is a linear map. □

PROPOSITION 23 (Dimension of sums). $\dim_F (U + V) = \dim_F U + \dim_F V - \dim_F (U \cap V)$.

PROOF. Consider the map f of Lemma 22. It is surjective by Lemma 20. Moreover $\text{Ker } f = \{(\underline{u}, \underline{v}) \in U \oplus V \mid \underline{u} + \underline{v} = \underline{0}_W\}$, that is

$$\text{Ker } f = \{(\underline{w}, -\underline{w}) \mid \underline{w} \in U \cap V\} \simeq U \cap V.$$

Since $\dim_F \text{Ker } f + \dim_F \text{Im } f = \dim (U \oplus V)$ the claim now follows from Lemma 17. □

REMARK 24. This was a review of that formula. Alternative proof by starting from a basis of $U \cap V$ and extending to bases of U, V , which is basically revisiting the proof of the formula.

DEFINITION 25 (Internal direct sum). We say the sum is *direct* if f is an isomorphism.

THEOREM 26. For subspaces $U, V \subset W$ TFAE

- (1) The sum $U + V$ is direct and equals W ;
- (2) $U + V = W$ and $U \cap V = \{0\}$
- (3) Every vector $\underline{w} \in W$ can be uniquely written in the form $\underline{w} = \underline{u} + \underline{v}$.

PROOF. (1) \Rightarrow (2): $U + V = W$ by assumption, $U \cap V = \text{Ker } f$.

(2) \Rightarrow (3): the first assumption gives existence, the second uniqueness.

(3) \Rightarrow (1): existence says f is surjective, uniqueness says f is injective. □

1.1.3. Finite direct sums (Lecture 4). Three possible notions: $(U \oplus V) \oplus W$, $U \oplus (V \oplus W)$, vector space structure on $U \times V \times W$. These are *all the same*. Not just isomorphic (that is, not just same dimension), but also isomorphic when considering the extra structure of the copies of U, V, W . How do we express this?

DEFINITION 27. W is the *internal direct sum* of its subspaces $\{V_i\}_{i \in I}$ if it spanned by them and each vector has a unique representation as a sum of elements of V_i (either as a finite sum of non-zero vectors or as a zero-extended sum).

REMARK 28. This generalizes the notion of “linear independence” from vectors to subspaces.

LEMMA 29. Each of the three candidates contains an embedded copy of U, V, W and is the internal direct sum of the three images.

PROOF. Easy. □

PROPOSITION 30. Let A, B each be the internal direct sum of embedded copies of U, V, W . Then there is a unique isomorphism $A \rightarrow B$ respecting this structure.

PROOF. Construct. □

REMARK 31. (1) Proof only used result of Lemma, not specific structure; but (2) proof implicitly relies on isomorphism to $U \times V \times W$; (3) We used the fact that a map can be defined using values on copies of U, V, W (4) Exactly same proof as the facts that a function on 3d space can be defined on bases, and that that all 3d spaces are isomorphic.

- Dimension by induction.

DEFINITION 32. Abstract arbitrary direct sum.

- Block diagonality.
- Block upper-triangularity. [first structural result].

1.2. Quotients (Lecture 5)

Recall that for a group G and a normal subgroup N , we can endow the quotient G/N with group structure $(gN)(hN) = (gh)N$.

- This is well-defined, gives group.
- Have quotient map $q: G \rightarrow G/N$ given by $g \mapsto gN$.

- Homomorphism theorem: any $f: G \rightarrow H$ factors as $G \rightarrow G/\text{Ker}(f)$ follows by isomorphism.
- If $N < M < G$ with both N, M normal then $q(M) \simeq M/N$ is normal in G/N and $(G/N)/(M/N) \simeq (G/M)$.

Now do the same for vector spaces.

LEMMA 33. Let V be a vector space, W a subspace. Let $\pi: V \rightarrow V/W$ be the quotient as abelian groups. Then there is a unique vector space structure on V/W making π a surjective linear map.

PROOF. We must set $\alpha(\underline{v} + W) = \alpha\underline{v} + W$. This is well-defined and gives the isomorphism. Use quotient map to verify vector space axioms. \square

EXAMPLE 34. $V = C^1(0, 1)$, space of continuously differentiable functions. Then $\frac{d}{dx}: V \rightarrow C(0, 1)$ vanishes on $\mathbb{R}\mathbb{1}$ and hence induces a map $\frac{d}{dx}: (C^1(0, 1)/\mathbb{R}\mathbb{1}) \rightarrow C(0, 1)$. Note that the inverse of this map is what we call “indefinite integral” – whose images is exactly an equivalence class “function+c”.

EXAMPLE 35. The definite integral is a linear function which vanishes on functions which are non-zero at countably many points. More generally, integration works on functions modulo functions which are zero a.e.

FACT 36. The properties above persist for vector spaces.

- How to use quotients: “kill off” part of the vector spaces that is irrelevant (linear maps vanish there) or already understood.

1.3. Hom spaces and duality (Lectures 6-8)

1.3.1. Hom spaces (Lecture 5 continued or start of lecture 6).

DEFINITION 37. $\text{Hom}_F(U, V)$ will denote the space of F -linear maps $U \rightarrow V$.

LEMMA 38. $\text{Hom}_F(U, V) \subset V^U$ is a subspace, hence a vector space.

DEFINITION 39. $V' = \text{Hom}_F(V, F)$ is called the dual space.

Motivation 1: in PDE. Want solutions in some function space V . Use that V' is much bigger to find solutions in V' , then show they are represented by functions.

1.3.2. The dual space, finite dimensions.

NOTE 40. In lecture ignore infinite dimensions (but make statements which are correct in general).

CONSTRUCTION 41 (Dual basis). Let $B = \{\underline{b}_i\}_{i \in I} \subset V$ be a basis. Write $\underline{v} \in V$ uniquely as $\underline{v} = \sum_{i \in I} a_i \underline{b}_i$ (almost all $a_i = 0$) and set $\varphi_i(\underline{v}) = a_i$.

LEMMA 42. These are linear functionals.

PROOF. Represent $\alpha\underline{v} + \underline{v}'$ in the basis. \square

EXAMPLE 43. $V = F^n$ with standard basis, get $\varphi_i(\underline{x}) = x_i$. Note every functional has the form $\varphi(\underline{x}) = \sum_{i=1}^n \varphi(\underline{e}_i) \varphi_i(\underline{x})$.

REMARK 44. Alternative construction: φ_i is the unique linear map to F satisfying $\varphi_i(\underline{b}_j) = \delta_{i,j}$.

LEMMA 45. *The dual basis is linearly independent. It is spanning iff $\dim_F V < \infty$.*

PROOF. Evaluate a linear combination at \underline{b}_j .

If V is finite-dimensional, enumerate the basis as $\{\underline{b}_i\}_{i=1}^n$. Then for any $\varphi \in V'$ and any $\underline{v} \in V$ write $\underline{v} = \sum_i a_i \underline{b}_i$ and then

$$\varphi(\underline{v}) = \sum_i a_i \varphi(\underline{b}_i) = \sum_i (\varphi(\underline{b}_i)) \varphi_i(\underline{v})$$

so

$$\varphi = \sum_i (\varphi(\underline{b}_i)) \varphi_i \in \text{Span}_F \{\varphi_i\}.$$

In the infinite-dimensional case let $\phi = \sum_{i \in I} \varphi_i$. Then ϕ is a well-defined linear functional which depends on every coordinate hence not in the span of the $\{\varphi_i\}$. \square

REMARK 46. This isomorphism $V \rightarrow V'$ is not canonical: the functional φ_i depends on the whole basis B and not only on \underline{b}_i , and the dual basis transforms differently from the original basis under change-of-basis.

The argument above used evaluation – let’s investigate that more.

PROPOSITION 47 (Double dual). *Given $\underline{v} \in V$ consider the evaluation map $e_{\underline{v}}: V' \rightarrow F$ given by $e_{\underline{v}}(\varphi) = \varphi(\underline{v})$. Then $\underline{v} \mapsto e_{\underline{v}}$ is a linear injection $V \hookrightarrow V''$, an isomorphism iff V is finite-dimensional.*

PROOF. The vector space structure on V' (and on F^V in general) is such that $e_{\underline{v}}$ is linear. That the map $\underline{v} \mapsto e_{\underline{v}}$ is linear follows from the linearity of the elements of V' . For injectivity let $\underline{v} \in V$ be non-zero. Extending \underline{v} to a basis, let $\varphi_{\underline{v}}$ be the element of the dual such that $\varphi_{\underline{v}}(\underline{v}) = 1$. Then $e_{\underline{v}}(\varphi_{\underline{v}}) \neq 0$ so $e_{\underline{v}} \neq 0$. If $\dim_F V = n$ then $\dim_F V' = n$ and thus $\dim_F V'' = n$ and we have an isomorphism. \square

The map $V \hookrightarrow V''$ is natural: the image $e_{\underline{v}}$ of \underline{v} is intrinsic and does not depend on a choice of basis.

1.3.3. The dual space, infinite dimensions (Lecture 7).

LEMMA 48 (Interaction with past constructions). *We have*

- (1) $(V/U)' \hookrightarrow V'$ as $\{\varphi \in V' \mid \varphi(U) = \{0\}\}$.
- (2) $(U \oplus V)' \simeq U' \oplus V'$.

PROOF. Universal property. \square

COROLLARY 49. *Since $(F)' \simeq F$, it follows by induction that $(F^n)' \simeq F^n$.*

What about infinite sums?

- The universal property gives a bijection $(\bigoplus_{i \in I} V_i)' \xleftrightarrow{\sim} \prod_{i \in I} V_i'$, more generally

$$\text{Hom}_F \left(\bigoplus_{i \in I} V_i, Z \right) \xleftrightarrow{1:1} \prod_{i \in I} \text{Hom}_F(V_i, Z).$$

- LHS has a vector space structure – should get one on the right.
- Which leads us to observe that:

- Any Cartesian product $\times_{i \in I} W_i$ has a natural vector space structure, coming from point-wise addition and scalar multiplication.
 - Note that the underlying set is

$$\begin{aligned} \times_{i \in I} W_i &= \{f \mid f \text{ is a function with domain } I \text{ and } \forall i \in I : f(i) \in W_i\} \\ &= \left\{ f : I \rightarrow \bigcup_{i \in I} W_i \mid f(i) \in W_i \right\}. \end{aligned}$$

* RMK: AC means Cartesian products nonempty, but our sets have a distinguished element so this is not an issue.

- Define $\alpha(w_i)_{i \in I} + (w'_i)_{i \in I} \stackrel{\text{def}}{=} (\alpha w_i + w'_i)_{i \in I}$. This gives a vector space structure.
- Denote the resulting vector space $\prod_{i \in I} W_i$ and called it the *direct product* of the W_i .
- The bijection $(\bigoplus_{i \in I} V_i)' \longleftrightarrow \prod_{i \in I} V_i'$ is now a linear isomorphism [in fact, the vector space structure on the right is the one transported by the isomorphism].

We now investigate $\prod_i W_i$ in general.

- Note that it contains a copy of each W_i (map $\underline{w} \in W_i$ to the sequence which has \underline{w} in the i th position, and $\underline{0}$ at every other position).
- And these copies are linearly independent: if a sum of such vectors from distinct W_i is zero, then every coordinate was zero.
- Thus $\prod_i W_i$ contains $\bigoplus_{i \in I} W_i$ as an internal direct sum.
 - This subspace is exactly the subset $\{\underline{w} \in \prod_i W_i \mid \text{supp}(\underline{w}) \text{ is finite}\}$.
 - And in fact, that subspace proves that $\bigoplus_{i \in I} W_i$ exists.
 - But $\prod_i W_i$ contains many other vectors – it is much bigger.

EXAMPLE 50. $\mathbb{R}^{\oplus \mathbb{N}} \subset \mathbb{R}^{\mathbb{N}}$ – and the latter is the dual!

COROLLARY 51. *The dual of an infinite-dimensional space is much bigger than the sum of the duals, and the double dual is bigger yet.*

1.3.4. Question: so we only have *finite* sums in linear algebra. What about infinite sums?

Answer: no infinite sums in algebra. Definition of $\sum_{n=1}^{\infty} a_n = A$ from real analysis relies on *analytic properties* of A (close to partial sums), not algebraic properties.

But, calculating sums can be understood in terms of linear functionals.

LEMMA 52 (Results from Calc II, reinterpreted). *Let $S \subset \mathbb{R}^{\mathbb{N}}$ denote the set of sequences \underline{a} such that $\sum_{n=1}^{\infty} a_n$ converges.*

- (1) $\mathbb{R}^{\oplus \mathbb{N}} \subset S \subset \mathbb{R}^{\mathbb{N}}$ is a linear subspace.
- (2) $\Sigma : S \rightarrow \mathbb{R}$ given by $\Sigma(\underline{a}) = \sum_{n=1}^{\infty} a_n$ is a linear functional.

Philosophy: Calc I,II made element-by-element statements, but using linear algebra we can express them as statements on the whole space.

Now questions about summing are questions about *intelligently* extending the linear functional Σ to a bigger subspace. BUT: if an extension is to satisfy every property of summing series, it is actually the trivial (no) extension.

For more information let's talk about limits of sequences instead (once we can generalize limits just apply that to partial sums of a series).

DEFINITION 53. Let $c \subset \ell^\infty \subset \mathbb{R}^\mathbb{N}$ be the sets of convergent, respectively bounded sequences.

LEMMA 54. $c \subset \ell^\infty$ are subspaces, and $\lim_{n \rightarrow \infty}: c \rightarrow \mathbb{R}$ is a linear functional.

EXAMPLE 55. Let $C: \mathbb{R}^\mathbb{N} \rightarrow \mathbb{R}^\mathbb{N}$ be the Cesàro map $(C\underline{a})_N = \frac{1}{N} \sum_{n=1}^N a_n$. This is clearly linear. Let $CS = C^{-1}(c)$ be the set of sequences which are Cesàro-convergent, and set $L \in CS'$ by $L(\underline{a}) = \lim_{n \rightarrow \infty}(C\underline{a})$. This is clearly linear (composition of linear maps). For example, the sequence $(0, 1, 0, 1, \dots)$ now has the limit $\frac{1}{2}$.

LEMMA 56. If $\underline{a} \in c$ then $C\underline{a} \in c$ and they have the same limit. Thus L above is an extension of $\lim_{n \rightarrow \infty}$.

THEOREM 57. There are two functionals $LIM, \lim_\omega \in (\ell^\infty)'$ (“Banach limit”, “limit along ultrafilter”, respectively) such that:

- (1) They are positive (map non-negative sequences to non-negative sequences);
- (2) Agree with $\lim_{n \rightarrow \infty}$ on c ;
- (3) And, in addition
 - (a) $LIM \circ S = LIM$ where $S: \ell^\infty \rightarrow \ell^\infty$ is the shift.
 - (b) $\lim_\omega (a_n b_n) = (\lim_\omega a_n) (\lim_\omega b_n)$.

1.3.5. Pairings and bilinear forms (Lecture 8). Goal: identify the dual of a vector space in concrete terms. For this we need an abstract notion of dual not tied to the particular realization V' (similar to how we have an abstract notion of direct sum as “space generated by independent copies of V_i ” which is not tied to the concrete realization as the subspace of tuples of finite support in $\prod_i V_i$).

OBSERVATION 58. The evaluation map $V \times V' \rightarrow F$ given by

$$\langle \underline{v}, \varphi \rangle = \varphi(\underline{v})$$

is *bilinear* (=linear in each variable).

Note that linearity in the first variable is equivalent to the linearity of φ , while linearity in the second variable is equivalent to the definition of the vector space structure on V' .

DEFINITION 59 (Pairings / bilinear maps). For any two vector spaces U, V a (bilinear) *pairing* between U, V is a map

$$\langle \cdot, \cdot \rangle: U \times V \rightarrow F$$

which is linear in each variable separately. Similarly we define a *bilinear map* $U \times V \rightarrow Z$.

EXAMPLE 60. The standard inner product on F^n : $\langle \underline{u}, \underline{v} \rangle = \sum_i u_i v_i$. More generally, given $B \in M_{m \times n}(F)$ have a pairing on $F^m \times F^n$ given by

$$\langle \underline{u}, \underline{v} \rangle = \sum_i u_i B_{ij} v_j.$$

More generally, given any bilinear pairing of U, V choose bases $\{\underline{u}_i\}_{i \in I} \subset U, \{\underline{v}_j\}_{j \in J} \subset V$ and define the *Gram matrix* by

$$B_{ij} = \langle \underline{u}_i, \underline{v}_j \rangle.$$

We can then compute the pairing of any two vectors: by the distributive law (“FOIL”)

$$\left\langle \sum_i a_i \underline{u}_i, \sum_j b_j \underline{v}_j \right\rangle = \sum_{i,j} a_j B_{ij} b_j.$$

Conversely, any matrix B defines a bilinear pairing (aside: this is a linear bijection if you give pairings the obvious vector space structure).

1.3.6. Pairings: duality and degeneracy. Fix a bilinear form $\langle \cdot, \cdot \rangle : U \times V \rightarrow F$. Then for any $\underline{u} \in U$ we get a map $\varphi_{\underline{u}} : V \rightarrow F$ by $\varphi_{\underline{u}}(\underline{v}) = \langle \underline{u}, \underline{v} \rangle$.

(1) $\varphi_{\underline{u}}$ is linear ($\in V'$) iff the pairing is linear in the second variable.

(2) The map $U \rightarrow V'$ given by $\underline{u} \rightarrow \varphi_{\underline{u}}$ is linear iff the pairing is linear in the first variable.

We conclude that every pairing gives a map $U \rightarrow V'$, and equivalently also a map $V \rightarrow U'$.

LEMMA 61. We have a linear bijection $\{\text{pairings on } U \times V\} \longleftrightarrow \text{Hom}_F(U, V')$

PROOF. The inverse map associates to each $f \in \text{Hom}_F(U, V')$ the bilinear form

$$\langle \underline{u}, \underline{v} \rangle_f = (f(\underline{u}))(\underline{v}).$$

□

DEFINITION 62. Call the bilinear map *non-degenerate* if both maps $U \rightarrow V'$, $V \rightarrow U'$ are embeddings.

LEMMA 63. A pairing is non-degenerate iff for every non-zero $\underline{u} \in U$ there is $\underline{v} \in V$ such that $\langle \underline{u}, \underline{v} \rangle \neq 0$ and conversely.

Key idea: if the map $V \rightarrow U'$ associated to a pairing is bijective, then we can use V as a *model* for U' via the pairing.

EXAMPLE 64. The dot product is a non-degenerate pairing $F^n \times F^n$ hence identifies $(F^n)'$ with F^n .

Two further examples from functional analysis:

First we fix a compact topological space X . Then for any finite Borel measure μ on X and any continuous $f \in C(X)$ we have the *integral*

$$\int f d\mu.$$

This is a bilinear pairing $C(X) \times \{\text{finite signed measures on } X\}$ which is non-degenerate.

THEOREM 65 (Riesz representation theorem). *Let X be compact. Then every continuous linear functional on $C(X)$ is given by a finite measure (that is, the continuous dual $C(X)'$ can be represented by the space of measures).*

Second, the inner product on Hilbert space is a non-degenerate pairing!

THEOREM 66 (Riesz representation theorem). *Let \mathcal{H} be a Hilbert space. Then every continuous linear functional on \mathcal{H} is of the form $\langle \underline{u}, \cdot \rangle$.*

1.3.7. The dual of a linear map (Lecture 8, continued).

CONSTRUCTION 67. Let $T \in \text{Hom}(U, V)$. Set $T' \in \text{Hom}(V', U')$ by $(T' \varphi)(\underline{v}) = \varphi(T\underline{v})$.

LEMMA 68. This is a linear map $\text{Hom}(U, V) \rightarrow \text{Hom}(V', U')$. An isomorphism if U, V finite-dimensional.

LEMMA 69. $(TS)' = S'T'$

PROOF. PS3

□

1.4. Multilinear algebra and tensor products (Lectures 9-14)

1.4.1. Bilinear forms (Lecture 9).

DEFINITION 70. Let $\{V_i\}_{i \in I}$ be vector spaces, W another vector space. A function $f: \times_{i \in I} V_i \rightarrow W$ is said to be *multilinear* if it is linear in each variable.

EXAMPLE 71 (Bilinear maps). (1) $f(x, y) = xy$ is bilinear $F^2 \rightarrow F$.

(2) The map $(T, \underline{v}) \mapsto T\underline{v}$ is a multilinear map $\text{Hom}(V, W) \times V \rightarrow W$.

(3) For a matrix $A \in M_{n,m}(F)$ have $(\underline{x}, \underline{y}) \mapsto {}^t \underline{x} A \underline{y}$ on $F^n \times F^m$.

(4) For $\varphi \in U'$, $\psi \in V'$ have $(\underline{u}, \underline{v}) \mapsto \varphi(\underline{u})\psi(\underline{v})$, and finite combinations of those.

REMARK 72. A *bilinear function* on $U \times V$ is not the same as a *linear function on $U \oplus V$* . For example: is $f(a\underline{u}, a\underline{v})$ equal to $a f(\underline{u}, \underline{v})$ or to $a^2 f(\underline{u}, \underline{v})$? That said, $\bigoplus V_i$ was *universal* for maps from V_i . It would be nice to have a space which is universal for multilinear maps. We only discuss the finite case.

EXAMPLE 73. A multilinear function $B: U \times \{0\} \rightarrow F$ has $B(\underline{u}, \underline{0}) = B(\underline{u}, \underline{0} \cdot \underline{0}) = \underline{0} \cdot B(\underline{u}, \underline{0}) = 0$. A multilinear function $B: U \times F \rightarrow F$ has $B(\underline{u}, x) = B(\underline{u}, x \cdot 1) = x B(\underline{u}, 1) = x \varphi(\underline{u})$ where $\varphi(\underline{u}) = B(\underline{u}, 1) \in U'$.

We can reduce everything to Example 71(3): Fix bases $\{\underline{u}_i\}, \{\underline{v}_j\}$. Then

$$B \left(\sum_i x_i \underline{u}_i, \sum_j y_j \underline{v}_j \right) = \sum_{i,j} x_i B(\underline{u}_i, \underline{v}_j) y_j = {}^t \underline{x} B \underline{y}$$

where $B_{ij} = B(\underline{u}_i, \underline{v}_j)$. Note: $x_i = \varphi_i(\underline{u})$ where $\{\varphi_i\}$ is the dual basis. Conclude that

$$(1.4.1) \quad B = \sum_{i,j} B(\underline{u}_i, \underline{v}_j) \varphi_i \psi_j.$$

Easy to check that this is an expansion in a basis (check against $(\underline{u}_i, \underline{v}_j)$). We have shown:

PROPOSITION 74. *The set $\{\varphi_i \psi_j\}_{i,j}$ is a basis of the space of bilinear forms $U \times V \rightarrow F$.*

COROLLARY 75. *The space of bilinear forms on $U \times V$ has dimension $\dim_F U \cdot \dim_F V$.*

REMARK 76. Also works in infinite dimensions, since can have the sum (1.4.1) be infinite – every pair of vectors only has finite support in the respective bases.

1.4.2. The tensor product (Lecture 10-11). Now let's fix U, V and try to construct a space that will classify bilinear maps on $U \times V$.

- Our space will be generated by terms $\underline{u} \otimes \underline{v}$ on which we can evaluate f to get $f(\underline{u}, \underline{v})$.
- Since f is multilinear, $f(a\underline{u}, b\underline{v}) = ab f(\underline{u}, \underline{v})$ so need $(a\underline{u}) \otimes (b\underline{v}) = ab(\underline{u} \otimes \underline{v})$.
- Similarly, since $f(\underline{u}_1 + \underline{u}_2, \underline{v}) = f(\underline{u}_1, \underline{v}) + f(\underline{u}_2, \underline{v})$ want $(\underline{u}_1 + \underline{u}_2) \otimes (\underline{v}_1 + \underline{v}_2) = \underline{u}_1 \otimes \underline{v}_1 + \underline{u}_2 \otimes \underline{v}_1 + \underline{u}_1 \otimes \underline{v}_2 + \underline{u}_2 \otimes \underline{v}_2$.

CONSTRUCTION 77 (Tensor product). *Let U, V be spaces. Let $X = F^{\oplus(U \times V)}$ be the formal span of all expressions of the form $\{\underline{u} \otimes \underline{v}\}_{(\underline{u}, \underline{v}) \in U \times V}$. Let $Y \subset X$ be the subspace spanned by*

$$\{(a\underline{u}) \otimes (b\underline{v}) - ab(\underline{u} \otimes \underline{v}) \mid a, b \in F, (\underline{u}, \underline{v}) \in U \times V\}$$

and

$$\{(\underline{u}_1 + \underline{u}_2) \otimes (\underline{v}_1 + \underline{v}_2) - (\underline{u}_1 \otimes \underline{v}_1 + \underline{u}_2 \otimes \underline{v}_1 + \underline{u}_1 \otimes \underline{v}_2 + \underline{u}_2 \otimes \underline{v}_2) \mid **\}.$$

Then set $U \otimes V = X/Y$ and let $\iota: U \times V \rightarrow U \otimes V$ be the map $\iota(\underline{u}, \underline{v}) = (\underline{u} \otimes \underline{v}) + Y$.

THEOREM 78. ι is a bilinear map. For any space W any any bilinear map $f: U \times V \rightarrow W$, there is a unique linear map $\tilde{f}: U \otimes V \rightarrow W$ such that $f = \tilde{f} \circ \iota$.

PROOF. Uniqueness is clear, since $\tilde{f}(\underline{u} \otimes \underline{v}) = f(\underline{u}, \underline{v})$ fixes \tilde{f} on a generating set. For existence we need to show that if $\tilde{f}: X \rightarrow W$ is defined by $\tilde{f}(\underline{u} \otimes \underline{v}) = f(\underline{u}, \underline{v})$ then \tilde{f} vanishes on Y and hence descends to $U \otimes V$. \square

PROPOSITION 79. Let B_U, B_V be bases for U, V respectively. Then $\{\underline{u} \otimes \underline{v} \mid \underline{u} \in B_U, \underline{v} \in B_V\}$ is a basis for $U \otimes V$.

PROOF. Spanning: use bilinearity of ι . Independence: for $\underline{u} \in B_U$ let $\{\varphi_{\underline{u}}\}_{\underline{u} \in B_U} \subset U'$, $\{\psi_{\underline{v}}\}_{\underline{v} \in B_V} \subset V'$ be the dual bases. Then $\varphi_{\underline{u}} \psi_{\underline{v}}$ is a bilinear map $U \times V \rightarrow F$, and the sets $\{\underline{u} \otimes \underline{v}\}_{(\underline{u}, \underline{v}) \in B_U \times B_V}$ and $\{\widehat{\varphi_{\underline{u}} \psi_{\underline{v}}}\}_{(\underline{u}, \underline{v}) \in B_U \times B_V}$ are dual bases. \square

COROLLARY 80. $\dim_F(U \otimes V) = \dim_F U \cdot \dim_F V$.

EXAMPLE 81. Examples of tensor products

- (1) $\mathbb{R}^n \otimes \mathbb{R}^m$ encoded as matrices.
- (2) Polynomial algebra: $F[x] \otimes F[y] \simeq F[x, y]$
- (3) Functions on product spaces: let X, Y be compact then $C(X) \otimes C(Y)$ dense in $C(X \times Y)$.
 - Note Fubini's Theorem can be obtained from this.
- (4) Quantum mechanics: state space for two particles is the (completion of the) tensor product of the state spaces for the individual particles.
 - Note that many states are not "pure tensors".

1.4.3. The Universal Property (Lecture 12). Abstract view of tensor products (note the indefinite article!)

DEFINITION 82 (Abstract tensor product). A *tensor product* of the spaces U, V is a pair (W, ι) where W is a vector space, $\iota: U \times V \rightarrow W$ is bilinear, and for every bilinear map $f: U \times V \rightarrow Z$ there is a unique $\tilde{f} \in \text{Hom}_F(W, Z)$ such that $f = \tilde{f} \circ \iota$.

REMARK 83. Informally, ι is the "most general bilinear map on $U \times V$ ".

EXAMPLE 84. We show that the image of ι spans W . Indeed if not there would be a non-zero functional $\tilde{f} \in W'$ vanishing on the span of that image, and then $\tilde{f} \circ \iota = 0 \circ \iota$ would both be the zero bilinear form, violating uniqueness.

Before we explain how to use this property, we return to the example of direct sum.

DEFINITION 85 (Abstract direct sum). A *direct sum* of $\{V_i\}_{i \in I}$ is a space W and maps $e_i \in \text{Hom}_F(V_i, W)$ such that for every system of maps $f_i \in \text{Hom}_F(V_i, Z)$ there is a unique $\tilde{f}: W \rightarrow Z$ such that $f_i = \tilde{f} \circ e_i$ for each i .

PROPOSITION 86. *Direct sums are unique up to a unique isomorphism.*

PROOF. Suppose W' is another direct sum with system of inclusions $\{e'_i\}_{i \in I}$. Then W' is a space with system of maps, so by hypothesis there is a unique $\tilde{f}': W' \rightarrow W'$ such that $e'_i = \tilde{f}' \circ e_i$ (note that this is the key requirement from an isomorphism preserving the structure, so we see both that such a map exists and that it is unique, but we don't know it is an isomorphism yet).

By symmetry there is also $\bar{f}: W' \rightarrow W$ such that $e_i = \bar{f} \circ e'_i$.
 Next, note that $\bar{f} \circ \bar{f}'$ and id_W are both maps $W \rightarrow W$ and they satisfy

$$\begin{aligned} (\bar{f} \circ \bar{f}') \circ e_i &= \bar{f} \circ (\bar{f}' \circ e_i) \\ &= \bar{f} \circ e'_i && \text{choice of } \bar{f}' \\ &= e_i && \text{choice of } \bar{f} \\ &= \text{id}_W \circ e_i \end{aligned}$$

so by the universal property (applied to the system of maps e_i with target W), $\bar{f} \circ \bar{f}' = \text{id}_W$. By symmetry we also have $\bar{f}' \circ \bar{f} = \text{id}_{W'}$ and we are done. \square

Consider now direct products. Reversing all the arrows (on the board, modify the definition in red ink), we get:

DEFINITION 87 (Abstract direct product). A *direct product* of $\{V_i\}_{i \in I}$ is a space W and maps $\pi_i \in \text{Hom}_F(W, V_i)$ such that for every system of maps $f_i \in \text{Hom}_F(Z, V_i)$ there is a unique $\bar{f}: Z \rightarrow W$ such that $f_i = \pi_i \circ \bar{f}$ for each i .

PROPOSITION 88. *Direct products are unique up to a unique isomorphism.*

PROOF. Reverse all the arrows in the previous proof. \square

PROPOSITION 89. *Tensor products are unique up to a unique isomorphism.*

PROOF. Let $(W, \iota), (W', \iota')$ be two tensor products of U, V . Then since $\iota': U \times V \rightarrow W'$ is bilinear there is a unique $\bar{f}': W \rightarrow W'$ such that $\iota' = \bar{f}' \circ \iota$.

REMARK. Note that this is a basic requirement for an “isomorphism of tensor products”: it must identify the vector representing $\underline{u} \otimes \underline{v}$ on both sides. Using the universal property we saw both that we can actually identify these vectors and that this identification extends to a linear map of the tensor product spaces.

Continuing with the proof, for the same reason there is $\bar{f}: W' \rightarrow W$ such that $\iota = \bar{f} \circ \iota'$.

REMARK. One can now finish the proof by noting, for example, that $\bar{f} \circ \bar{f}'$ fixes elements of the form $\underline{u} \otimes \underline{v}$ and that these span the tensor product. But we prefer a proof which doesn’t “look under the hood” and use the vectors in the vector space.

Finally, we have

$$\begin{aligned} (\bar{f} \circ \bar{f}') \circ \iota &= \bar{f} \circ (\bar{f}' \circ \iota) \\ &= \bar{f} \circ \iota' && \text{choice of } \bar{f}' \\ &= \iota && \text{choice of } \bar{f} \\ &= \text{id}_W \circ \iota \end{aligned}$$

We have shown that the bilinear map $\iota: U \times V \rightarrow W$ is represented by both homomorphisms $\bar{f} \circ \bar{f}'$ and id_W so they must be equal, and by symmetry we conclude that $\bar{f}' \circ \bar{f} = \text{id}_{W'}$ as well so that \bar{f}, \bar{f}' are the desired isomorphisms. \square

REMARK 90. This point of view leads to “category theory” where one forgets about the specific algebraic structure under consideration (here vector spaces) and considers only the statements about objects and homomorphisms. This way theorems about “direct sums”, say, apply for any construction of direct sum regardless of the underlying algebraic structures.

For example, we get direct sums (and direct products) of groups, rings, modules, vector spaces. But we also get direct sums of topological spaces (this turns out to be the disjoint union) and direct products of topological spaces (this is the Tychonoff product).

1.4.4. Symmetric and antisymmetric tensor products (Lecture 13). For motivation, think of $U \otimes U$ as the state space of a pair of identical quantum particles. What happens when we swap them? Represent swapping them by the obvious map $T \in \text{End}_F(U \otimes U)$.

FACT 91. *Some fundamental particles (“Bosons”) always have states in the +1-eigenspace. Other particles (“Fermions”) always have states in the -1-eigenspace. Note that a state like $\underline{u} \otimes \underline{u}$ is permitted to Bosons, but prohibited (!) to Fermions. This is called the “Fermi exclusion principle”.*

ASSUMPTION 92. *For this section, $\text{char}(F) = 0$.*

Let $(12) \in S_2$ act on $V \otimes V$ by exchanging the factors (why is this well-defined?).

LEMMA 93. *Let $T \in \text{End}_F(U)$ satisfy $T^2 = \text{Id}$. Then U is the direct sum of the two eigenspaces.*

DEFINITION 94. $\text{Sym}^2 V$ and $\wedge^2 V$ are the eigenspaces.

PROPOSITION 95. *Generating sets and bases.*

In general, let S_k act on $V^{\otimes k}$.

- What do we mean by that? Well, this classifies n -linear maps $V \times \cdots \times V \rightarrow Z$. Universal property gives isom of $(U \otimes V) \otimes W, U \otimes (V \otimes W)$.
- Why action well-defined? After all, the set of pure tensors is nonlinear. So see first as multilinear map $V^n \rightarrow V^{\otimes n}$.
- Single out $\text{Sym}^k V, \wedge^k V$. Note that there are other representations.
- Claim: bases

1.4.5. Bases of Sym^k, \wedge^k , determinants (Lecture 14).

EXAMPLE 96. $\wedge^2 \mathbb{R}^3$ is three-dimensional, which explains the cross-product as giving anti-symmetric 2-tensors rather than vectors.

PROPOSITION 97. *Symmetric/antisymmetric tensors are generating sets; bases coming from subsets of basis.*

Tool: the maps $P_k^\pm : V^{\otimes k} \rightarrow V^{\otimes k}$ given by $P_k^\pm(\underline{v}_1 \otimes \cdots \otimes \underline{v}_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (\pm)^\sigma (\underline{v}_{\sigma(1)} \otimes \cdots \otimes \underline{v}_{\sigma(k)})$.

LEMMA 98. *These are well defined (extensions of linear maps). Fix elements of $\text{Sym}^k V, \wedge^k V$ respectively, images are in those subspaces (check $\tau \circ P_k^\pm = (\pm)^\tau P_k^\pm$). Conclude that image is spanned by image of basis.*

EXAMPLE 99. Exterior forms of top degree and determinants.

CHAPTER 2

Structure Theory: The Jordan Canonical Form

2.1. Introduction (Lecture 15)

2.1.1. The two paradigmatic problems. Fix a vector space V of dimension $n < \infty$ (in this chapter, all spaces are finite-dimensional unless stated otherwise), and a map $T \in \text{End}(V)$. We will try for two kinds of structural results:

- (1) [“decomposition”] $T = RS$ where $R, S \in \text{End}(V)$ are “simple”
- (2) [“canonical form”] There is a basis $\{\underline{v}_i\}_{i=1}^n \subset V$ in which the matrix of T is “simple”.

EXAMPLE 100. (From 1st course)

- (1) (Gaussian elimination) Every matrix $A \in M_n(F)$ can be written in the form $A = E_1 \cdots E_k \cdot A_{\text{rr}}$ where E_i are “elementary” (row operations or rescaling) and A_{rr} is row-reduced.
- (2) (Spectral theory) Suppose T is diagonalizable. Then there is a basis in which T is diagonal.

As an example of how to use (1), suppose $\det(A)$ is defined for matrices by column expansion. Then can show (Lemma 1) that $\det(EX) = \det(E)\det(X)$ whenever E is elementary and that (Lemma 2) $\det(A_{\text{rr}}X) = \det(A)\det(X)$ whenever A is row-reduced. One can then prove

THEOREM 101. For all A, B , $\det(AB) = \det(A)\det(B)$.

PROOF. Let $\mathcal{D} = \{A \mid \forall X : \det(AX) = \det(A)\det(X)\}$. Then we know that all $A_{\text{rr}} \in \mathcal{D}$ and that if $A \in \mathcal{D}$ then for any elementary E , $\det((EA)X) = \det(E(AX)) = \det(E)\det(AX) = \det(E)\det(A)\det(X) = \det(EA)\det(X)$ so $EA \in \mathcal{D}$ as well. It now follows from Gauss’s Theorem that \mathcal{D} is the set of all matrices. \square

2.1.2. Triangular matrices.

DEFINITION 102. $A \in M_n(F)$ is upper (lower) triangular if ...

Significance: these are very good for computation. For example:

LEMMA 103. The lower-triangular matrix L is invertible iff its diagonal entries are non-zero.

The proof is:

ALGORITHM 104 (Forward-substitution). Let L be lower-triangular with non-zero diagonal entries. Then the solution to $L\underline{x} = \underline{b}$ is given by $x_i = \frac{b_i - \sum_{j=1}^{i-1} l_{ij}x_j}{l_{ii}}$ for $i = 1, 2, \dots, n$.

REMARK 105. Note that the algorithm does exactly as many multiplications as non-zero entries in U . Hence better than Gaussian elimination for general matrix ($O(n^3)$), really good for sparse matrix, and doesn’t require storing the matrix entries only the way to calculate u_{ij} (in particular no need to find inverse).

EXERCISE 106. (1) Express this as a formula for the inverse of a lower-triangular matrix (2) Develop the backward-substitution algorithm for upper-triangular matrices and find a formula for their inverses.

COROLLARY 107. *If $A = LU$ we can efficiently solve $A\underline{x} = \underline{b}$.*

Note that we don't like to store inverses. For example, because they are generally dense matrices even if L, U are sparse.

We now try to look for a vector-space interpretation of being triangular. For this note that if $U \in M_n(F)$ is upper-triangular then

$$\begin{aligned} U\underline{e}_1 &= u_{11}\underline{e}_1 \in \text{Span}\{\underline{e}_1\} \\ U\underline{e}_2 &= u_{12}\underline{e}_1 + u_{22}\underline{e}_2 \in \text{Span}\{\underline{e}_1, \underline{e}_2\} \\ &\vdots = \vdots \\ U\underline{e}_k &\in \text{Span}\{\underline{e}_1, \dots, \underline{e}_k\} \\ &\vdots = \vdots \end{aligned}$$

In particular, we found a family of subspaces $V_i = \text{Span}\{\underline{e}_1, \dots, \underline{e}_i\}$ such that $U(V_i) \subset V_i$, such that $\{\underline{0}\} = V_0 \subset V_1 \subset \dots \subset V_n = F^n$ and such that $\dim V_i = i$.

EXERCISE 108 (Cholesky decomposition). If A is positive-definite, then $A = LL^t$ for a lower-triangular matrix L .

THEOREM 109. *$T \in \text{End}(V)$ has an upper-triangular matrix wrt some basis iff there are T -invariant subspaces $\{\underline{0}\} = V_0 \subset V_1 \subset \dots \subset V_n = F^n$ with $\dim V_i = i$.*

PROOF. We just saw necessity. For sufficiency, given V_i choose for $1 \leq i \leq n$, $\underline{v}_i \in V_i \setminus V_{i-1}$. These exist (the dimension increases by 1), are a linearly independent set (each vector is independent of its predecessors) and the first i span V_i (by dimension count). Finally for each i , $T\underline{v}_i \in T(V_i) \subset V_i = \text{Span}\{\underline{v}_1, \dots, \underline{v}_i\}$ so the matrix of T in this basis is upper triangular. \square

2.2. The minimal polynomial (Lecture 16)

Recall we have an n -dimensional F -vector space V .

- A key tool for studying linear maps is studying polynomials in the maps (we saw how to analyze maps satisfying $T^2 = \text{Id}$, for example).
- We will construct a gadget (the “minimal polynomial”) attached to every linear map on V . It is a polynomial, and will tell us a lot about the map.
- Computationally speaking, this polynomial cannot be found efficiently. It is a tool of theorem-proving in abstract algebra.

DEFINITION 110. Given a polynomial $f \in F[x]$, say $f = \sum_{i=0}^d a_i x^i$ and a map $T \in \text{End}(V)$ set (with $T^0 = \text{Id}$)

$$f(T) = \sum_{i=0}^d a_i T^i.$$

LEMMA 111. *Let $f, g \in F[x]$. Then $(f+g)(T) = f(T) + g(T)$ and $(fg)(T) = f(T)g(T)$. In other words, the map $f \mapsto f(T)$ is a linear map $F[x] \rightarrow \text{End}(V)$, also respecting multiplication (“a map of F -algebras”, but this is beyond our scope).*

PROOF. Do it yourself. □

- Given a linear map our first instinct is to study the kernel and the image. [Aside: the kernel is an *ideal* in the algebra].
- We'll examine the kernel and leave the image for later.

LEMMA 112. *There is a non-zero polynomial $f \in F[x]$ such that $f(T) = 0$. In fact, there is such f with $\deg f \leq n^2$.*

PROOF. $F[x]$ is infinite-dimensional while $\text{End}_F(V)$ is finite-dimensional. Specifically, $\dim_F F[x]^{\leq n^2} = n^2 + 1$ while $\dim_F \text{End}_F(V) = n^2$. □

REMARK 113. We will later show (Theorem of Cayley–Hamilton) that the *characteristic polynomial* $P_T(x) = \det(x\text{Id} - T)$ from basic linear algebra has this property.

- Warning: we are about to divide polynomials with remainder.

PROPOSITION 114. *Let $I = \{f \in F[x] \mid f(T) = 0\}$. Then I contains a unique non-zero monic polynomial of least degree, say $m(x)$, and $I = \{g(x)m(x) \mid g \in F[x]\}$ is the set of multiples of m .*

PROOF. Let $m \in I$ be a non-zero member of least degree. Dividing by the leading coefficient we may assume m monic. Now suppose m' is another such. Then $m - m' \in I$ (this is a subspace) is of strictly smaller degree. It must therefore be the zero polynomial, and m is unique. Clearly if $g \in F[x]$ then $(gm)(T) = g(T)m(T) = 0$. Conversely, given any $f \in I$ we can divide with remainder and write $f = gm + r$ for some $q, r \in F[x]$ with $\deg r < \deg m$. Evaluating at T we find $r(T) = 0$, so $r = 0$ and $f = gm$. □

DEFINITION 115. Call $m(x) = m_T(x)$ the *minimal polynomial* of T .

REMARK 116. We will later prove directly that $\deg m_T(x) \leq n$.

EXAMPLE 117. (Minimal polynomials)

(1) $T = \text{Id}, m(x) = x - 1$.

(2) $T = \begin{pmatrix} 1 \\ \end{pmatrix}, T^2 = 0$ but $T \neq 0$ so $m_T(x) = x^2$.

(3) $T = \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}, T^2 = \begin{pmatrix} 1 & 2 \\ & 1 \end{pmatrix}$ so $(T^2 - \text{Id}) = 2(T - \text{Id})$ so $T^2 - 2T + \text{Id} = 0$ so $m_T(x) \mid (x - 1)^2$. But $T - \text{Id} \neq 0$ so $m_T(x) = (x - 1)^2$.

(4) $T = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, T^2 = \text{Id}$ so $m_T(x) = x^2 - 1 = (x - 1)(x + 1)$.

- In the eigenbasis $\left\{ \begin{pmatrix} 1 \\ \pm 1 \end{pmatrix} \right\}$ the matrix is $\begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$ – we saw this in a previous class.

(5) $T = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}, T^2 = -\text{Id}$ so $m_T(x) = x^2 + 1$.

(a) If $F = \mathbb{Q}$ or $F = \mathbb{R}$ this is irreducible. No better basis.

(b) If $F = \mathbb{C}$ (or $\mathbb{Q}(i)$) then factor $m_T(x) = (x - i)(x + i)$ and in the eigenbasis $\left\{ \begin{pmatrix} 1 \\ \pm i \end{pmatrix} \right\}$

the matrix has the form $\begin{pmatrix} -i & \\ & i \end{pmatrix}$.

- (6) $V = F[x]^{<n}$ (polynomials of degree less than n), $T = \frac{d}{dx}$. Then $T^n = 0$ but $T^{n-1} \neq 0$ (why?) so $m_T(x) = x^n$.
- (7) [To be proved in problem set] Let $D = \text{diag}(a_1, \dots, a_n)$ be diagonal, its entries being the distinct numbers $\{b_1, \dots, b_r\}$ (perhaps with repetition). Then its minimal polynomial is $\prod_{i=1}^r (x - b_i)$ [cf (1),(4),(5)]

We now connect the minimal polynomial with the spectrum.

LEMMA 118 (Spectral calculus). *Suppose that $T\underline{v} = \lambda\underline{v}$. Then $f(T)\underline{v} = f(\lambda)\underline{v}$.*

PROOF. Work it out at home. □

REMARK 119. The same proof shows that if the subspace W is T -invariant ($T(W) \subset W$) then W is $f(T)$ -invariant for all polynomials f .

COROLLARY 120. *If λ is an eigenvalue of T then $m_T(\lambda) = 0$. In particular, if $m_T(0) \neq 0$ then T is invertible (0 is not an eigenvalue)*

We now use the *minimality* of the minimal polynomial.

THEOREM 121. *T is invertible iff $m_T(0) \neq 0$.*

PROOF. Suppose that T is invertible and that $\sum_{i=1}^d a_i T^i = 0$ [note $a_0 = 0$ here]. Then this is not the minimal polynomial since multiplying by T^{-1} also gives

$$\sum_{i=0}^{d-1} a_{i+1} T^i = 0.$$

□

COROLLARY 122. *$\lambda \in F$ is an eigenvalue of T iff λ is a root of $m_T(x)$.*

PROOF. Let $S = T - \lambda \text{Id}$. Then $m_S(x) = m_T(x + \lambda)$. Then $\lambda \in \text{Spec}_F(T) \iff S$ not invertible $\iff m_S(0) = 0 \iff m_T(\lambda) = 0$. □

REMARK 123. The characteristic polynomial $P_T(x)$ also has this property – this is how eigenvalues are found in basic linear algebra.

2.3. Generalized eigenspaces and Cayley–Hamilton (Lectures 17-18)

REMARK 124. A slower schedule covers 2.3.1 in one lecture, 2.3.2 and half of 2.3.3 in another, and finishes 2.3.3 in a third lecture.

2.3.1. Generalized Eigenspaces (Lecture 17). Continue with $T \in \text{End}_F(V)$, $\dim_F(V) = n$. Recall that T is diagonalizable iff V is the direct sum of the eigenspace. For non-diagonalizable maps we need something more sophisticated.

PROBLEM 125. Find a matrix $A \in M_2(F)$ which only has a 1-d eigenspace.

DEFINITION 126. Call $\underline{v} \in V$ a *generalized eigenvector* of T if for some $\lambda \in F$ and $k \geq 1$, $(T - \lambda)^k \underline{v} = \underline{0}$. Let $V_\lambda \subset V$ denote the set of generalized λ -eigenvectors and $\underline{0}$. Call λ a *generalized eigenvalue* of T if $V_\lambda \neq \{\underline{0}\}$.

In particular, if $T\underline{v} = \lambda\underline{v}$ then $\underline{v} \in V_\lambda$.

- PROPOSITION 127 (Generalized eigenspaces). (1) Each V_λ is a T -invariant subspace.
 (2) Let $\lambda \neq \mu$. Then $(T - \mu)$ is invertible on V_λ .
 (3) $V_\lambda \neq \{0\}$ iff $\lambda \in \text{Spec}_F(T)$.

PROOF. Let $\underline{v}, \underline{v}' \in V_\lambda$ be killed by $(T - \lambda)^k, (T - \lambda)^{k'}$ respectively. Then $\alpha \underline{v} + \beta \underline{v}'$ is killed by $(T - \lambda)^{\max\{k, k'\}}$. Also, $(T - \lambda)^k T \underline{v} = T (T - \lambda)^k \underline{v} = \underline{0}$ so $T \underline{v} \in V_\lambda$ as well.

Let $\underline{v} \in \text{Ker}(T - \mu)$ be non-zero. By Lemma 118, for any k we have $(T - \lambda)^k \underline{v} = (\mu - \lambda)^k \underline{v} \neq \underline{0}$ so $\underline{v} \notin V_\lambda$.

Finally, given λ and non-zero $\underline{v} \in V_\lambda$ let k be minimal such that $(T - \lambda)^k \underline{v} = \underline{0}$. Then $(T - \lambda)^{k-1} \underline{v}$ is non-zero and is an eigenvector of eigenvalue λ . \square

THEOREM 128. The sum $\bigoplus_{\lambda \in \text{Spec}_F(T)} V_\lambda \subset V$ is direct.

PROOF. Let $\sum_{i=1}^r \underline{v}_i = \underline{0}$ be a minimal dependence with $\underline{v}_i \in V_{\lambda_i}$ for distinct λ_i . Applying $(T - \lambda_r)^k$ for k large enough to kill \underline{v}_r , we get the dependence.

$$\sum_{i=1}^{r-1} (T - \lambda_r)^k \underline{v}_i = \underline{0}.$$

Now $(T - \lambda_r)^k \underline{v}_i \in V_{\lambda_i}$ since these are T -invariant subspaces, and for $1 \leq i \leq r-1$ is non-zero since $T - \lambda_r$ is invertible there. This shorter dependence contradicts the minimality. \square

REMARK 129. The sum may very well be empty – there are non-trivial maps without eigenvalues (for example $\begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(\mathbb{R})$).

2.3.2. Algebraically closed fields. We all know that sometimes linear maps fail to have eigenvalues, even though they “should”. In this course we’ll blame the field, not the map, for this deficiency.

DEFINITION 130. Call the field F *algebraically closed* if every non-constant polynomial $f \in F[x]$ has a root in F . Equivalently, if every non-constant polynomial can be written as a product of linear factors.

FACT 131 (Fundamental theorem of algebra). \mathbb{C} is algebraically closed.

REMARK 132. Despite the title, this is a theorem of *analysis*.

Discussion. The goal is to create enough eigenvalues so that the generalized eigenspaces explain all of V . The first point of view is that we can simply “define the problem away” by restricting to the case of algebraically closed fields. But this isn’t enough, since sometimes we are given maps over other fields. This already appears in the diagonalizable case, dealt with in 223: we can view $\begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(\mathbb{R})$ instead as $\begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(\mathbb{C})$, at which point it becomes diagonalizable. In other words, we can take a *constructive* point of view:

- Starting with any field F we can “close it” by repeatedly adding roots to polynomial equations until we can’t, obtaining an “algebraic closure” \bar{F} [the difficulty is in showing the process eventually stops].
 - This explains the “closed” part of the name – it’s closure under an operation.

- [Q: do you need the full thing? A: In fact, it's enough to pass to the splitting field of the minimal polynomial]
- We now make this work for linear maps, with three points of view:
 - (1) (matrices) Given $A \in M_n(F)$ view it as $A \in M_n(\bar{F})$, and apply the theory there.
 - (2) (linear maps) Given $T \in \text{End}_F(V)$, fix a basis $\{\underline{v}_i\}_{i=1}^n \subset V$, make the formal span $\bar{V} = \bigoplus_{i=1}^n \bar{F}\underline{v}_i$ and extend T to \bar{V} by the property of having the same matrix.
 - (3) (coordinate free) Given V over F set $\bar{V} = \bar{F} \otimes_F V$ (considering \bar{F} as an F -vectorspace), and extend T (by $\bar{T} = \text{Id}_{\bar{F}} \otimes_F T$).

2.3.3. The direct sum decomposition and Cayley–Hamilton (Lecture 18).

LEMMA 133. *Suppose F is algebraically closed and that $1 \leq \dim_F V < \infty$. Then every $T \in \text{End}_F(V)$ has an eigenvector.*

PROOF. $m_T(x)$ has roots. □

We suppose now that F is algebraically closed, in other words that every linear map has an eigenvalue. The following is the key structure theorem for linear maps:

THEOREM 134. *(with F algebraically closed) We have $V = \bigoplus_{\lambda \in \text{Spec}_F(T)} V_\lambda$.*

PROOF. Let $m_T(x) = \prod_{i=1}^r (x - \lambda_i)^{k_i}$ and let $W = \bigoplus_{i=1}^r V_{\lambda_i}$. Supposing that $W \neq V$, let $\bar{V} = V/W$ and consider the quotient map $\bar{T} \in \text{End}_F(\bar{V})$ defined by $\bar{T}(\underline{v} + W) = T\underline{v} + W$. Since $\dim_F \bar{V} \geq 1$, \bar{T} has an eigenvalue there. We first check that this eigenvalue is one of the λ_i . Indeed, for any polynomial $f \in F[x]$, $f(\bar{T})(\underline{v} + W) = (f(T)\underline{v}) + W$, and in particular $m_T(\bar{T}) = 0$ and hence $m_{\bar{T}} | m_T$.

Renumbering the eigenvalues, we may assume $\bar{V}_{\lambda_r} \neq \{0\}$, and let $\underline{v} \in V$ be such that $\underline{v} + W \in \bar{V}_{\lambda_r}$ is non-zero, that is $\underline{v} \notin W$. Since $\prod_{i=1}^{r-1} (\bar{T} - \lambda_i)^{k_i}$ is invertible on \bar{V}_{λ_r} , $\underline{u} = \prod_{i=1}^{r-1} (T - \lambda_i)^{k_i} \underline{v} \notin W$. But $(T - \lambda_r)^{k_r} \underline{u} = m_T(T)\underline{v} = 0$ means that $\underline{u} \in V_{\lambda_r} \subset W$, a contradiction. □

PROPOSITION 135. *In $m_T(x) = \prod_{i=1}^r (x - \lambda_i)^{k_i}$, the number k_i is the minimal k such that $(T - \lambda_i)^k = 0$ on V_{λ_i} .*

PROOF. Let T_i be the restriction of T to V_{λ_i} . Then $(T_i - \lambda_i)^k$ is the minimal polynomial by assumption. But $m_T(T_i) = 0$. It follows that $(x - \lambda_i)^k | m_T$ and hence that $k \leq k_i$. Conversely, since $\prod_{j \neq i} (T - \lambda_j)^{k_j}$ is invertible on V_{λ_i} , we see that $(T - \lambda_i)^{k_i} = 0$ there, so $k_i \geq k$. □

Summary of the construction so far:

- F algebraically closed field, $\dim_F V = n$, $T \in \text{End}_F(V)$.
- $m_T(x) = \prod_{i=1}^r (x - \lambda_i)^{k_i}$ the minimal polynomial.
- Then $V = \bigoplus_{i=1}^r V_{\lambda_i}$ where on V_{λ_i} we have $(T - \lambda_i)^{k_i} = 0$ but $(T - \lambda_i)^{k_i-1} \neq 0$.

We now study the restriction of T to each V_{λ_i} , via the map $N = T - \lambda_i$, which is *nilpotent* of degree k_i .

DEFINITION 136. A map $N \in \text{End}_F(V)$ such that $N^k = 0$ for some k is called *nilpotent*. The smallest such k is called its *degree of nilpotence*.

LEMMA 137. *Let $N \in \text{End}_F(V)$ be nilpotent. Then its degree of nilpotence is at most $\dim_F V$.*

PROOF. Exercise. □

PROOF. Define subspaces V_k by $V_0 = V$ and $V_{i+1} = N(V_i)$. Then $V = V_0 \supset V_1 \cdots \supset V_i \supset \cdots$. If at any stage $V_i = V_{i+1}$ then $V_{i+j} = V_i$ for all $j \geq 1$, and in particular $V_i = \{0\}$ (since $V_k = 0$). It follows that for $i < k$, $\dim V_{i+1} < \dim V_i$ and the claim follows. \square

COROLLARY 138 (Cayley–Hamilton Theorem). *Suppose F is algebraically closed. Then $m_T(x) \mid p_T(x)$ and, equivalently, $p_T(T) = 0$. In particular, $\deg m_T \leq \dim_F V$.*

Recall that the *characteristic polynomial* of T is the polynomial $p_T(x) = \det(x\text{Id} - T)$ of degree $\dim_F V$, and that is also has the property that $\lambda \in \text{Spec}_F(T)$ iff $p_T(\lambda) = 0$.

PROOF. The linear map $x\text{Id} - T$ respects the decomposition $V = \bigoplus_{i=1}^r V_{\lambda_i}$. We thus have $p_T(x) = \prod_{i=1}^r p_{T|_{V_{\lambda_i}}}(x)$. Since $p_{T|_{V_{\lambda_i}}}(x)$ has the unique root λ , it is the polynomial $(x - \lambda)^{\dim_F V_{\lambda_i}}$, so

$$p_T(x) = \prod_{i=1}^r (x - \lambda_i)^{\dim V_{\lambda_i}}.$$

Finally, k_i is the degree of nilpotence of $(T - \lambda_i)$ on V_{λ_i} . Thus $k_i \leq \dim_F V_{\lambda_i}$. \square

We now resolve a lingering issue:

LEMMA 139. *The minimal polynomial is independent of the choice of the field. In particular, the Cayley–Hamilton Theorem holds over any field.*

PROOF. Whether $\{1, T, \dots, T^{d-1}\} \subset \text{End}_F(V)$ are linearly dependent or not does not depend on the field. \square

THEOREM 140 (Cayley–Hamilton). *Over any field we have $m_T(x) \mid p_T(x)$ or, equivalently, $p_T(T) = 0$.*

PROOF. Extend scalars to an algebraic closure. This does not change either of the polynomials m_T, p_T . \square

2.4. Nilpotent maps and Jordan blocks (Lectures 19-20)

2.4.1. Jordan blocks (Lecture 19). We finally turn to the problem of finding good bases for linear maps, starting with the nilpotent case. Here F can be an arbitrary field.

To start with, consider the filtration $\{0\} \subsetneq \text{Ker}(N) \subsetneq \text{Ker}(N^2) \subsetneq \cdots \subsetneq V = \text{Ker}(N^k)$ where k is the degree of nilpotence. Choose a basis as follows:

- (1) Choose a basis of size m_1 in $\text{Ker}(N)$.
- (2) Add m_2 vectors to get a basis of $\text{Ker}(N^2) \supset \text{Ker}(N)$
- \vdots
- (3) Add m_k vectors to get a basis of $V = \text{Ker}(N^k)$.

What is the matrix of N in this basis? It is block-upper triangular. On the diagonal the blocks are $m_i \times m_i$, all zero. Above each block is a block of size $m_{i-1} \times m_i$ where every column is non-zero (every if $\underline{v} \in \text{Ker}(N^r) \setminus \text{Ker}(N^{r-1})$ then $N\underline{v} \in \text{Ker}(N^{r-1}) \setminus \text{Ker}(N^{r-2})$). Above that no information.

We now try to improve this by carefully choosing the basis to encode more of the action of N .

LEMMA 141. *Let $N \in \text{End}(V)$ be nilpotent. Let $B \subset V$ be a set of vectors such that $N(B) \subset B \cup \{0\}$. Then B is linearly independent iff $B \cap \text{Ker}(N)$ is.*

PROOF. One direction is clear. For the converse, let $\sum_{i=1}^r a_i \underline{v}_i = \underline{0}$ be a minimal dependence in B . Apply N^s where s is maximal such that for some i , $N^s \underline{v}_i \neq \underline{0}$ (perhaps $s = 0$). Then

$$\sum_{i=1}^r a_i N^s \underline{v}_i = \underline{0}$$

where each $N^s \underline{v}_i \in B \cup \{\underline{0}\}$. If some $N^s \underline{v}_i = \underline{0}$ the new dependence is shorter. Thus they are all non-zero then by the maximality of s they are all in $\text{Ker}(N)$, so we obtain a dependence among $B \cap \text{Ker}(N)$, a contradiction. \square

COROLLARY 142. Let $N \in \text{End}(V)$ and let $\underline{v} \in V$ be non-zero such that $N^k \underline{v} = \underline{0}$ for some k (wlog minimal). Then $\{N^i \underline{v}\}_{i=0}^{k-1}$ is linearly independent.

PROOF. N is nilpotent on $\text{Span}\{N^i \underline{v}\}_{i=0}^{k-1}$, this set is invariant, and its intersection with $\text{Ker} N$ is exactly $\{N^{k-1} \underline{v}\} \neq \{\underline{0}\}$. \square

2.4.2. Jordan canonical form for nilpotent maps (Lecture 20). Our goal is now to decompose V as a direct sum of N subspaces (“Jordan blocks”) each of which has a basis as in the Corollary.

THEOREM 143 (Jordan form for nilpotent maps). Let $N \in \text{End}_F(V)$ be nilpotent. We then have a decomposition $V = \bigoplus_{j=1}^r V_j$ where each V_j is an N -invariant Jordan block.

EXAMPLE 144. $A = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \\ -1 & -2 & -3 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} (1 \ 2 \ 3)$.

- $A^2 = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} (1 \ 2 \ 3) \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} (1 \ 2 \ 3) = 0$, so A is nilpotent. The characteristic polynomial must be x^3 .
- The image of A is $\text{Span}\left\{\begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}\right\}$. Since $A \begin{pmatrix} 3 \\ -1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$, $\text{Span}\left\{\begin{pmatrix} 3 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}\right\}$ is a block.
- Taking any other vector in the kernel (say, $\begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$) we get the basis $\left\{\begin{pmatrix} 3 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}\right\}$ in which A has the matrix

$$\begin{pmatrix} \begin{pmatrix} 0 & 1 \\ & 0 \end{pmatrix} & \\ & (0) \end{pmatrix}.$$

PROOF. Let N have degree of nilpotence d and kernel W . For $1 \leq k \leq d$ define $W_k = \text{Im}(N^k) \cap W$, so that $W_0 = W \supset W_1 \supset W_d = \{0\}$. Now choose a basis C of W compatible with this decomposition – in other words choose subsets $C_k \subset W_k$ such that $\cup_{k \geq k'} C_k$ is a basis for $W_{k'}$. Let $C = \cup_{k=0}^{d-1} C_k = \{\underline{v}_i\}_{i \in I}$ and for each i define k_i by $\underline{v}_i \in C_{k_i}$. Choose \underline{u}_i such that $N^{k_i} \underline{u}_i = \underline{v}_i$, and for

$1 \leq j \leq k_i$ set $\underline{v}_{i,j} = N^{k_i-j} \underline{u}_i$ so that $\underline{v}_{i,1} = \underline{u}_i$ and in general $N \underline{v}_{i,j} = \begin{cases} \underline{v}_{i,j-1} & j \geq 1 \\ \underline{0} & j = 1 \end{cases}$. It is clear that $\text{Span}_F \{ \underline{v}_{i,j} \}_{j=1}^{k_i}$ is a Jordan block, and that $B = \{ \underline{u}_{i,j} \}_{i,j}$ is a union of Jordan blocks. \square

- **The set B is linearly independent:** by construction, $N(B) \subset B \cup \{ \underline{0} \}$ and $B \cap W = C$ is independent.
- **The set B is spanning:** We prove by induction on $k \leq d$ that $\text{Span}_F(B) \supset \text{Ker}(N^k)$. This is clear for $k = 0$; suppose the result for $0 \leq k < d$, and let $\underline{v} \in \text{Ker}(N^{k+1})$. Then $N^k \underline{v} \in W_k$, so we can write

$$\begin{aligned} N^k \underline{v} &= \sum_{i:k_i \geq k} a_i \underline{v}_i \\ &= \sum_{i:k_i \geq k} a_i N^k (\underline{v}_{i,k+1}) . \end{aligned}$$

It follows that

$$N^k \left(\underline{v} - \sum_{i:k_i \geq k} a_i \underline{v}_{i,k+1} \right) = \underline{0} .$$

By induction, $\underline{v} - \sum_{i:k_i \geq k} a_i \underline{v}_{i,k} \in \text{Span}_F(B)$, and it follows that $\underline{v} \in \text{Span}_F(B)$.

DEFINITION 145. A *Jordan basis* is a basis as in the Theorem.

LEMMA 146. Any Jordan basis for N has exactly $\dim_F W_{k-1} - \dim_F W_k$ blocks of length k . Equivalently, up to permuting the blocks, N has a unique matrix in Jordan form.

PROOF. Let $\{ \underline{v}_{i,j} \}$ be a Jordan basis. Then $\text{Ker} N = \text{Span} \{ \underline{v}_{i,1} \}$, while $\{ \underline{v}_{i,j} \mid k_i \geq k, j \leq k_i - k \}$ is a basis for $\text{Im}(N^k)$. Clearly $\{ \underline{v}_{i,1} \mid k_i \geq k \}$ then spans W_k and the claim follows. \square

2.5. The Jordan canonical form (Lecture 21)

THEOREM 147 (Jordan canonical form). Let $T \in \text{End}_F(V)$ and suppose that m_T splits into linear factors in F (for example, that F is algebraically closed). Then there is a basis $\{ \underline{v}_{\lambda,i,j} \}_{\lambda,i,j}$ of V

such that $\{ \underline{v}_{\lambda,i,j} \}_{i,j} \subset V_\lambda$ is a basis, and such that $(T - \lambda) \underline{v}_{\lambda,i,j} = \begin{cases} \underline{v}_{\lambda,i,j-1} & j \geq 1 \\ \underline{0} & j = 1 \end{cases}$. Furthermore,

writing $W_\lambda = \text{Ker}(T - \lambda)$ for the eigenspace, we have for each λ , that $1 \leq i \leq \dim_F W_\lambda$ and that the number of i such that $1 \leq j \leq k$ is exactly $\dim_F ((T - \lambda)^{k-1} V_\lambda \cap W_\lambda) - \dim_F ((T - \lambda)^k V_\lambda \cap W_\lambda)$. Equivalently, T has a unique matrix in Jordan canonical form up to permuting the blocks.

COROLLARY 148. The algebraic multiplicity of λ is $\dim_F V_\lambda$. The geometric multiplicity is the number of blocks.

EXAMPLE 149 (Jordan forms). (1) $A_1 = \begin{pmatrix} 2 & 2 & 3 \\ 1 & 3 & 3 \\ -1 & -2 & -2 \end{pmatrix} = I + A$. This has character-

istic polynomial $(x - 1)^3$, $A_1 - I = A$ and we are back in example 144.

(2) (taken from Wikibooks:Linear Algebra) $p_B(x) = (x-6)^4$.

$$\text{Let } B = \begin{pmatrix} 7 & 1 & 2 & 2 \\ 1 & 4 & -1 & -1 \\ -2 & 1 & 5 & -1 \\ 1 & 1 & 2 & 8 \end{pmatrix}, B' = B - 6I = \begin{pmatrix} 1 & 1 & 2 & 2 \\ 1 & -2 & -1 & -1 \\ -2 & 1 & -1 & -1 \\ 1 & 1 & 2 & 2 \end{pmatrix}. \text{ Gaussian elimina-}$$

$$\text{tion shows } B' = E \begin{pmatrix} 3 & -3 & 0 & 0 \\ 1 & -2 & -1 & -1 \\ -3 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, B'^2 = \begin{pmatrix} 0 & 3 & 3 & 3 \\ 0 & 3 & 3 & 3 \\ 0 & -6 & -6 & -6 \\ 0 & 3 & 3 & 3 \end{pmatrix} \text{ and } B'^3 = 0. \text{ Thus}$$

$\text{Ker } B' = \{(x, y, z, w)^t \mid x = y = -(z + w)\}$ is two-dimensional. We see that the image of B'^2 is spanned by $(3, 3, -6, 3)^t$, which is (say) $B'(2, -1, -1, 2)^t$ which (being the last column) was $B'(0, 0, 0, 1)^t$. Another vector in the kernel is $(-1, -1, 1, 0)^t$, and we get the

$$\text{Jordan basis } \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ -1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 3 \\ -6 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ -1 \\ 1 \\ 0 \end{pmatrix} \right\}.$$

$$(3) C = \begin{pmatrix} 4 & 0 & 1 & 0 \\ 2 & 2 & 3 & 0 \\ -1 & 0 & 2 & 0 \\ 4 & 0 & 1 & 2 \end{pmatrix} \text{ acting on } V = \mathbb{R}^4 \text{ with } p_C(x) = (x-2)^2(x-3)^2. \text{ Then } C - 2I =$$

$$\begin{pmatrix} 2 & 0 & 1 & 0 \\ 2 & 0 & 3 & 0 \\ -1 & 0 & 0 & 0 \\ 4 & 0 & 1 & 0 \end{pmatrix}, C - 3I = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 2 & -1 & 3 & 0 \\ -1 & 0 & -1 & 0 \\ 4 & 0 & 1 & -1 \end{pmatrix}, (C - 3I)^2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -3 & 1 & -4 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 2 & 1 \end{pmatrix}. \text{ Thus}$$

$\text{Ker}(C - 2I) = \text{Span}\{\underline{e}_2, \underline{e}_4\}$, which must be the 2d generalized eigenspace V_2 giving two 1×1 blocks. For $\lambda = 3$, $\text{Ker}(C - 3I) = \{(x, y, z, w)^t \mid z = y = -x, w = 3x\} = \text{Span}\{(1, -1, -1, 3)^t\}$. This isn't the whole generalized eigenspace, and

$$\text{Ker}(C - 3I)^2 = \{(x, y, z, w)^t \mid y = 3x + 4z, w = x - 2z\} = \text{Span}\{(1, -1, -1, 3)^t, (1, 3, 0, 1)^t\}.$$

This must be the generalized eigenspace V_3 , since it's 2d. We need to find the image of $(C - 3I)[V_3]$. One vector is in the kernel, so we try the other one, and indeed $(C - 3I)(1, 3, 0, 1)^t = (1, -1, -1, 3)$. This gives us a 2x2 block, so in the basis

$$\left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 0 \\ 1 \end{pmatrix} \right\} \text{ the matrix has the form } \begin{pmatrix} (2) & & & \\ & (2) & & \\ & & (3 & 1) \\ & & & (3) \end{pmatrix}. \text{ Note}$$

how the image of $(C - 3I)^2$ is exactly V_2 (why?)

(4) $V = \mathbb{R}^6$. $p_D(x) = t^6 + 3t^5 - 10t^3 - 15t^2 - 9t - 2 = (t+1)^5(t-2)$:

$$D = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & -1 \\ 0 & -8 & 4 & -3 & 1 & -3 \\ -3 & 13 & -8 & 6 & 2 & 9 \\ -2 & 14 & -7 & 4 & 2 & 10 \\ 1 & -18 & 11 & -11 & 2 & -6 \\ -1 & 19 & -11 & 10 & -2 & 7 \end{pmatrix}, D+I = \begin{pmatrix} 1 & 0 & 0 & 0 & -1 & -1 \\ 0 & -7 & 4 & -3 & 1 & -3 \\ -3 & 13 & -7 & 6 & 2 & 9 \\ -2 & 14 & -7 & 5 & 2 & 10 \\ 1 & -18 & 11 & -11 & 3 & -6 \\ -1 & 19 & -11 & 10 & -2 & 8 \end{pmatrix},$$

$$(D+I)^2 = \begin{pmatrix} 1 & -1 & 0 & 1 & -2 & -3 \\ -2 & -16 & 9 & -11 & 4 & -3 \\ -1 & 37 & -18 & 17 & 2 & 21 \\ 1 & 35 & -18 & 19 & -2 & 15 \\ -1 & -53 & 27 & -28 & 2 & -24 \\ 2 & 52 & -27 & 29 & -4 & 21 \end{pmatrix}, (D+I)^3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -54 & 27 & -27 & 0 & -27 \\ 0 & 108 & -54 & 54 & 0 & 54 \\ 0 & 108 & -54 & 54 & 0 & 54 \\ 0 & -162 & 81 & -81 & 0 & -81 \\ 0 & 162 & -81 & 81 & 0 & 81 \end{pmatrix}.$$

(5) First, V_2 must be a 1-dimensional eigenspace. Gaussian elimination finds the eigenvector $(01, -2, -2, 3, -3)^t$. Next, V_{-1} must be 5-dimensional. Row-reduction gives: $D+I \rightarrow$

$$\begin{pmatrix} 1 & 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 1 & 0 & -1/2 \\ 0 & 1 & 0 & 0 & 1 & 3/2 \\ 0 & 0 & 1 & 0 & 2 & 3/2 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, (D+I)^2 \rightarrow \begin{pmatrix} 2 & 0 & -1 & 3 & -4 & -5 \\ 0 & 2 & -1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \text{ So the Ker}(D+I)$$

is two-dimensional (since $(D+I)^2 \neq 0$ there will be a block of size at least 3; since $(D+I)^3$ has rank one, it has the 5d kernel $V_{-1} = \{x \mid x_3 = 2x_2 + x_4 + x_6\}$ so the largest block is 3, and so the other block must have size 2. We need a vector from the generalized eigenspace in the image of $(D+I)^2$. Since $(D+I)^3 e_1 = 0$ but the first column of $(D+I)^2$ is non-zero, we see that $(D+I)^2 e_1 = (1, -2, -1, 1, -1, 2)^t$ has preimage $(D+I) e_1 = (1, 0, -3, -2, 1, -1)^t$, and we obtain our first block. Next, we need an eigenvector in the kernel and image of $D+I$, but any vector in the kernel is also in the image (no blocks of size 1), so we can take any vector in $\text{Ker}(D+I)$ independent of the one we already have. Using the row-reduced form we see that $(1, -1, -2, 0, 1, 0)^t$ is such a vector. Then we solve

$$\begin{pmatrix} 1 & 0 & 0 & 0 & -1 & -1 \\ 0 & -7 & 4 & -3 & 1 & -3 \\ -3 & 13 & -7 & 6 & 2 & 9 \\ -2 & 14 & -7 & 5 & 2 & 10 \\ 1 & -18 & 11 & -11 & 3 & -6 \\ -1 & 19 & -11 & 10 & -2 & 8 \end{pmatrix} \underline{x} = \begin{pmatrix} 1 \\ -1 \\ -2 \\ 0 \\ 1 \\ 0 \end{pmatrix},$$

finding for example the vector $(1, 0, -1, -1, 0, 0)^t$ and our second block. We conclude

that in the basis $\left\{ \begin{pmatrix} 0 \\ 1 \\ -2 \\ -2 \\ 3 \\ -3 \end{pmatrix}, \begin{pmatrix} 1 \\ -2 \\ -1 \\ 1 \\ -1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -3 \\ -2 \\ 1 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ -2 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right\}$ the matrix has the

form

$$\left(\begin{array}{c} (2) \\ \left(\begin{array}{cc} -1 & 1 \\ & -1 & 1 \\ & & -1 \end{array} \right) \\ \left(\begin{array}{cc} -1 & 1 \\ & -1 \end{array} \right) \end{array} \right)$$

CHAPTER 3

Vector and matrix norms

For the rest of the course our field of scalars is either \mathbb{R} or \mathbb{C} .

3.1. Norms on vector spaces (Lecture 22)

3.1.1. Review of metric spaces.

DEFINITION 150. A *metric space* is a pair (X, d_X) where X is a set, and $d_X : X \times X \rightarrow \mathbb{R}_{\geq 0}$ is a function such that for all $x, y, z \in X$, $d_X(x, y) = 0$ iff $x = y$, $d_X(x, y) = d_X(y, z)$ and (the *triangle inequality*) $d_X(x, z) \leq d_X(x, y) + d_X(y, z)$.

NOTATION 151. For $x \in X$ and $r \geq 0$ we write $B_X(x, r) = \{y \in X \mid d_X(x, y) \leq r\}$ for the closed ball of radius r around x , $B_X^\circ(x, r) = \{y \in X \mid d_X(x, y) < r\}$ for the open ball.

DEFINITION 152. Let $(X, d_X), (Y, d_Y)$ be metric spaces and let $f : X \rightarrow Y$ be a function.

- (1) We say f is *continuous* if $\forall x \in X : \forall \varepsilon > 0 : \exists \delta > 0 : f(B_X(x, \delta)) \subset B_Y(f(x), \varepsilon)$.
- (2) We say f is *uniformly continuous* $\forall \varepsilon > 0 : \exists \delta > 0 : \forall x \in X : f(B_X(x, \delta)) \subset B_Y(f(x), \varepsilon)$.
- (3) We say f is *Lipschitz continuous* if in (2) we can take $\delta = L\varepsilon$, in other words if for all $x \neq x' \in X$,

$$d_Y(f(x), f(x')) \leq L d_X(x, x').$$

In that case we let $\|f\|_{\text{Lip}}$ denote the smallest L for which this holds.

Clearly (3) \Rightarrow (2) \Rightarrow (1).

LEMMA 153. *The composition of two functions of type (1),(2),(3) is again a function of that type. In particular, $\|f \circ g\|_{\text{Lip}} \leq \|f\|_{\text{Lip}} \|g\|_{\text{Lip}}$.*

DEFINITION 154. We call the metric space (X, d_X) *complete* if every Cauchy sequence converges.

3.1.2. Norms. Fix a vector space V .

DEFINITION 155. A *norm* on V is a function $\|\cdot\| : V \rightarrow \mathbb{R}_{\geq 0}$ such that $\|\underline{v}\| = 0$ iff $\underline{v} = \underline{0}$, $\|\alpha \underline{v}\| = |\alpha| \|\underline{v}\|$ and $\|\underline{u} + \underline{v}\| \leq \|\underline{u}\| + \|\underline{v}\|$. A *normed space* is a pair $(V, \|\cdot\|)$.

LEMMA 156. *Let $\|\cdot\|$ be a norm on V . Then the function $d(\underline{u}, \underline{v}) = \|\underline{u} - \underline{v}\|$ is a metric.*

EXERCISE 157. The map $\|\cdot\| \mapsto d$ is a bijection between norms on V and metrics on V which are (1) translation-invariant $d(\underline{u}, \underline{v}) = d(\underline{u} + \underline{w}, \underline{v} + \underline{w})$ and (2) 1-homogenous: $d(\alpha \underline{u}, \alpha \underline{v}) = |\alpha| d(\underline{u}, \underline{v})$.

The restriction of a norm to a subspace is a norm.

3.1.3. Finite-dimensional examples.

EXAMPLE 158. Standard norms on \mathbb{R}^n and \mathbb{C}^n :

- (1) The *supremum norm* $\|\underline{v}\|_\infty = \max\{|v_i|\}_{i=1}^n$, parametrizing *uniform convergence*.
- (2) $\|\underline{v}\|_1 = \sum_{i=1}^n |v_i|$.
- (3) The *Euclidean norm* $\|\underline{v}\|_2 = \left(\sum_{i=1}^n |v_i|^2\right)^{1/2}$, connected to the *inner product* $\langle \underline{u}, \underline{v} \rangle = \sum_{i=1}^n \bar{u}_i v_i$ (prove Δ inequality from this by squaring norm of sum).
- (4) For $1 < p < \infty$, $\|\underline{v}\|_p = \left(\sum_{i=1}^n |v_i|^p\right)^{1/p}$.

PROOF. These functions are clearly homogeneous, and clearly are non-zero if $\underline{v} \neq 0$; the only non-trivial part is the triangle inequality (“Minkowsky’s inequality”). This is easy for $p = 1, \infty$, well-known for $p = 2$. Other cases resolved in supplement to PS8. \square

EXERCISE 159. Show that $\lim_{p \rightarrow \infty} \|\underline{v}\|_p = \|\underline{v}\|_\infty$.

We have a geometric interpretation. The *unit ball* of a norm is the set $B = B(\underline{0}, 1) = \{\underline{v} \in V \mid \|\underline{v}\| \leq 1\}$. This determines the norm ($\frac{1}{\|\underline{v}\|}$ is the largest α such that $\alpha \underline{v} \in B$). Now applying a linear map to B gives a the ball of a new norm.

EXERCISE 160. Draw the unit balls of

PROPOSITION 161 (Pullback). *Let $T : U \hookrightarrow V$ be an injectivel linear map. Let $\|\cdot\|_V$ be a norm on V . Then $\|\underline{u}\| \stackrel{\text{def}}{=} \|T\underline{u}\|_V$ defines a norm on U .*

PROOF. Easy check. \square

3.1.4. Infinite-dimensional examples.

Now the norm comes first, the space second.

EXAMPLE 162. For a set X let $\ell^\infty(X) = \{f \in F^X \mid \sup\{|f(x)| : x \in X\} < \infty\}$, $\|f\|_\infty = \sup_{x \in X} |f(x)|$.

PROOF. The map $\|\cdot\|_\infty : X^F \rightarrow [0, \infty]$ satisfies the axioms of a norm, suitably extended to include the value ∞ . That the set of vectors of finite norm is a subspace follows from the scaling and triangle inequalities. \square

REMARK 163. A vector space with basis B can be embedded into $\ell^\infty(B)$ (we’ve basically seen this).

EXAMPLE 164. $\ell^p(\mathbb{N}) = \{\underline{a} \in F^\mathbb{N} : \sum_{i=1}^\infty |a_i|^p < \infty\}$ with the obvious norm.

In the continuous case we a construction from earlier in the course:

DEFINITION 165. $L^p(\mathbb{R}) = \{f : \mathbb{R} \rightarrow F \text{ [measurable]} \mid \int_{\mathbb{R}} |f(x)|^p dx < \infty\} / \{f \mid f = 0 \text{ a.e.}\}$ with the natural norm.

REMARK 166. The quotient is *essential*: for actualy functions, can have $\int |f(x)|^p dx = 0$ without $f = 0$ exactly. In particular, elements of $L^p(\mathbb{R})$ don’t have specific values.

FACT 167. *In each equivalence class in $L^p(\mathbb{R})$ there is at most one continuous representative.*

So part of PDE is about whether an L^p solution can be promoted to a continuous functiuon. We give an example theorem:

THEOREM 168 (Elliptic regularity). *Let $\Omega \subset \mathbb{R}^2$ be a domain, and let $f \in L^2(\Omega)$ satisfy $\Delta f = \lambda f$ distributionally: for $g \in C_c^\infty(\Omega)$, $\int_\Omega f \Delta g = \lambda \int_\Omega f g$. Then there is a smooth function \bar{f} such that $\Delta \bar{f} = \lambda \bar{f}$ pointwise and such that $f = \bar{f}$ almost everywhere.*

3.1.5. Converges in the norm. While there are many norms on \mathbb{R}^n , it turns out that there is only one notion of convergence.

LEMMA 169. *Every norm on \mathbb{R}^n is a continuous function.*

PROOF. Let $M = \max_i \|e_i\|$. Then

$$\|\underline{x}\| = \left\| \sum_{i=1}^n x_i e_i \right\| \leq \sum_{i=1}^n |x_i| \|e_i\| \leq M \|\underline{x}\|_1.$$

In particular,

$$\left| \|\underline{x}\| - \|\underline{y}\| \right| \leq \|\underline{x} - \underline{y}\| \leq M \|\underline{x} - \underline{y}\|_1.$$

□

DEFINITION 170. Call two norms equivalent if there are $0 < m \leq M$ such that $m \|\underline{x}\| \leq \|\underline{x}\|' \leq M \|\underline{x}\|$ holds for all $\underline{x} \in V$.

EXERCISE 171. This is an equivalence relation. The norms are equivalent iff the same sequences of vectors satisfy $\lim_{n \rightarrow \infty} \underline{x}_n = \underline{0}$.

THEOREM 172. *All norms on \mathbb{R}^n (and \mathbb{C}^n) are equivalent.*

PROOF. It is enough to show that they are all equivalent to $\|\cdot\|_1$. Accordingly let $\|\cdot\|$ be any other norm. Then the Lemma shows that there is M such that

$$\|\underline{x}\| \leq M \|\underline{x}\|_1.$$

Next, the “sphere” $\{\underline{x} \mid \|\underline{x}\|_1 = 1\}$ is closed and bounded, hence compact. Accordingly let $m = \min \{\|\underline{x}\| \mid \|\underline{x}\|_1 = 1\}$. Then $m > 0$ since $\|\underline{0}\|_1 = 0 \neq 1$. Finally, for any $\underline{x} \neq \underline{0}$ we have

$$\frac{\|\underline{x}\|}{\|\underline{x}\|_1} = \left\| \frac{\underline{x}}{\|\underline{x}\|_1} \right\| \geq m$$

since $\left\| \frac{\underline{x}}{\|\underline{x}\|_1} \right\|_1 = 1$. It follows that

$$m \|\underline{x}\|_1 \leq \|\underline{x}\| \leq M \|\underline{x}\|_1$$

□

3.2. Norms on matrices (Lectures 23-24)

DEFINITION 173. Let U, V be normed spaces. A map $T: U \rightarrow V$ is called *bounded* if there is $M \geq 0$ such that $\|T\underline{u}\|_V \leq M \|\underline{u}\|_U$ for all $\underline{u} \in U$. The smallest such M is called the (*operator*) *norm* of T .

REMARK 174. Motivation: Let U be the space of initial data for an evolution equation (say wave, or heat). Let V be the space of possible states at time t . Let T be “time evolution”. Then a key part of PDE is finding norms in which T is bounded as a map from U to V . This shows that solution exist, and that they are unique.

EXAMPLE 175. The identity map has norm 1. Now consider the matrix $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ acting on \mathbb{R}^2 .

(1) As a map from $\ell^1 \rightarrow \ell^1$ we have

$$\left\| A \begin{pmatrix} x \\ y \end{pmatrix} \right\|_1 = |x+y| + |y| \leq 2 \left\| \begin{pmatrix} x \\ y \end{pmatrix} \right\|_1,$$

with equality if $x = 0$. Thus $\|A\|_1 = 2$.

(2) Next,

$$\left\| A \begin{pmatrix} x \\ y \end{pmatrix} \right\|_2^2 = |x+y|^2 + |y|^2 \leq \frac{3+\sqrt{5}}{2} |x^2 + y^2|.$$

(3) Finally,

$$\left\| A \begin{pmatrix} x \\ y \end{pmatrix} \right\|_\infty = \max\{|x+y|, |y|\} \leq 2 \max\{|x|, |y|\}$$

with equality if $x = y$, Thus $\|A\|_\infty = 2$.

EXAMPLE 176. Consider $D_x: C_c^\infty(\mathbb{R}) \rightarrow C_c^\infty(\mathbb{R})$. This is not bounded in any norm (consider $f(x) = e^{2\pi i k x}$).

LEMMA 177. *Every map of finite-dimensional spaces is bounded.*

PROOF. Identify U with \mathbb{R}^n . Then the $\|\cdot\|_U$ is equivalent with $\|\cdot\|_1$, so there is A such that $\|\underline{u}\|_1 \leq A \|\underline{u}\|_U$. Now the map $\underline{u} \mapsto \|T\underline{u}\|_V$ is 1-homogenous and satisfies the triangle inequality, so by the proof of Lemma 169 there is B so that $\|T\underline{u}\|_V \leq B \|\underline{u}\|_1 \leq (AB) \|\underline{u}\|_U$. \square

LEMMA 178. *Let T, S be bounded and composable. Then ST is bounded and $\|ST\| \leq \|S\| \|T\|$.*

PROOF. For any $\underline{u} \in U$, $\|ST\underline{u}\|_W \leq \|S\| \|T\underline{u}\|_V \leq \|S\| \|T\| \|\underline{u}\|_U$. \square

PROPOSITION 179. *The operator norm is a norm on $\text{Hom}_b(U, V)$, the space of bounded maps $U \rightarrow V$.*

PROOF. For any $S, T \in \text{Hom}_b(U, V)$, $|\alpha| \|T\| + \|S\|$ is a bound for $\alpha T + S$. Since the zero map is bounded it follows that $\text{Hom}_b(U, V) \subset \text{Hom}(U, V)$ is a subspace, and setting $\alpha = 1$ gives the triangle inequality. If $T \neq 0$ then there is \underline{u} such that $T\underline{u} \neq \underline{0}$ at which point

$$\|T\| \geq \frac{\|T\underline{u}\|}{\|\underline{u}\|} > 0.$$

Finally, $\|(\alpha T)\underline{u}\| = |\alpha| \|T\underline{u}\| \leq |\alpha| \|T\| \|\underline{u}\|$ so $\|\alpha T\| \leq |\alpha| \|T\|$. But then

$$\|T\| = \left\| \frac{1}{|\alpha|} \alpha T \right\| \leq \frac{1}{|\alpha|} \|\alpha T\|$$

gives the reverse inequality. \square

3.3. Example: eigenvalues and the power method (Lecture 25)

Let A be diagonalizable. Want eigenvalues of A . Raising A to large powers selects the eigenvalue with largest component.

- Algorithm: multiply by A and renormalize.
- Advantage: if A sparse only need to multiply by A .
- Rate of convergence related to spectral gap.

3.4. Sequences and series of vectors and matrices (Lectures 26-27)

3.4.1. Completeness (Lecture 26).

DEFINITION 180. A metric space is complete if every Cauchy sequence in it converges.

EXAMPLE 181. \mathbb{R}, \mathbb{R}^n in any norm. $\text{Hom}(\mathbb{R}^n, \mathbb{R}^m)$ (because isom to \mathbb{R}^{mn}).

FACT 182. Any metric space has a completion. [note associated universal property and hence uniqueness]

THEOREM 183. Let $(U, \|\cdot\|_U), (V, \|\cdot\|_V)$ be normed spaces with V complete. Then $\text{Hom}_b(U, V)$ is complete with respect to the operator norm.

PROOF. Let $\{T_n\}_{n=1}^\infty$ be a Cauchy sequences of linear maps. For fixed $\underline{u} \in U$, the sequence $\{T_n \underline{u}\}$ is Cauchy: $\|(T_n \underline{u} - T_m \underline{u})\|_V \leq \|T_n - T_m\| \|\underline{u}\|$. It is therefore convergent – call the limit $T \underline{u}$. This is linear since $\alpha T_n \underline{u} + T_n \underline{u}'$ converges to $\alpha T \underline{u} + T \underline{u}'$ while $T_n(\alpha \underline{u} + \underline{u}')$ converges to $T(\alpha \underline{u} + \underline{u}')$.

Since $\| \|T_n\| - \|T_m\| \| \leq \|T_n - T_m\|$, the norms themselves are a Cauchy sequences of real numbers, in particular a convergent sequence. Now for fixed \underline{u} , we have $\|T \underline{u}\|_V = \lim_{n \rightarrow \infty} \|T_n \underline{u}\|_V$. We have the pointwise bound $\|T_n \underline{u}\| \leq \|T_n\| \|\underline{u}\|_U$. Passing to the limit we find

$$\|T \underline{u}\|_V \leq \left(\lim_{n \rightarrow \infty} \|T_n\| \right) \|\underline{u}\|_U$$

so T is bounded. Finally, given ε let N be such that if $m, n \geq N$ then $\|T_n - T_m\| \leq \varepsilon$. Then for any $\underline{u} \in U$,

$$\|T_n \underline{u} - T_m \underline{u}\| \leq \|T_n - T_m\| \|\underline{u}\|_U \leq \varepsilon \|\underline{u}\|_U.$$

Letting $m \rightarrow \infty$ and using the continuity of the norm, we get that if $n \geq N$ then

$$\|T_n \underline{u} - T \underline{u}\| \leq \varepsilon \|\underline{u}\|_U.$$

Since \underline{u} was arbitrary this shows that $\|T_n - T\| \leq \varepsilon$ for $n \geq N$ and we are done. \square

EXAMPLE 184. Let K be a compact space. Then $C(K)$, the space of continuous functions on K , is complete wrt $\|\cdot\|_\infty$.

PROOF. Continuous functions on a compact space are bounded. Let $\{f_n\}_{n=1}^\infty \subset C(K)$ be a Cauchy sequence. Then for fixed $x \in X$, $\{f_n(x)\}_{n=1}^\infty \subset \mathbb{C}$ is a Cauchy sequence, hence convergent to some $f(x) \in \mathbb{C}$. To see the convergence is in the norm, give $\varepsilon > 0$ let N be such that $\|f_n - f_m\|_\infty \leq \varepsilon$ for $n, m \geq N$. Then for any x ,

$$|f_n(x) - f_m(x)| \leq \varepsilon.$$

Letting $m \rightarrow \infty$ we find for all $n \geq N$ that $|f_n(x) - f(x)| \leq \varepsilon$, that is

$$\|f_n - f\|_\infty \leq \varepsilon.$$

Finally, we need to show that f is continuous. Given $x \in X$ and $\varepsilon > 0$ let N be as above and let $n \geq N$. For any x , the continuity of f_n gives a neighbourhood of x where $|f_n(x) - f_n(y)| \leq \varepsilon$. Then

$$|f(x) - f(y)| \leq |f(x) - f_n(x)| + |f_n(x) - f_n(y)| + |f_n(y) - f(y)| \leq 3\varepsilon$$

in that neighbourhood, so f is continuous at x . \square

EXERCISE 185. Generalize this example:

- (1) Show that $\ell^\infty(X)$ is complete for any set X .

(2) For a general (topological) space X , show that $C_b(X) = C(X) \cap \ell^\infty(X)$ is complete with respect to the supremum norm.

Now fix a complete normed space V .

(3) For a set X write $\ell^\infty(X; V)$ for the space of bounded functions $X \rightarrow V$. Then $\ell^\infty(X; V)$ is complete.

(4) $C_b(X; V) = C(X; V) \cap \ell^\infty(X; V)$ is complete.

3.4.2. Series of vectors and matrices (Lecture 27).

Fix a complete normed space V .

DEFINITION 186. Say the series $\sum_{n=1}^\infty v_n$ converges *absolutely* if $\sum_{n=1}^\infty \|v_n\|_V < \infty$.

PROPOSITION 187. If $\sum_{n=1}^\infty v_n$ converges absolutely it converges, and $\|\sum_{n=1}^\infty v_n\|_V \leq \sum_{n=1}^\infty \|v_n\|_V$.

PROOF. Standard. □

EXAMPLE 188 (Exponential series). Let V be a complete normed space and let $T \in \text{End}_b(V)$. Then $\exp(T) \stackrel{\text{def}}{=} \sum_{n=0}^\infty \frac{1}{n!} T^n$ converges absolutely.

Another instance of this phenomenon:

THEOREM 189 (Weierstrass's M -test). Let X be a (topological) space, $f_n: X \rightarrow V$ continuous. Suppose that we have M_n such that $\|f_n(x)\|_V \leq M_n$ holds for all $x \in X$. Suppose that $M = \sum_{n=1}^\infty M_n < \infty$. Then $\sum_n f_n$ converges uniformly to a continuous function $F: X \rightarrow V$.

PROOF. By the Proposition this amounts to showing that the space $C_b(X, V)$ (continuous functions $X \rightarrow V$ with $\|f(x)\|_V$ bounded) is complete with respect to the supremum norm $\|f\|_\infty = \sup\{\|f(x)\|_V : x \in X\}$. □

We will apply this to power series of matrices.

EXAMPLE 190. Let $\|\cdot\|$ be some operator norm on $M_n(\mathbb{R})$, and let $A \in M_n(\mathbb{R})$. For $0 < R < \frac{1}{\|A\|}$ (any $R > 0$ if $A = 0$) and $z \in \mathbb{C}$ with $|z| \leq R$ consider the series

$$\sum_{n=0}^{\infty} z^n A^n.$$

We have $\|A^n\| \leq \|A\|^n$ (operator norm!) so that $\|z^n A^n\| \leq (R\|A\|)^n$. Since $\sum_{n=0}^\infty (R\|A\|)^n$ converges, we see that our series converges and the sum is continuous in z (and in A). Taking the union we get convergence in $|z| < \frac{1}{\|A\|}$. The limit is $(\text{Id} - zA)^{-1}$ (incidentally showing this is invertible).

REMARK 191. In fact, the radius of convergence is $\frac{1}{\rho(A)}$.

3.4.3. Vector-valued limits and derivatives.

We recall facts about vector-valued limits.

LEMMA 192 (Limit arithmetic). Let U, V, W be normed spaces. Let $u_i(x): X \rightarrow U$, $\alpha_i(x): X \rightarrow F$, $T(x): X \rightarrow \text{Hom}_b(U, V)$, $S(x): X \rightarrow \text{Hom}_b(V, W)$. Then, in each case supposing the limits on the right exist, the limits on the left exist and equality holds:

- (1) $\lim_{x \rightarrow x_0} (\alpha_1(x)u_1(x) + \alpha_2(x)u_2(x)) = (\lim_{x \rightarrow x_0} \alpha_1(x)) (\lim_{x \rightarrow x_0} u_1(x)) + (\lim_{x \rightarrow x_0} \alpha_2(x)) (\lim_{x \rightarrow x_0} u_2(x))$.
- (2) $\lim_{x \rightarrow x_0} T(x)u(x) = (\lim_{x \rightarrow x_0} T(x)) (\lim_{x \rightarrow x_0} u(x))$.
- (3) $\lim_{x \rightarrow x_0} S(x)T(x) = (\lim_{x \rightarrow x_0} S(x)) (\lim_{x \rightarrow x_0} T(x))$.

PROOF. Same as in \mathbb{R} , replacing $|\cdot|$ with $\|\cdot\|_V$. □

We can also differentiate vector-valued functions (see Math 320 for details)

DEFINITION 193. Let $X \subset \mathbb{R}^n$ be open. Say that $f: X \rightarrow V$ is *strongly differentiable* at x_0 if there is a bounded linear map $L: \mathbb{R}^n \rightarrow V$ such that

$$\lim_{h \rightarrow 0} \frac{\|f(x_0 + h) - f(x_0) - Lh\|_V}{\|h\|_{\mathbb{R}^n}} = 0.$$

In that case we write $Df(x_0)$ for L .

It is clear that differentiability at x_0 implies continuity at x_0 .

LEMMA 194 (Derivatives). Let U, V, W be normed spaces. Let $\underline{u}_i(x): X \rightarrow U$, $T(x): X \rightarrow \text{Hom}_b(U, V)$, $S(x): X \rightarrow \text{Hom}_b(V, W)$ be differentiable at x_0 . Then the derivatives on the left exist and take the following values:

- (1) $D(\underline{u}_1 + \underline{u}_2)(x_0) = D\underline{u}_1(x_0) + D\underline{u}_2(x_0)$.
- (2) $D(T\underline{u})(x_0)(\underline{h}) = (DT(x_0)(\underline{h}) \cdot \underline{u}(x_0)) + T(x_0) \cdot D\underline{u}(x_0)(\underline{h})$.
- (3) $D(ST)(x_0)(\underline{h}) = (DS(x_0)(\underline{h}) \cdot T(x_0)) + (S(x_0) \cdot DT(x_0)(\underline{h}))$.

PROOF. Same as in \mathbb{R} , replacing $|\cdot|$ with $\|\cdot\|_V$. □

3.5. The exponential series (Lecture 28)

We apply Weierstrass's M -test to *power series*.

THEOREM 195. Let X be a (topological) space, $f_n: X \rightarrow V$ continuous. Suppose that we have M_n such that $\|f_n(x)\|_V \leq M_n$ holds for all $x \in X$ with $M = \sum_{n=1}^{\infty} M_n < \infty$. Then $\sum_n f_n$ converges uniformly to a continuous function $F: X \rightarrow V$ and $\|F(x)\|_V \leq M$ for all $x \in X$.

COROLLARY 196. Let V be a complete normed space, and let $\sum_n a_n z^n$ be a power series with radius of convergence R . Then for any $A \in \text{End}_b(V)$, $\sum_n a_n A^n$ converges absolutely if $\|A\| < R$, uniformly in $\{\|A\| \leq R - \varepsilon\}$

PROOF. Let $X = V = \text{End}_b(V)$, $f_n(A) = a_n A^n$, so that $\|f_n(A)\| \leq |a_n| \|A\|^n$. For $T < R$ we have $\sum_n |a_n| T^n < \infty$ and hence uniform convergence in $\{\|A\| \leq T\}$. □

We therefore fix a normed space V , and plug matrices $A \in \text{End}_b(V)$ into power series.

EXAMPLE 197. $\exp(A) = \sum_k \frac{A^k}{k!}$ converges everywhere.

REMARK 198. We'll look at two kinds of matrix-valued series:

- (1) Power series with matrix coefficients: $f(t) = \sum_{n=0}^{\infty} A_n t^n$. Here, t is a scalar and $A_n \in \text{End}_b(V)$.
- (2) Plugging in matrices into power series: given $f(z) = \sum_{n=0}^{\infty} a_n z^n$ set $f(A) = \sum_{n=0}^{\infty} a_n A^n$.

3.5.1. Basic properties.

LEMMA 199. $\exp(tA)\exp(sA) = \exp((t+s)A)$.

PROOF. The series converge absolutely, so the product converges in any order. We thus have

$$\begin{aligned}
\exp(tA)\exp(sA) &= \left(\sum_{k=0}^{\infty} \frac{(tA)^k}{k!} \right) \left(\sum_{l=0}^{\infty} \frac{(sA)^l}{l!} \right) = \sum_{k,l} \frac{t^k s^l A^{k+l}}{k!l!} \\
&= \sum_{m=0}^{\infty} \sum_{k+l=m} \frac{t^k s^l A^{k+l}}{k!l!} = \sum_{m=0}^{\infty} \frac{A^m}{m!} \sum_{k+l=m} \frac{m!}{k!l!} t^k s^l \\
&= \sum_{m=0}^{\infty} \frac{A^m}{m!} (t+s)^m = \exp((t+s)A).
\end{aligned}$$

□

Recap: multiplication of absolutely convergent series.

LEMMA 200. *Let $\sum_n \underline{u}_n$ converge absolutely, Then it converges in any reordering and its sum is unchanged.*

PROOF. Let $\sigma \in S_{\mathbb{N}}$. Given $N > 0$ let $K > \max \sigma^{-1}(\{0, 1, \dots, N\})$. Then

$$\begin{aligned}
\left\| \sum_{k=0}^K \underline{u}_{\sigma(k)} - \sum_{n=0}^N \underline{u}_n \right\| &= \left\| \sum_{\substack{0 \leq k \leq K \\ \sigma(k) > N}} \underline{u}_{\sigma(k)} \right\| \\
&\leq \sum_{\substack{0 \leq k \leq K \\ \sigma(k) > N}} \|\underline{u}_{\sigma(k)}\| \\
&\leq \sum_{n > N} \|\underline{u}_n\|.
\end{aligned}$$

Now given $\varepsilon > 0$ let N be large enough such that $\sum_{n > N} \|\underline{u}_n\| < \varepsilon$ (exists by absolute convergence). Then for K large enough as above,

$$\begin{aligned}
\left\| \sum_{k=0}^K \underline{u}_{\sigma(k)} - \sum_{n=0}^{\infty} \underline{u}_n \right\| &\leq \left\| \sum_{n > N} \underline{u}_n \right\| + \sum_{n > N} \|\underline{u}_n\| \\
&\leq 2 \sum_{n > N} \|\underline{u}_n\| \leq 2\varepsilon.
\end{aligned}$$

□

PROPOSITION 201. *Let $A = \sum_n a_n$, $B = \sum_m b_m$ be convergent series of positive real numbers. Then $\sum_{n,m} a_n b_m$ converges to AB .*

PROOF. Let $S \subset \mathbb{N}^2$ be finite so that $\sum_{(n,m) \in S} a_n b_m$ is a partial sum. Then for N large enough we have $S \subset \{0, 1, \dots, N\}^2$ so that

$$\sum_{(m,n) \in S} a_n b_m \leq \sum_{0 \leq n, m \leq N} a_n b_m = \left(\sum_{n \leq N} a_n \right) \left(\sum_{m \leq N} b_m \right) \leq AB$$

and the series converges. To evaluate the limit it's enough to note that the subsequence of partial sums

$$\sum_{0 \leq n, m \leq N} a_n b_m = \left(\sum_{n \leq N} a_n \right) \left(\sum_{m \leq N} b_m \right)$$

evidently converges to AB . □

THEOREM 202. *Let $A = \sum_{n=0}^{\infty} A_n$, $B = \sum_{m=0}^{\infty} B_m$ be absolutely convergent ($A_n \in \text{Hom}_b(V, W)$, $B_m \in \text{Hom}_b(U, V)$). Then $\sum_{m, n \geq 0} A_n B_m$ converges absolutely to AB .*

PROOF. Since $\|A_n B_m\| \leq \|A_n\| \|B_m\|$ absolute convergence follows from the Proposition and the convergence of $\sum_n \|A_n\|$ and $\sum_m \|B_m\|$. To evaluate the sum we again take the “square” partial sums:

$$\sum_{0 \leq n, m \leq N} A_n B_m = \left(\sum_{n \leq N} A_n \right) \left(\sum_{m \leq N} B_m \right) \xrightarrow{N \rightarrow \infty} AB.$$

□

3.5.2. Differentiation and application to constant-coefficient differential equations.

COROLLARY 203. $\frac{d}{dt} \exp(tA) = A \exp(tA) = \exp(tA)A$.

PROOF. At $t = 0$ we have $\frac{\exp(hA) - \text{Id}}{h} = A + \sum_{k=1}^{\infty} \frac{h^k}{(k+1)!} A^{k+1}$ and

$$\left\| \sum_{k=1}^{\infty} \frac{h^k}{(k+1)!} A^{k+1} \right\| \leq \sum_{k=1}^{\infty} \frac{|h|^k}{(k+1)!} \|A\|^{k+1} \leq \frac{\exp(|h| \|A\|) - 1 - \|A\| |h|}{|h|} \xrightarrow{h \rightarrow 0} 0.$$

In general we have

$$\frac{\exp((t+h)A) - \exp(tA)}{h} = \exp(tA) \frac{\exp(hA) - \text{Id}}{h} \xrightarrow{h \rightarrow 0} \exp(tA)A.$$

That $A \exp(tA) = \exp(tA)A$ follows from considering partial sums. □

Consider the system of differential equations

$$\begin{cases} \frac{d}{dt} \underline{v}(t) = A \underline{v}(t) \\ \underline{v}(0) = \underline{v}_0 \end{cases}$$

where A is a bounded map.

PROPOSITION 204. *The system has the unique solution $\underline{v}(t) = \exp(At) \underline{v}_0$.*

PROOF. We saw $\frac{d}{dt} \exp(At) \underline{v}_0 = A (\exp(At) \underline{v}_0)$. Conversely, suppose $\underline{v}(t)$ is any solution. Then

$$\begin{aligned} \frac{d}{dt} \left(e^{-At} \underline{v}(t) \right) &= \left(e^{-At} (-A) \right) (\underline{v}(t)) + \left(e^{-At} \right) (A \underline{v}(t)) \\ &= e^{-At} (-A + A) \underline{v}(t) = 0. \end{aligned}$$

It remains to prove: □

LEMMA 205. *Let $f: [0, 1] \rightarrow V$ be differentiable. If $f'(t) = 0$ for all t then f is constant.*

PROOF. Suppose $f(t_0) \neq f(0)$. Let $\varphi \in V'$ be a *bounded* linear functional such that $\varphi(f(t_0) - f(0)) \neq 0$. Then $\varphi \circ f: [0, 1] \rightarrow \mathbb{R}$ is differentiable and its derivative is 0:

$$\lim_{h \rightarrow 0} \frac{\varphi(f(t+h)) - \varphi(f(t))}{h} = \lim_{h \rightarrow 0} \varphi \left(\frac{f(t+h) - f(t)}{h} \right) = \varphi \left(\lim_{h \rightarrow 0} \frac{f(t+h) - f(t)}{h} \right) = \varphi(f'(t)).$$

But $(\varphi \circ f)(t_0) - (\varphi \circ f)(0) = \varphi(f(t_0) - f(0)) \neq 0$, a contradiction. \square

REMARK 206. If V is finite-dimensional, every linear functional is bounded. If V is infinite-dimensional the existence of φ is a serious fact (“Hahn–Banach Theorem”)

Now consider a *linear ODE with constant coefficients*:

$$\begin{cases} \frac{d^n}{dt^n} u(t) = \sum_{k=0}^{n-1} a_k u^{(k)}(t) \\ u^{(k)}(0) = w_k \end{cases} \quad 0 \leq k \leq n-1.$$

We solve this system via the auxiliary vector

$$\underline{v}(t) = \left(u(t), u'(t), \dots, u^{(n-1)}(t) \right).$$

We then have

$$\frac{d\underline{v}(t)}{dt} = A\underline{v}$$

where A is the *companion matrix*

$$A = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 0 \\ a_0 & a_1 & \dots & a_{n-2} & a_{n-1} \end{pmatrix}.$$

(companion to the polynomial $x^n - \sum_{k=0}^{n-1} a_k x^k$). It follows that

$$\underline{v}(t) = e^{At} \underline{w}.$$

Idea: bring A to Jordan form so easier to take exponential.

3.6. Invertibility and the resolvent (Lecture 29)

Say we have a matrix A we'd like to invert. Idea: write $A = D + E$ where we know to invert D . Then $A = D(I + D^{-1}E)$, so if $\|D^{-1}E\| < 1$ we have

$$(I + D^{-1}E)^{-1} = \sum_{n=0}^{\infty} (-D^{-1}E)^n$$

and

$$A^{-1} = \sum_{n=0}^{\infty} (-D^{-1}E)^n D^{-1}$$

(in particular, A is invertible).

3.6.1. Application: Gauss-Seidel and Jacobi iteration.

3.6.2. Application: the resolvent. Let V be a complete normed space. Let T be an operator on V . Define the *resolvent set* of T to be the set of $z \in \mathbb{C}$ for which $T - z\text{Id}$ has a bounded inverse. Define the *spectrum* $\sigma(T)$ to be the complement of the resolvent set. This contains the actual eigenvalues (λ such that $\text{Ker}(T - \lambda)$ is non-trivial) but also λ where $T - \lambda$ is not surjective, and λ where an inverse to $T - \lambda$ exists but is unbounded).

THEOREM 207. *The resolvent set is open, and the function (“resolvent function”) $\rho(T) \rightarrow \text{End}_b(V)$ given by $z \mapsto R(z) = (z\text{Id} - T)^{-1}$ is holomorphic.*

PROOF. Suppose $z_0 - T$ has a bounded inverse. We need to invert $z - T$ for z close to z_0 . Indeed, if $|z - z_0| < \frac{1}{\|(z_0 - T)^{-1}\|}$ then

$$- \sum_{n=0}^{\infty} (T - z_0)^{n+1} (z - z_0)^n$$

converges and furnishes the requisite inverse. It is evidently holomorphic in z in the indicated ball. \square

EXAMPLE 208. Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with nice boundary, $\Delta = \frac{d^2}{dx^2} + \frac{d^2}{dy^2}$ the Laplace operator (say defined on $f \in C^\infty(\Omega)$ vanishing on the boundary). Then Δ is unbounded, but its resolvent is nice. For example, $R(i\varepsilon)$ only has eigenvalues. It follows that the spectrum of Δ consists of eigenvalues, that is for $\lambda \in \sigma(\Delta)$ there is $f \in L^2(\Omega)$ with $\Delta f = \lambda f$ (and $f \in C^\infty$ by elliptic regularity).

3.7. Holomorphic calculus

DEFINITION 209. Let $f(z) = \sum_n a_n z^n$. Define $f(A) = \sum_{n=0}^{\infty} a_n A^n$.

LEMMA 210. $Sf(A)S^{-1} = f(SAS^{-1})$.

PROPOSITION 211. $(f \circ g)(A) = f(g(A))$ if it all works.

THEOREM 212. $\det(\exp(A)) = \exp(\text{Tr}(A))$.

CHAPTER 4

Vignettes

Sketches of applications of linear algebra to group theory.

Key Idea: *linearization* – use linear tools to study non-linear objects.

4.1. The exponential map and structure theory for $\mathrm{GL}_n(\mathbb{R})$

Our goal is to understand the (topologically) closed subgroups of $G = \mathrm{GL}_n(\mathbb{R})$.

Idea: to a subgroup H assign the logarithms of the elements of H . If H was commutative this would be a subspace.

DEFINITION 213. $\mathrm{Lie}(H) = \{X \in M_n(\mathbb{R}) \mid \forall t : \exp(tX) \in H\}$.

REMARK 214. Clearly this is invariant under scaling. In fact, enough to take small t , and even just a sequence of t tending to zero (since $\{t \mid \exp(tX) \in H\}$ is a closed subgroup of \mathbb{R}).

THEOREM 215. $\mathrm{Lie}(H)$ is a subspace of $M_n(\mathbb{R})$, closed under $[X, Y]$.

PROOF. For $t \in \mathbb{R}$ and $m \in \mathbb{Z}_{\geq 1}$, $(\exp(\frac{tX}{m}) \exp(\frac{tY}{m}))^m = \left(\mathrm{Id} + \frac{tX+tY}{m} + O(\frac{1}{m^2})\right)^m \xrightarrow{m \rightarrow \infty} \exp(tX + tY)$.
Thus If $X, Y \in \mathrm{Lie}(H)$ then also $X + Y \in \mathrm{Lie}(H)$. \square

- Classify subgroups of G containing A by action on Lie algebra and finding eigenspaces.

4.2. Representation Theory of Groups

EXAMPLE 216 (Representations). (1) Structure of $\mathrm{GL}_n(\mathbb{R})$: let A act on $M_n(\mathbb{R})$.

(2) M manifold, G acting on M , thus acting on $H_k(M)$ and $H^k(M)$.

(3) *Angular momentum*: $O(3)$ acting by rotation on $L^2(\mathbb{R}^3)$.

Bibliography