

---

# Positive Rational Solutions to $x^y = y^{mx}$ : A Number-Theoretic Excursion

---

Michael A. Bennett and Bruce Reznick

---

**1. PROLOGUE.** Late in the last millennium, the second author ran a seminar course for undergraduates that was intended to introduce them to problem-solving and question-asking in the context of mathematical research. He led them through the classic “difficult” equation

$$x^y = y^x \quad (x, y > 0), \quad (1)$$

whose solution is much easier than one would think at first glance. Solutions were sought, first over  $\mathbb{R}$ , then over  $\mathbb{Z}$ , and finally over  $\mathbb{Q}$ . In the context of this seminar, it was natural to consider the variant equation

$$x^y = y^{2x} \quad (x, y > 0), \quad (2)$$

which does not seem to have appeared in the literature. It turns out that there are solutions to (2) that do not fit the well-known parametric pattern of (1) (see (5)). For example,

$$x = \left(\frac{4}{5}\right)^{128}, \quad y = \left(\frac{4}{5}\right)^{125} \quad (3)$$

is a solution to (2). This preposterous fact is trivial to verify: simply substitute (3) into (2), take logs, and transpose:

$$\frac{2x}{y} = 2 \left(\frac{4}{5}\right)^3 = \frac{128}{125} = \frac{\log x}{\log y}.$$

As we shall see, the ultimate explanation for this identity is that  $2 \cdot 4^3 = 5^3 + 3$ . Upon discovering (3), the second author realized he was in over his head and contacted the first author. This paper is the result.

**2.  $x^y = y^x$ .** First, we review the familiar, but beautiful solution to (1), reserving historical references to the last paragraph of the section. We acknowledge the solutions  $x = y$  and now let  $y = tx$ , where  $t \neq 1$ , so that

$$x^y = y^x \iff (x^t)^x = (tx)^x \iff x^t = tx \iff x = t^{\frac{1}{t-1}}. \quad (4)$$

The positive real solutions to (1) are thus

$$(x, y) = (u, u), \quad (x, y) = (t^{\frac{1}{t-1}}, t^{\frac{t}{t-1}}). \quad (5)$$

(We might equally well have set  $y = x^r$  in (1) and drawn essentially the same conclusion.) Since (1) implies that  $x^{1/x} = y^{1/y}$  and since  $f(u) = u^{1/u}$  increases on  $(1, e)$  and decreases on  $(e, \infty)$ , for each  $x$  in  $(1, e)$  there is exactly one  $y$  in  $(e, \infty)$  so that (1)

holds. There are at most two solutions to (1), one of which is  $x = y$ . In particular, the only integer solution to (1) with  $x \neq y$  is  $x = 2$  and  $y = 4$  (or vice versa).

Euler already noted that if  $t = 1 + n^{-1}$  for integral  $n$ , then (4) gives a rational solution to (1); namely,

$$x_n = \left(1 + \frac{1}{n}\right)^n, \quad y_n = \left(1 + \frac{1}{n}\right)^{n+1}. \tag{6}$$

(Observe that, as  $n \rightarrow \infty$ , we have  $t \rightarrow 1$  and  $(x_n)$  and  $(y_n)$  increase and decrease monotonically to  $e$ , as is familiar from calculus.)

To show that these are the only rational solutions, we need an elementary lemma, whose proof, relying upon the Fundamental Theorem of Arithmetic, we omit:

**Lemma 1.** *Suppose  $a, b, m$ , and  $n$  are integers satisfying  $\gcd(a, b) = \gcd(m, n) = 1$  and  $b, n \neq 0$ . Then  $(m/n)^{a/b}$  is rational if and only if  $m$  and  $n$  are  $|b|$ th powers of integers.*

Let us now proceed to find all rational solutions to (1) with  $x \neq y$ . By symmetry, we may assume that  $y > x$ , so  $t > 1$ . If  $x$  and  $y$  are rational, then so is  $t = y/x$ . Write  $t$  in lowest terms as

$$t = \frac{p}{q} := \frac{q+d}{q}$$

( $d, q > 0$ ), so that  $1/(t-1) = q/d$  and  $t/(t-1) = (q+d)/q$ . With this substitution, we have

$$x = \left(\frac{q+d}{q}\right)^{q/d}, \quad y = \left(\frac{q+d}{q}\right)^{q/d+1}.$$

Since  $\gcd(d, q) = \gcd(q, q+d) = \gcd(d, q+d) = 1$ , the integers  $q$  and  $q+d$  must both be  $d$ th powers (Lemma 1). This causes no problem when  $d = 1$ , of course, and, setting  $q = n$ , we recover (6). Suppose that  $d > 1$ , and write  $q = a^d$ ,  $q+d = b^d$  for positive integers  $a$  and  $b$  with  $a < b$ , so that  $b^d - a^d = d$ . Observe that, for such positive integers  $a$  and  $b$ ,

$$b^d - a^d \geq (a+1)^d - a^d \geq 1 + da \geq 1 + d > d,$$

so there are no solutions with  $d > 1$ . Finally, we remark that if  $n = -r$  is allowed to be negative in (6), then  $(x_{-r}, y_{-r}) = (y_{r-1}, x_{r-1})$ :

$$\left(1 - \frac{1}{r}\right)^{-r} = \left(1 + \frac{1}{r-1}\right)^r, \quad \left(1 - \frac{1}{r}\right)^{-r+1} = \left(1 + \frac{1}{r-1}\right)^{r-1}.$$

Thus, by removing the restriction that  $n$  be positive, we can eliminate the constraint  $y > x$  in (6).

A historical discussion of  $x^y = y^x$  occupies a paragraph in chapter 23 of Dickson [3, p.687]. The first reference to (1) was in a letter from D. Bernoulli to C. Goldbach dated June 29, 1728. Bernoulli asserts, without proof (see [2, p.263]), that this equation has only one solution in positive integers, and infinitely many rational solutions. The first person to write about (1) in detail was Euler (see [4, pp. 340–341]). Euler made

the substitution  $y = tx$  and solved the equation over  $\mathbb{R}_+$  and  $\mathbb{Z}_+$ , and he presented the rational solutions (6), without claiming that they were the only ones.

Dickson mentions other writers who covered the same ground, and adds that “\*A. Flechsenhaar and R. Schimmack discussed the rational solutions” [3, p. 687]. The cited papers appeared in 1911 and 1912 in the expository journal *Unterrichtsblätter für Mathematik und Naturwissenschaften*. Dickson writes: “The reports in Chapters XI–XXVI have been checked by the original papers in case they are to be found in Chicago” [3, p. xxi]; the asterisk means that the paper was “not available for report” [3, p. xxii]. We were able to examine [6] at the magnificent Mathematics Library of the University of Illinois at Urbana-Champaign and can report that Flechsenhaar appears to deserve credit as the first author to solve (1) over the positive rationals. In 1967, S. Hurwitz [9] gave the first readily accessible analysis of the rational solutions; subsequent work on this equation, including generalizations to algebraic solutions, can be found in [8], [14], [17], [18], [21], and [22].

**3.  $x^y = y^{mx}$ .** Let us now consider the generalization of Euler’s equation to

$$x^y = y^{mx} \tag{7}$$

where  $m > 1$  is a fixed positive integer. We again restrict our attention to positive solutions  $(x, y)$ . If  $x = 1$  or  $y = 1$ , then necessarily  $(x, y) = (1, 1)$ . Supposing that  $x, y \neq 1$  and taking logarithms in (7), we find that

$$\frac{m \log y}{y} = \frac{\log x}{x},$$

so  $x \neq y$ . Write  $y = x^r$  with  $x \neq 1$  and  $r \neq 1$ . Then  $x^{r-1} = mr$ . Therefore, the positive real solutions to (7) are given by  $x = y = 1$  and

$$x = (mr)^{\frac{1}{r-1}}, \quad y = (mr)^{\frac{r}{r-1}}.$$

We now restrict our attention to positive rational solutions. Since  $y/x = mr$ , it follows that  $r > 0$ ; further, since  $x$  and  $y$  are rational, we have  $r$  in  $\mathbb{Q}$ . Let us write  $r = a/b$  where  $a$  and  $b$  are natural numbers with  $\gcd(a, b) = 1$ , and set  $k = |a - b|$ . To have  $x$  in  $\mathbb{Q}$ , we require

$$x = \left(\frac{ma}{b}\right)^{\frac{b}{a-b}} \in \mathbb{Q}.$$

Suppose that  $\gcd(b, m) = d$ , and write  $b = db'$  and  $m = dm'$ . Then  $\gcd(am', b') = 1$ , so we need  $am'$  and  $b'$  to be  $k$ th powers of integers. If

$$am' = u^k, \quad b' = v^k,$$

it follows that

$$|u^k - mv^k| = |am' - mb'| = \left| \frac{am}{d} - \frac{mb}{d} \right| = \frac{m}{d} \cdot k = m'k. \tag{8}$$

For a given positive integer  $m$ , we classify the set  $S(m)$  of positive rational solutions to (7) as follows: write

$$S(m) = \bigcup_{k=0}^{\infty} S_k(m),$$

where  $S_k(m)$  represents the set of solutions  $(x, y)$  corresponding to equation (8) for  $m'$  ranging over all positive integral divisors of  $m$ . Here,  $S_0(m)$  denotes the solutions with  $x = y$  (i.e., just  $(x, y) = (1, 1)$  for  $m > 1$ ).

The remainder of this paper is devoted to analyzing  $S(m)$ . As a consequence, we will show how this set may be completely characterized for any given  $m$ . In case  $m = 2$  or 3 (where all features of interest for general  $m$  may in fact be observed), we have the following results:

**Theorem 2.** *If  $x$  and  $y$  are positive rational numbers for which  $x^y = y^{2x}$ , then either*

$$(a) \ x = \left(2 + \frac{2}{n}\right)^n \text{ and } y = \left(2 + \frac{2}{n}\right)^{1+n} \text{ for } n \text{ in } \mathbb{Z}, n \neq 0, -1;$$

or

$$(b) \ (x, y) = (1, 1), (2, 16), \text{ or } \left(\left(\frac{4}{5}\right)^{128}, \left(\frac{4}{5}\right)^{125}\right).$$

**Theorem 3.** *If  $x$  and  $y$  are positive rational numbers for which  $x^y = y^{3x}$ , then either*

$$(a) \ x = \left(3 + \frac{3}{n}\right)^n \text{ and } y = \left(3 + \frac{3}{n}\right)^{1+n} \text{ for } n \text{ in } \mathbb{Z}, n \neq 0, -1;$$

or

$$(b) \ x = \left(\frac{3w_n}{v_n}\right)^{v_n^2} \text{ and } y = \left(\frac{3w_n}{v_n}\right)^{3w_n^2} \text{ for } n = 0, 1, 2, \dots;$$

or

$$(c) \ x = \left(\frac{w_n}{v_n}\right)^{3w_n^2} \text{ and } y = \left(\frac{w_n}{v_n}\right)^{v_n^2} \text{ for } n = 0, 1, 2, \dots$$

In (b) and (c),  $v_n$  and  $w_n$  are the integers defined by

$$v_n + w_n\sqrt{3} = (1 + \sqrt{3})(2 + \sqrt{3})^n.$$

Our proofs of Theorems 2 and 3 (perhaps rather surprisingly) involve techniques from transcendental number theory and Diophantine approximation. Moreover, they provide us with an opportunity to illustrate a fairly diverse grab bag of methods for solving Diophantine problems.

**4. THE CASES  $k = 1$  AND  $k = 2$ .** The set  $S_1(m)$  is easily computed. Indeed, we immediately find that  $k = 1$  implies that either

$$x = \left(m + \frac{m}{n}\right)^n, \quad y = \left(m + \frac{m}{n}\right)^{1+n}$$

or

$$x = \left(m - \frac{m}{n}\right)^{-n}, \quad y = \left(m - \frac{m}{n}\right)^{1-n}$$

for  $n$  a positive integer (with  $n \geq 2$  in the latter case). We note that these solutions correspond to the parametrized family (6) of solutions to Euler's original equation and provide us with part (a) of Theorems 2 and 3.

If  $k \geq 2$ , the situation becomes more interesting, though the set  $S_2(m)$  is still not too difficult to describe: it is either empty or infinite. The following lemmata provide sufficient conditions for the former to occur. As usual, for  $x$  in  $\mathbb{N}$ , let  $v_2(x)$  be the largest integer such that  $2^{v_2(x)}$  divides  $x$ .

**Lemma 4.** *If  $v_2(m)$  is odd and  $k$  is even, then  $S_k(m)$  is empty.*

*Proof.* We have  $\gcd(a, b) = 1$ , and since  $k = |a - b|$  is even,  $a$  and  $b$  must both be odd. It follows that  $d = \gcd(b, m)$  is odd as well. Write, as before,  $b = db'$ ,  $m = dm'$ ,  $am' = u^k$ ,  $b' = v^k$ . Then  $v_2(m) = v_2(d) + v_2(m')$  is odd, hence so is  $v_2(am') = v_2(u^k) = kv_2(u)$ , which contradicts  $k$  being even. ■

**Lemma 5.** *If  $m = 2^{v_2(m)} \cdot m_1$  is a positive integer for which  $m_1 \equiv 1 \pmod{4}$ , then  $S_2(m)$  is empty.*

*Proof.* If  $v_2(m)$  is odd, this is a special case of Lemma 4. Since  $k = 2$  is even, we may conclude as before that  $a$  and  $b$  are both odd. Suppose  $v_2(m)$  is even, say  $v_2(m) = 2t$ . It follows from (8) that

$$u^2 - 2^{2t} m_1 v^2 = \pm 2^{2t+1} \cdot \frac{m_1}{d},$$

where  $u$  and  $v$  are coprime and  $d$  divides  $m$ . Because  $d \mid b$ , it follows that  $d$  is odd, so  $d \mid m_1$ . We thus see that  $2^t \mid u$ , say  $u = 2^t u_1$ , whereby

$$u_1^2 - m_1 v^2 = \pm 2 \cdot \frac{m_1}{d}. \quad (9)$$

Since the right-hand side of this equation is even,  $u_1$  and  $v$  have the same parity and, since  $u$  and  $v$  are coprime, are both necessarily odd. Since  $m_1 \equiv 1 \pmod{4}$ , this implies that the left-hand side of (9) is divisible by 4. Since the right-hand side of this equation is congruent to 2 modulo 4, this yields the desired contradiction. ■

It is a well-known fact that if  $m$  is a positive nonsquare integer and  $c$  is a nonzero integer, then a single solution in positive integers  $x$  and  $y$  to the equation  $x^2 - my^2 = c$  implies the existence of infinitely many such solutions. In fact, one can find a finite collection of pairs of positive integers, say

$$(x_1, y_1), (x_2, y_2), \dots, (x_r, y_r)$$

with  $x_i^2 - my_i^2 = c$  for  $i = 1, 2, \dots, r$ , such that every solution in positive integers  $(x, y)$  to the equation  $x^2 - my^2 = c$  satisfies

$$x + y\sqrt{m} = (x_i + y_i\sqrt{m}) \cdot (u_1 + v_1\sqrt{m})^k, \quad (10)$$

where  $k$  is a nonnegative integer,  $i$  belongs to  $\{1, 2, \dots, r\}$ , and  $(u_1, v_1)$  is the smallest positive integer solution to the equation  $u^2 - mv^2 = 1$ . The integer  $r$  here depends upon  $c$  and, potentially, upon  $m$ .

It follows that if  $S_2(m)$  is not empty, then it is infinite. From the theory of Pell equations (or Fermat-Pell equations if one likes) these correspond to elements of finitely many recurrence sequences (see Nagell [15] for a nice, affordable exposition of such matters). This fact is also a pretty easy consequence of (10). We note that consideration of the case  $m = 39$  demonstrates that the foregoing lemmata do not in fact provide necessary conditions for  $S_2(m)$  to be empty. A routine computation shows that, when  $2 \leq m \leq 50$ ,  $S_2(m)$  is infinite precisely for

$$m \in \{3, 7, 11, 12, 15, 19, 23, 27, 28, 31, 35, 43, 44, 47, 48\}.$$

It is also not hard to provide sufficient conditions for  $S_2(m)$  to be nonempty. An example of such a result is as follows:

**Lemma 6.** *If  $p \equiv 3 \pmod{4}$  and  $p$  is prime, then  $S_2(p)$  is infinite.*

*Proof.* From a venerable (after citing Euler, we can hardly call this old!) result of Petr [16], precisely one of the equations

$$x^2 - py^2 = -2, \quad x^2 - py^2 = 2$$

is solvable in integers  $x$  and  $y$ . This implies, in either case, that  $S_2(p)$  is infinite. ■

We leave to our gentle reader the (rather nontrivial) task of deriving *necessary* conditions for  $S_2(m)$  to be nonempty. In the next section, we observe that the behavior of  $S_k(m)$  is dramatically different when  $k \geq 3$ .

**5. THUE EQUATIONS.** A famous theorem of the Norwegian mathematician Axel Thue [23] asserts that, if  $\theta$  is an algebraic number of degree  $k \geq 3$  and  $\epsilon > 0$ , then the inequality

$$\left| \theta - \frac{x}{y} \right| < \frac{1}{|y|^{k/2+1+\epsilon}}$$

has at most finitely many solutions in integers  $x$  and  $y$  with  $y \neq 0$ . Note that, if  $\theta = \sqrt[k]{m}$ , then we have the algebraic identity

$$|x^k - my^k| = y^k \left| \theta - \frac{x}{y} \right| \cdot \left( \left( \frac{x}{y} \right)^{k-1} + \theta \left( \frac{x}{y} \right)^{k-2} + \dots + \theta^{k-1} \right).$$

If  $m$  is not a perfect  $k$ th power and  $c \neq 0$  is an integer, then it follows that the equation

$$x^k - my^k = c \tag{11}$$

has at most finitely many solutions in integers  $x$  and  $y$ . Such equations are nowadays termed *Thue equations* (a nice exposition of this active area of research may be found in the book of Fel'dman and Nesterenko [5]).

In our situation, this immediately implies that  $S_k(m)$  is finite for each fixed integer  $k \geq 3$ . As we shall see in the next two sections, it is possible to determine each such  $S_k(m)$  “effectively” (and to derive an upper bound for  $k$  that depends solely upon  $m$ ).

**6. LINEAR FORMS IN LOGARITHMS.** The sets  $S_k(m)$  are defined by equalities of the shape

$$u^k - mv^k = \pm \frac{m}{d}k$$

for  $u$  and  $v$  positive integers. In all cases, we thus have

$$|u^k - mv^k| \leq mk.$$

It follows that

$$\left| m^{-1} \left( \frac{u}{v} \right)^k - 1 \right| \leq \frac{k}{v^k}$$

and hence the *linear form in logarithms*  $|k \log(u/v) - \log m|$  is small. A classical result of Gel'fond [7] (extending his work on Hilbert's seventh problem) indicates that, for any given nonzero algebraic numbers  $\alpha_1, \alpha_2, \beta_1$ , and  $\beta_2$  with  $\log \alpha_1$  and  $\log \alpha_2$  linearly independent over the rationals, we have

$$|\beta_1 \log \alpha_1 - \beta_2 \log \alpha_2| \neq 0.$$

Moreover, Gel'fond provides lower bounds for such a form. Applying state-of-the-art versions of these bounds, say those due to Laurent, Mignotte, and Nesterenko [11], we may conclude as in Theorem 3 of Mignotte [13] that, in our situation,  $k < 600$  (if  $m = 2$ ) and, more generally, as in Theorem 2 of [13], that

$$k < 10676 \log m.$$

It follows that

$$\bigcup_{k=3}^{\infty} S_k(m)$$

is finite. In the remaining sections, we describe a strategy for explicitly determining this set and illustrate it in the cases  $m = 2$  and  $m = 3$ .

**7. SOLVING THE REMAINING EQUATIONS.** For small values of  $k$ , it is possible to use standard computational techniques based upon lower bounds for linear forms in logarithms, combined with lattice basis reduction, to solve the Thue equations that occur (a reasonably accessible book that covers this field is Smart [20]). If, however,  $k$  is moderately large, this becomes computationally infeasible due to the difficulty in finding systems of independent units in the ring of integers of  $\mathbb{Q}(\sqrt[k]{m})$ . (For instance, it is an interesting challenge to find the fundamental units in, say,  $\mathbb{Q}(\sqrt[101]{2})$ .)

In our situation, though, we are able to find local obstructions to solvability (i.e., appropriate moduli  $m$  so that the equations have no admissible solutions modulo  $m$ ) for virtually all values of  $k$  under consideration, obviating the need for extensive computations. For the sake of simplicity, let us restrict our attention to the case  $m = 2$  (where, as mentioned previously, we may assume that  $k < 600$ ). Here the equations to be studied are of the shape

$$x^k - 2y^k = \pm 2^\delta k,$$

where  $\delta$  is 0 or 1. Lemma 4 also allows us to suppose that  $k$  is odd. For each such  $k$ , we consider primes of the form  $p = 2nk + 1$  for  $n$  in  $\mathbb{N}$ . For these  $p$ , there are at most  $(2n + 1)^2$  values of  $x^k - 2y^k$  modulo  $p$ . If none of these are congruent to  $\pm k$  or  $\pm 2k$  modulo  $p$ , we have found a local obstruction to solvability and can thus conclude that  $S_k(2)$  is empty. If, for example,  $k = 13$ , we consider the equation(s) modulo 53. Noting that  $x^{13} \equiv 0, \pm 1, \pm 23 \pmod{53}$ , it follows that

$$x^{13} - 2y^{13} \equiv 0, \pm 1, \pm 2, \pm 3, \pm 6, \pm 7, \pm 8, \pm 16, \pm 21, \pm 23, \pm 25 \pmod{53},$$

hence we may conclude that  $S_{13}(2)$  is empty. Similar arguments suffice to eliminate all equations (with  $k < 600$ ) except for

$$\begin{aligned} x^3 - 2y^3 &= \pm 3, & x^3 - 2y^3 &= \pm 6, & x^5 - 2y^5 &= \pm 10, \\ x^7 - 2y^7 &= \pm 7, & x^7 - 2y^7 &= \pm 14, & x^{11} - 2y^{11} &= \pm 22 \end{aligned}$$

(to verify this, the reader may wish to write her own piece of code; we used Maple [12], being careful to note that the “msolve” routine is somewhat unreliable).

Similarly, when  $m = 3$ , we easily deal with all the equations encountered, with the exceptions of

$$x^3 - 3y^3 = \pm 3, \quad x^5 - 3y^5 = \pm 15, \quad x^7 - 3y^7 = \pm 21$$

(though, in this case, we need to treat all  $k$  up to  $k = 11,728$ , including even values).

**8. ENDGAME.** As mentioned in the last section, various techniques from Diophantine approximation may be used to solve these remaining equations. The symbolics package Kash [10], for example, has a built-in Thue solver that can handle all the equations under consideration in a matter of minutes on a Sun Ultra. In any case, nowadays it is a routine matter to verify that, of the equations that have so far evaded our net, only those with  $k = 3$  possess solutions, corresponding to  $5^3 - 2 \cdot 4^3 = -3$ ,  $2^3 - 2 \cdot 1^3 = 6$ , and  $3^3 - 3 \cdot 2^3 = 3$ . By way of example, to solve the Diophantine equations  $|x^3 - 2y^3| = 3$  or  $|x^3 - 2y^3| = 6$ , one can appeal to the inequality

$$|x^3 - 2y^3| \geq \sqrt{|x|},$$

valid for all integers  $x$  and  $y$  (see, for example, [1]).

The equation  $5^3 - 2 \cdot 4^3 = -3$  maps back to the solution  $(x, y)$  to  $x^y = y^{2x}$  given in (3). Similarly,  $2^3 - 2 \cdot 1^3 = 6$  yields the solution  $(x, y) = (2, 16)$ , also to  $x^y = y^{2x}$ . The equation  $3^3 - 3 \cdot 2^3 = 3$ , which potentially yields a solution to  $x^y = y^{3x}$ , leads to  $a = 27$  and  $b = 24$ , contradicting the coprimality of  $a$  and  $b$ .

These arguments suffice to solve  $x^y = y^{mx}$  completely, with a modicum of computation, for all values of  $m$  up to 40 or so. As  $m$  increases, we are faced with the prospect of handling Thue equations of higher and higher degree, an apparently formidable task. We note that  $S_k(m)$  can be nonempty for arbitrarily large  $k$  (for example, this is always the case for  $S_k(2^k - k)$  if  $k$  is odd).

The reader might reasonably wonder about a generalization to  $x^{ny} = y^{mx}$ . This Diophantine equation was studied by Asher Kach, a student in that undergraduate seminar course who is currently a graduate student at the University of Wisconsin-Madison. He and the authors are preparing an article on this subject.

**9. POSTSCRIPT: ON LOCAL-GLOBAL PRINCIPLES.** A standard heuristic employed in the field of Diophantine problems is that an equation should be solvable over  $\mathbb{Q}$  precisely when it is solvable over  $\mathbb{R}$  and over the  $p$ -adic fields  $\mathbb{Q}_p$  for all primes  $p$ . Such *local-global* (or *Hasse*) principles are known to be true in various settings, but false in general.

One of the first instances where local-global principles were shown to fail occurred in a 1942 paper of Skolem [19], where he demonstrated that the equation  $x^3 - 3y^3 = 22$  can be solved modulo  $p^\alpha$  for every prime power  $p^\alpha$  but has no solution over the integers.

In our analysis, we encounter numerous instances for which a like conclusion holds, with equations of degree up to (at least) 19. Such is the case, for example, for the equations

$$\begin{aligned} x^3 - 5y^3 = 15, \quad x^5 - 2y^5 = 10, \quad x^5 - 3y^5 = 15, \quad x^7 - 2y^7 = 7, \\ x^7 - 2y^7 = 14, \quad x^7 - 3y^7 = 21, \quad x^{11} - 2y^{11} = 22. \end{aligned}$$

To prove that the Hasse principle fails on these occasions is an interesting exercise that we leave to our loyal (and, at this stage, perhaps rather fatigued) reader!

**ACKNOWLEDGMENTS.** We thank Karim Belabas, Jeff Lagarias, Gerry Myerson, and the anonymous referees for their insightful comments on an earlier version of this paper. Some computations in this paper were performed by using Maple.

## REFERENCES

1. M. Bennett, Effective measures of irrationality for certain algebraic numbers, *J. Austral. Math. Soc.* **62** (1997) 329–344.
2. D. Bernoulli, *Corresp. Math. Phys.*, vol. 2, P.-H. Fuss, ed., St. Pétersbourg, 1843.
3. L. E. Dickson, *History of the Theory of Numbers*, vol. 2, Carnegie Institute, Washington, 1919; reprinted by Chelsea, New York, 1966.
4. L. Euler, *Introduction to Analysis of the Infinite* (trans. J. D. Blanton), Springer-Verlag, New York, 1990.
5. N. I. Fel'dman, and Yu. V. Nesterenko, *Transcendental Numbers*, Number Theory IV, EMS, Springer-Verlag, New York, 1998.
6. A. Flechsenaar, Über die Gleichung  $x^y = y^x$ , *Unterrichts. für Math.* **17** (1911) 70–73.
7. A. O. Gel'fond, On Hilbert's seventh problem, *Dokl. Akad. Nauk SSSR* **2** (1934) 1–6.
8. A. Hausner, Algebraic number fields and the Diophantine equation  $m^n = n^m$ , this MONTHLY **68** (1961) 856–861.
9. S. Hurwitz, On the rational solutions to  $m^n = n^m$  with  $m \neq n$ , this MONTHLY **74** (1967) 298–300.
10. Kant-Gruppe, KASH, <http://www.math.tu-berlin.de/~kant>, 1999.
11. M. Laurent, M. Mignotte, and Y. Nesterenko, Formes linéaires en deux logarithmes et déterminants d'interpolation, *J. Number Theory* **55** (1995) 285–321.
12. M. B. Monagan et al., *Maple 8 Advanced Programming Guide*, Waterloo Maple, Toronto, 2002.
13. M. Mignotte, A note on the equation  $ax^n - by^n = c$ , *Acta Arith.* **75** (1996) 287–295.
14. G. Mitás, Über die Lösungen der Gleichung  $a^b = b^a$  in rationalen und in algebraischen Zahlen, *Mitt. Math. Gesellsch. Hamburg* **10** (1976) 249–254.
15. T. Nagell, *Introduction to Number Theory*, John Wiley and Sons, New York, 1951; reprinted by Chelsea, New York, 1981.
16. K. Petr, Über die Pellsche Gleichung, *Časopis Pěst. Mat. Fys.* **56** (1927) 57–66.
17. D. Sato, Algebraic solution of  $x^y = y^x$  ( $0 < x < y$ ), *Proc. Amer. Math. Soc.* **31** (1972) 316.
18. ———, A characterization of two real algebraic numbers  $x, y$  such that  $x^y = y^x$ , *Sūgaku* **24** (1972) 223–226.
19. T. Skolem, Unlösbarkeit von Gleichungen, deren entsprechende Kongruenz für jeden Modul lösbar ist, *Avh. Norske Vid. Akad. Oslo* **4** (1942) 1–28.
20. N. Smart, *The Algorithmic Resolution of Diophantine Equations*, Cambridge University Press, Cambridge, 1998.
21. R. M. Sternheimer, A corollary to iterated exponentiation, *Fibonacci Quart.* **23** (1985) 146–148.
22. M. Sved, On the rational solutions of  $x^y = y^x$ , *Math. Mag.* **63** (1990) 30–33.
23. A. Thue, Über Annäherungswerte algebraischer Zahlen, *J. Reine Angew. Math.* **135** (1909) 284–305.

**MIKE BENNETT** studied mathematics at Dalhousie University and the University of British Columbia. He received his Ph.D. in 1993 as a student of David Boyd. After postdoctoral fellowships at Waterloo, Ann Arbor, and the Institute for Advanced Study, he was a faculty member at the University of Illinois, Urbana-Champaign until returning to Vancouver in 2001. He remains an unrepentant number theorist and favors the plucky underdog New York Yankees.

*Department of Mathematics, University of British Columbia, Vancouver, B.C.*  
*bennett@math.ubc.ca*

**BRUCE REZNICK** got his degrees at Caltech (B.S., 1973) and Stanford (Ph.D., 1976) and has been at the University of Illinois, Urbana-Champaign since 1979. He is interested in combinatorial problems in number theory, analysis, algebra, and geometry, often involving polynomials. He continues, somehow, to root for the Chicago Cubs.

*Department of Mathematics, University of Illinois, Urbana, IL 61801*  
*reznick@math.uiuc.edu*