

PROOF OF A DYNAMICAL BOGOMOLOV CONJECTURE FOR LINES UNDER POLYNOMIAL ACTIONS

DRAGOS GHIOCA AND THOMAS J. TUCKER

ABSTRACT. We prove a dynamical version of the Bogomolov conjecture in the special case of lines in \mathbb{A}^m under the action of a map (f_1, \dots, f_m) where each f_i is a polynomial in $\overline{\mathbb{Q}}[X]$ of the same degree.

1. INTRODUCTION

In 1998, Ullmo [Ull98] and Zhang [Zha98] proved the following conjecture of Bogomolov [Bog91].

Theorem 1.1. *Let A be an abelian variety defined over a number field with Néron-Tate height \widehat{h}_{nt} and let W be an irreducible subvariety of A that is not a torsion translate of an abelian subvariety of A . Then there exists an $\epsilon > 0$ such that the set*

$$\{x \in A(\overline{\mathbb{Q}}) \mid \widehat{h}_{\text{nt}}(x) \leq \epsilon\}$$

is not Zariski dense in W .

Earlier, Zhang [Zha95a] had proved a similar result for the multiplicative group \mathbb{G}_m^n . Zhang [Zha95b, Zha06] also proposed a more general conjecture for what he called *polarizable* morphisms; a morphism $\Phi : X \rightarrow X$ on a projective variety X is said to be polarizable if there is an ample line bundle \mathcal{L} on X such that $\Phi^*\mathcal{L} \cong q\mathcal{L}$ for some integer $q > 1$. When a polarizable map Φ is defined over a number field, it gives rise to a canonical height \widehat{h}_Φ with the property that $\widehat{h}_\Phi(\Phi(\alpha)) = q\widehat{h}_\Phi(\alpha)$ for all $\alpha \in X(\overline{\mathbb{Q}})$. Zhang makes the following Bogomolov-type conjecture in this more general context.

Conjecture 1.2. *(Zhang) Let $\Phi : X \rightarrow X$ be a polarizable morphism of a projective variety defined over a number field and let W be a subvariety of X that is not preperiodic under Φ . Then there exists an $\epsilon > 0$ such that the set*

$$\{x \in W(\overline{\mathbb{Q}}) \mid h_\Phi(x) \leq \epsilon\}$$

is not Zariski dense in W .

Date: April 30, 2009.

1991 Mathematics Subject Classification. Primary 14G25; Secondary 37F10, 11C08.

The first author was partially supported by NSERC. The second author was partially supported by NSA Grant 06G-067 and NSF Grant DMS-0801072.

The definition of preperiodicity for varieties here is the same as the usual definition of preperiodicity for points. More precisely, for any quasiprojective variety X , any endomorphism $\Phi : X \rightarrow X$, and any subvariety $V \subset X$, we say that V is Φ -preperiodic if there exists $N \geq 0$, and $k \geq 1$ such that $\Phi^N(V) = \Phi^{N+k}(V)$. Note that when A is an abelian variety and Φ is a multiplication-by- n map (for $n \geq 2$), an irreducible subvariety W is preperiodic if and only if it is a torsion translate of an abelian subvariety of A .

We note that if $X = (\mathbb{P}^1)^n$, and if Φ is given by the coordinatewise action of z^d (for some $d \geq 2$) on X , then Conjecture 1.2 reduces to the result proved by Zhang in [Zha95a].

In this paper, we prove the following special case of Conjecture 1.2.

Theorem 1.3. *Let $f_1, \dots, f_m \in \overline{\mathbb{Q}}[X]$ be polynomials of degree $d > 1$, let $\Phi := (f_1, \dots, f_m)$ be their coordinatewise action on \mathbb{A}^m , and let L be a line in \mathbb{A}^m defined over $\overline{\mathbb{Q}}$. If L is not Φ -preperiodic, then there exists an $\epsilon > 0$ such that*

$$S_{L, \Phi, \epsilon} := \{x \in L(\overline{\mathbb{Q}}) \mid \widehat{h}_\Phi(x) \leq \epsilon\}$$

is finite (see Section 2 for the definition of \widehat{h}_Φ).

Baker and Hsia [BH05, Theorem 8.10] previously proved Theorem 1.3 in the special case where $f_1 = f_2$ and $m = 2$.

In Section 2 we introduce our notation for canonical heights associated to polynomials, then in Section 3 we present some general results regarding polynomials which share the same Julia set. In Section 4 we prove Theorem 1.3, and then describe few examples of preperiodic lines under the coordinatewise action of polynomials of same degree.

Acknowledgements. We would like to thank Arman Mimar for helpful conversations.

2. PRELIMINARIES

Heights. Let $\mathbb{M}_{\mathbb{Q}}$ be the usual set of absolute values on \mathbb{Q} , normalized so that the archimedean absolute value is simply the absolute value $|\cdot|$ and $|p|_p = 1/p$ for each p -adic absolute value $|\cdot|_p$. For any finite extension K of \mathbb{Q} we define \mathbb{M}_K to be the set of absolute values on K that extend elements of $\mathbb{M}_{\mathbb{Q}}$. Then, for any $x \in \overline{\mathbb{Q}}$ we define the Weil height of x to be

$$h(x) = \frac{1}{[\mathbb{Q}(x) : \mathbb{Q}]} \cdot \sum_{v \in \mathbb{M}_{\mathbb{Q}(x)}} \sum_{\substack{w|v \\ w \in \mathbb{M}_{\mathbb{Q}(x)}}} \log \max\{|x|_w^{[\mathbb{Q}(x)_w : \mathbb{Q}_v]}, 1\}$$

where \mathbb{Q}_v and $\mathbb{Q}(x)_w$ are the completions of \mathbb{Q} and $\mathbb{Q}(x)$ at v and w respectively (see [BG06, Chapter 1] for details).

For a polynomial $f \in \overline{\mathbb{Q}}[X]$ of degree greater than 1, define the f -canonical height $\widehat{h}_f : \overline{\mathbb{Q}} \rightarrow \mathbb{R}_{\geq 0}$ by

$$(2.1) \quad \widehat{h}_f(x) = \lim_{n \rightarrow \infty} \frac{h(f^n(x))}{(\deg f)^n},$$

following Call-Silverman [CS93] (where f^n denotes the n -th iterate of f).

Let $f_1, \dots, f_m \in \overline{\mathbb{Q}}[X]$ be polynomials of degree $d > 1$, and let $\Phi := (f_1, \dots, f_m)$ be their coordinatewise action on \mathbb{A}^m ; that is,

$$\Phi(x_1, \dots, x_m) = (f_1(x_1), \dots, f_m(x_m)).$$

We define the Φ -canonical height $\widehat{h}_\Phi : \mathbb{A}^m(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}_{\geq 0}$ by

$$\widehat{h}_\Phi(x_1, \dots, x_m) = \sum_{i=1}^m \widehat{h}_{f_i}(x_i).$$

Note that while \mathbb{A}^m is not a projective variety, Φ extends to a map $\tilde{\Phi} : (\mathbb{P}^1)^m \rightarrow (\mathbb{P}^1)^m$. Furthermore, $\tilde{\Phi}$ is polarizable, since

$$\tilde{\Phi}^* \bigotimes_{i=1}^m \text{pr}_i^* \mathcal{O}_{\mathbb{P}^1}(1) \cong \bigotimes_{i=1}^m \text{pr}_i^* \mathcal{O}_{\mathbb{P}^1}(d),$$

where pr_i is the projection of $(\mathbb{P}^1)^m$ onto its i -th coordinate.

Remark. Theorem 1.3 is *not* true if one allows the polynomials f_i to have different degrees. This is easily seen, for example, in the case where $m = 2$, the line L is the diagonal, and $f_2 = f_1^2$. The map $\Phi = (f_1, \dots, f_m)$ is only polarizable when $\deg f_i = \deg f_j$, so this is not a counterexample to Conjecture 1.2.

3. SYMMETRIES OF THE JULIA SET

In this section we recall the main results regarding polynomials which share the same Julia set. For a polynomial $f \in \mathbb{C}[X]$, we let $J(f)$ denote the Julia set of f (see [Bea91, Chapter 3] or [Mil99] for the definition of a Julia set of a rational function over the complex numbers). As proved by Beardon [Bea90, Bea92], any family of polynomials which have the same Julia set \mathcal{J} is determined by the symmetries of \mathcal{J} .

Definition 3.1. *If $f \in \mathbb{C}[z]$ is a polynomial and $J(f)$ is its Julia set, then the symmetry group $\Sigma(f)$ of $J(f)$ is defined by*

$$\Sigma(f) = \{\sigma \in \mathfrak{C} : \sigma(J(f)) = J(f)\},$$

where \mathfrak{C} is the group of conformal Euclidean isometries.

Beardon [Bea90, Lemma 3] computed $\Sigma(f)$ for any $f \in \mathbb{C}[z]$.

Lemma 3.2. *The isometry group $\Sigma(f)$ is a group of rotations about some point $\zeta \in \mathbb{C}$, and it is either trivial, or finite cyclic group, or the group of all rotations about ζ .*

To prove the result above, Beardon uses the fact that each polynomial f of degree $d \geq 2$ is conjugate to a monic polynomial $\tilde{f} \in \mathbb{C}[z]$ which has no term in z^{d-1} . Clearly, if $\tilde{f} = \gamma \circ f \circ \gamma^{-1}$ where $\gamma \in \mathbb{C}[z]$ is a linear polynomial, then $J(\tilde{f}) = \gamma(J(f))$; thus it suffices to prove Lemma 3.2 for \tilde{f} instead of f . If $\tilde{f}(z) = z^d$, then $J(\tilde{f})$ is the unit circle, and $\Sigma(\tilde{f})$ is the group of all rotations about 0 (see also [Bea90, Lemma 4]). If \tilde{f} is not a monomial, then we choose $b \geq 1$ maximal such that we can find $a \geq 0$ and $\tilde{f}_1 \in \mathbb{C}[z]$ satisfying $\tilde{f}(z) = z^a \tilde{f}_1(z^b)$. In this case, $J(\tilde{f})$ is the finite cyclic group of rotations generated by the multiplication by $\exp(2\pi i/b)$ on \mathbb{C} (see [Bea90, Theorem 5]). Beardon also proves the following result (see [Bea90, Lemma 7]) which we will use later.

Lemma 3.3. *If $f \in \mathbb{C}[z]$ is a polynomial of degree $d \geq 2$ and $\sigma \in J(f)$, then $f \circ \sigma = \sigma^d \circ f$.*

The following classification of polynomials which have the same Julia set is proven by Beardon in [Bea92, Theorem 1].

Lemma 3.4. *If $f, g \in \mathbb{C}[z]$ have the same Julia set, then there exists $\sigma \in \Sigma(f)$ such that $g = \sigma \circ f$.*

4. PROOF OF OUR MAIN RESULT

Proof of Theorem 1.3. Suppose that for every $\epsilon > 0$, the set $S_{L, \Phi, \epsilon}$ is infinite. We will show that this implies that L must be Φ -preperiodic.

We first note that it suffices to prove the theorem for the line $L' = (\sigma_1, \dots, \sigma_m)(L)$ and the map

$$\Phi' = (\sigma_1 \circ f_1 \circ \sigma_1^{-1}, \dots, \sigma_m \circ f_m \circ \sigma_m^{-1})$$

for some linear automorphisms $\sigma_1, \dots, \sigma_m$ of \mathbb{A}^1 . This follows from the fact that L is preperiodic for Φ if and only if L' is preperiodic for Φ' along with the equality

$$(4.1) \quad \widehat{h}_{\Phi'}(\sigma_1 \alpha_1, \dots, \sigma_m \alpha_m) = \widehat{h}_{\Phi}(\alpha_1, \dots, \alpha_m).$$

Note that (4.1) is a simple consequence of Definition 2.1, since $|h(\sigma_i x) - h(x)|$ is bounded for all $x \in \overline{\mathbb{Q}}$.

We now proceed by induction on m ; the case $m = 1$ is obvious.

If the projection of L on any of the coordinates consists of only one point, we are done by the inductive hypothesis. Indeed, without loss of generality, assume the projection of L on the first coordinate equals $\{z_1\}$, then $L = \{z_1\} \times L_1$, where $L_1 \subset \mathbb{A}^{m-1}$ is a line, and $\widehat{h}_{f_1}(z_1) = 0$. Since only preperiodic points have canonical height equal to 0 (see [CS93, Cor. 1.1.1]), we conclude that z_1 is f_1 -preperiodic, and thus we are done by the induction hypothesis applied to L_1 .

Suppose now that L projects dominantly onto each coordinate of \mathbb{A}^m . For each $i = 2, \dots, m$, we let L_i be the projection of L on the first and the i -th coordinates of \mathbb{A}^m . Then L_i is a line given by an equation $X_1 = \sigma_i(X_i)$,

for some linear polynomial $\sigma_i \in \overline{\mathbb{Q}}[X]$. Clearly, it suffices to show that for each $i = 2, \dots, m$, the line L_i is preperiodic under the action of (f_1, f_i) on the corresponding two coordinates of \mathbb{A}^m . Indeed, if for each $i = 2, \dots, m$, we show that there exist $a_i, b_i \in \mathbb{N}$ (with $a_i < b_i$) such that $(f_1^{a_i}, f_i^{a_i})(L_i) = (f_1^{b_i}, f_i^{b_i})(L_i)$, then

$$(f_1^a, \dots, f_m^a)(L) = (f_1^{a+b}, \dots, f_m^{a+b})(L),$$

where $a := \max_{i=2}^m a_i$, and b is the least common multiple of all $(b_i - a_i)$ for $i = 2, \dots, m$.

Let $\tilde{f}_i := \sigma_i \circ f_i \circ \sigma_i^{-1}$ and let $\Delta = (x, x) \in \mathbb{A}^2$ be the diagonal on \mathbb{A}^2 . By our remarks at the beginning of the proof, it suffices to show that $(\text{id}, \sigma_i)(L_i) = \Delta$ is preperiodic under the action of (f_1, \tilde{f}_i) . Furthermore, the fact that we have an infinite sequence $(z_{n,1}, z_{n,i}) \in L_i(\overline{\mathbb{Q}})$ with

$$\lim_{n \rightarrow \infty} \widehat{h}_{f_1}(z_{n,1}) = \lim_{n \rightarrow \infty} \widehat{h}_{\tilde{f}_i}(z_{n,i}) = 0$$

implies that we have

$$\lim_{n \rightarrow \infty} \widehat{h}_{\tilde{f}_i}(z_{n,1}) = 0,$$

because of (4.1). Fix an embedding $\theta : \overline{\mathbb{Q}} \rightarrow \mathbb{C}$ and let f_1^θ and \tilde{f}_i^θ be the images of f_1 and \tilde{f}_i , respectively, in $\mathbb{C}[X]$ under this embedding. Then, by [BH05, Corollary 4.6], the Galois orbits of the points $\{z_{n,1}\}_{n \in \mathbb{N}}$ are equidistributed with respect to the equilibrium measures on the Julia sets of both f_1^θ and \tilde{f}_i^θ . Since the support of the equilibrium measure μ_g of a polynomial $g \in \mathbb{C}[X]$ is equal to the Julia set of g ([BH05, Section 4]), we must have $J(\tilde{f}_i^\theta) = J(f_1^\theta)$.

By Lemma 3.4 (see also [BE87, AH96]), there exists a conformal Euclidean symmetry $\mu_i : z \rightarrow a_i z + b_i$ such that $\mu_i(J(f_1^\theta)) = J(f_1^\theta)$ and $\tilde{f}_i^\theta = \mu_i \circ f_1^\theta$. Note that a_i and b_i must be in the image of $\overline{\mathbb{Q}}$ under θ since the coefficients of f_1^θ and \tilde{f}_i^θ are. Let τ_i be the map $\tau_i : z \rightarrow \theta^{-1}(a_i)z + \theta^{-1}(b_i)$. Then we have $\tilde{f}_i = \tau_i \circ f_1$.

If τ_i has infinite order, then it follows from [Bea90, Lemma 4] that there exist linear polynomials γ_1, γ_i such that $\gamma_1 \circ f_1 \circ \gamma_1^{-1} = \gamma_i \circ \tilde{f}_i \circ \gamma_i^{-1} = X^d$. In this case, we reduce our problem to the usual Bogomolov conjecture for \mathbb{G}_m^2 , proved by Zhang [Zha92]. Indeed, Zhang proves that if a curve C in \mathbb{G}_m^2 has an infinite family of algebraic points with height tending to zero, then it must be a torsion translate of an algebraic subgroup of \mathbb{G}_m^2 ; that is, $C = \xi A$ where ξ has finite order and A is an algebraic subgroup of \mathbb{G}_m^2 . Since $(\xi A)^n = \xi^n A$ and ξ has finite order, it is clear that such a curve is preperiodic under the map $(X, Y) \mapsto (X^d, Y^d)$.

We may suppose then that τ_i has finite order. By Lemma 3.3, we have $f_1 \circ \tau_i = \tau_i^d \circ f_1$. Thus, we have

$$\tilde{f}_i^k = \tau_i^{(d^k - 1)/(d - 1)} \circ f_1^k$$

for all $k \geq 1$. Since τ_i has finite order, we conclude that the set

$$\{\tau_i^{(d^k-1)/(d-1)}\}_{k \geq 0}$$

is finite. This implies that the set of curves of the form $(f_1^k, \tilde{f}_i^k)(\Delta)$ is finite, which means the diagonal subvariety Δ is preperiodic under the action of (f_1, \tilde{f}_i) , as desired. \square

4.1. Examples. Note that Δ is periodic under Φ only if there is some n such that $f^n = g^n$. Take, for example, $f(x) = x^3$, $g(x) = -x^3$; then we have $f^2 = g^2$, so Δ has period two under the action of Φ . More generally, there exists some n such that $f^n = g^n$ if and only if $f(x) = -\beta + \gamma h(x + \beta)$ and $g(x) = -\beta + h(x + \beta)$ for some $\gamma \in \mathbb{C}^*$, $\beta \in \mathbb{C}$ and $h \in x^r \mathbb{C}[x^s]$ (with $r, s \geq 0$) such that $\gamma^s = 1$ and $\gamma^{(d^n-1)/(d-1)} = 1$ (see [GTZ08, Prop. 6.3]). On the other hand, if $f(x) = h(x^s)$ for some positive integer $s \geq 2$ and $\zeta \in \mathbb{C}$ is an s -th root of unity, and $g(x) = \zeta \cdot f(x)$, then Δ is Φ -preperiodic, but it is not Φ -periodic.

We believe that it is possible to extend the methods of the proof of Theorem 1.3 to the case of arbitrary rational maps $\varphi_1, \dots, \varphi_m$ of the same degree, though the proof seems to be much more difficult, requiring in particular Mimar's [Mim97] results on arithmetic intersections of metrized line bundles and an analysis of Douady-Hubbard-Thurston's [DH93] classification of critically finite rational maps with parabolic orbifolds. We intend to treat this problem in a future paper.

REFERENCES

- [AH96] P. Atela and J. Hu, *Commuting polynomials and polynomials with same Julia set*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **6** (1996), no. 12A, 2427–2432.
- [BE87] I. N. Baker and A. Erëmenko, *A problem on Julia sets*, Ann. Acad. Sci. Fenn. Ser. A I Math. **12** (1987), no. 2, 229–236.
- [BH05] M. H. Baker and L.-C. Hsia, *Canonical heights, transfinite diameters, and polynomial dynamics*, J. reine angew. Math. **585** (2005), 61–92.
- [Bea90] A. F. Beardon, *Symmetries of Julia sets*, Bull. London Math. Soc. **22** (1990), no. 6, 576–582.
- [Bea91] A. F. Beardon, *Iteration of rational functions*, Springer-Verlag, New York, 1991.
- [Bea92] A. F. Beardon, *Polynomials with identical Julia sets*, Complex Variables Theory Appl. **17** (1992), no. 3–4, 195–200.
- [BG06] E. Bombieri and W. Gubler, *Heights in Diophantine geometry*, New Mathematical Monographs, vol. 4, Cambridge University Press, Cambridge, 2006.
- [Bog91] F. A. Bogomolov, *Abelian subgroups of Galois groups*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 1, 32–67.
- [CS93] G. S. Call and J. Silverman, *Canonical heights on varieties with morphism*, Compositio Math. **89** (1993), 163–205.
- [DH93] A. Douady and J. H. Hubbard, *A proof of Thurston's topological characterization of rational functions*, Acta Math. **171** (1993), no. 2, 263–297.
- [GTZ08] D. Ghioca, T. J. Tucker, and M. E. Zieve, *Intersections of polynomials orbits, and a dynamical Mordell-Lang conjecture*, Invent. Math. **171** (2008), no. 2, 463–483.
- [Mil99] J. Milnor, *Dynamics in one complex variable*, Vieweg, Braunschweig, 1999.

- [Mim97] A. Mimar, *On the preperiodic points of an endomorphism of $\mathbb{P}^1 \times \mathbb{P}^1$ which lie on a curve*, Ph.D. thesis, Columbia University, 1997.
- [Ull98] E. Ullmo, *Positivité et discrétion des points algébriques des courbes*, Ann. of Math. (2) **147** (1998), no. 1, 167–179.
- [Zha92] S. Zhang, *Positive line bundles on arithmetic surfaces*, Annals of Math **136** (1992), 569–587.
- [Zha95a] ———, *Positive line bundles on arithmetic varieties*, J. Amer. Math. Soc. **8** (1995), 187–221.
- [Zha95b] ———, *Small points and adelic metrics*, J. Algebraic Geometry **4** (1995), 281–300.
- [Zha98] S. Zhang, *Equidistribution of small points on abelian varieties*, Ann. of Math. (2) **147** (1998), no. 1, 159–165.
- [Zha06] S. Zhang, *Distributions in Algebraic Dynamics*, Survey in Differential Geometry, vol. 10, International Press, 2006, pp. 381–430.

DRAGOS GHIOCA, DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, UNIVERSITY OF LETHBRIDGE, LETHBRIDGE, AB T1K 3M4
E-mail address: `dragos.ghioca@uleth.ca`

THOMAS TUCKER, DEPARTMENT OF MATHEMATICS, HYLAN BUILDING, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627
E-mail address: `ttucker@math.rochester.edu`