

Computation of π

We use the formula

$$\frac{\pi}{4} = 4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{239}$$

which was discovered by the English mathematician John Machin. He used it to compute π to 100 decimal places in 1706.

Proof of Machin's Formula

Let $\beta = \tan^{-1} \frac{1}{5}$. Then $\tan \beta = \frac{1}{5}$. Using the double angle formula

$$\tan(2\theta) = \frac{\sin(2\theta)}{\cos(2\theta)} = \frac{2 \sin \theta \cos \theta}{\cos^2 \theta - \sin^2 \theta} = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

twice

$$\tan(2\beta) = \frac{2 \tan \beta}{1 - \tan^2 \beta} = \frac{2/5}{1 - 1/25} = \frac{10}{24} = \frac{5}{12}$$

$$\tan(4\beta) = \frac{2 \tan(2\beta)}{1 - \tan^2(2\beta)} = \frac{5/6}{1 - 25/144} = \frac{120}{119}$$

Then using the addition formula

$$\tan(x - y) = \frac{\sin(x-y)}{\cos(x-y)} = \frac{\sin x \cos y - \cos x \sin y}{\cos x \cos y + \sin x \sin y} = \frac{\tan x - \tan y}{1 + \tan x \tan y}$$

and $\tan \frac{\pi}{4} = 1$

$$\tan\left(4\beta - \frac{\pi}{4}\right) = \frac{\frac{120}{119} - 1}{1 + \frac{120}{119}} = \frac{120 - 119}{120 + 119} = \frac{1}{239}$$

Taking the arctan of both sides gives

$$4 \tan^{-1} \frac{1}{5} - \frac{\pi}{4} = \tan^{-1} \frac{1}{239}$$

which is what we want.

A Series Expansion for $\tan^{-1} x$

We already know that for $|s| < 1$

$$\frac{1}{1-s} = 1 + s + s^2 + s^3 + \dots$$

Subbing in $s = -t^2$ gives, for $|t| < 1$

$$\frac{1}{1+t^2} = 1 - t^2 + t^4 - t^6 + \dots$$

Integrating gives, for $|x| < 1$,

$$\tan^{-1} x = \int_0^x \frac{1}{1+t^2} dt = \int_0^x [1 - t^2 + t^4 - t^6 + \dots] dt = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

Furthermore, for $0 \leq x < 1$, the requirements of the alternating series test apply and truncating the series introduces an error between 0 and the first term omitted.

The Computation of π

$$\begin{aligned}\tan^{-1} \frac{1}{5} &= \frac{1}{5} - \frac{1}{3 \times 5^3} + \frac{1}{5 \times 5^5} \\ &\quad - \frac{1}{7 \times 5^7} + \frac{1}{9 \times 5^9} - \frac{1}{11 \times 5^{11}} \\ &= 0.2000000000 - 0.0026666667 + 0.0000640000 \\ &\quad - 0.0000018286 + 0.0000000569 - 0.0000000019 \\ &= 0.197395562 - e \text{ with } 0 \leq e \leq 1.9 \times 10^{-9} + 5 \times 0.5 \times 10^{-10} \\ \tan^{-1} \frac{1}{239} &= \frac{1}{239} - \frac{1}{3 \times 239^3} + \frac{1}{5 \times 239^5} \\ &= 0.0041841004 - 0.0000000244 + 2.6 \times 10^{-13} \\ &= 0.0041840760 + e' \text{ with } 0 \leq e' \leq 2.6 \times 10^{-13} + 3 \times 0.5 \times 10^{-10}\end{aligned}$$

The extra 10^{-10} terms take roundoff error into account. So

$$4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{239} = 0.785398172 - 4e - e'$$

and

$$\pi = 3.141592688 - 16e - 4e'$$

$$\pi = 3.14159269 - e'' \text{ with } e'' \text{ between } 0 \text{ and } 4 \times 10^{-8}$$