

1 **UNIFORM BOUND ON THE NUMBER OF PARTITIONS FOR**
2 **OPTIMAL CONFIGURATIONS OF THE OHTA-KAWASAKI**
3 **ENERGY IN 3D***

4 XIN YANG LU[†] AND JUN-CHENG WEI[‡]

5 **Abstract.** We study a 3D ternary system which combines an interface energy with a long range
6 interaction term. Several such systems were derived as a sharp-interface limit of the Nakazawa-
7 Ohta density functional theory of triblock copolymers. Both the binary case in 2D and 3D, and
8 the ternary case in 2D, are quite well understood, while very little is known about the ternary case
9 in 3D. In particular, it is even unclear whether minimizers are made of finitely many components.
10 In this paper we provide a positive answer to this, by proving that the number of components in
11 a minimizer is bounded from above by a computable quantity depending only on the total masses
12 and the interaction coefficients. There are two key difficulties, namely the impossibility to decouple
13 the long range interaction from the perimeter term, and the absence of a quantitative isoperimetric
14 inequality with two mass constraints in 3D. Therefore, the actual shape of minimizers is unknown,
15 even for small masses, making the construction of suitable competing configurations significantly
16 more delicate.

17 **Key words.** Pattern formation, small volume-fraction limit, triblock copolymers.

18 **AMS subject classifications.** 49S05, 35K30, 35K55

19 **1. Introduction.** Energy functionals entailing a direct competition between an
20 attractive short-range force and a repulsive Coulombic long-range force have been
21 studied intensively in recent years, to understand physical problems such as Gamow’s
22 liquid drop problem, and self-assembly of block copolymers. In Gamow’s liquid drop
23 model [10], the volume of the nucleus $\Omega \subset \mathbb{R}^3$ is fixed, i.e., $|\Omega| = m$ with the parameter
24 m being referred to as “mass”. The binding energy is given by

25
$$\mathcal{E}_{\text{liquid}}(\Omega) := \text{Per}(\Omega) + \frac{1}{8\pi} \int_{\Omega \times \Omega} \frac{dx dy}{|x - y|},$$

26 where the first term is the perimeter (or surface area) of Ω , which arises due to the
27 lower nucleon density near the nucleus boundary; the second term is a Coulomb-type
28 one, introduced to account for the presence of positively charged protons [3].

29 In Ohta and Kawasaki’s diblock copolymer model [21], the free energy is given by

30
$$\mathcal{E}_{\text{diblock}}(\Omega) := \text{Per}(\Omega) + \gamma \int_{\Omega \times \Omega} G(x, y) \, dx dy,$$

31 where the first term, i.e. the perimeter, favors a large ball; the second term prefers
32 splitting, and models long-range interactions between monomers due to the connec-
33 tivity of different subchains in copolymer molecules. Here

34
$$G(x, y) = \frac{1}{4\pi|x - y|} + R(x, y)$$

*SUBMITTED TO THE EDITORS 01/28/2022.

Funding: XYL ACKNOWLEDGES THE SUPPORT OF NSERC. THE RESEARCH OF J. WEI IS PARTIALLY SUPPORTED BY NSERC.

[†]Department of Mathematical Sciences, Lakehead University, Thunder Bay, ON, Canada (xlu8@lakeheadu.ca).

[‡]Department of Mathematics, University of British Columbia, Vancouver, BC, Canada (jcwei@math.ubc.ca).

35 is the zero-average Green's function of the Laplace operator in \mathbb{R}^3 , $R(x, y)$ is the
 36 regular part of $G(x, y)$, and γ is the long-range interaction coefficient, determined
 37 by the percentage of each type monomer, the total number of monomers in a chain
 38 molecule, the repulsion between different monomers, and the average distance between
 39 two adjacent monomers [6]. During each experiment, the total mass of each type
 40 monomer is fixed. So the energy is minimized under the mass constraint $|\Omega| = m$.

41 In this paper, we study a model in ternary systems, introduced by Nakazawa and
 42 Ohta to study triblock copolymers [20]. A triblock copolymer is a chain molecule
 43 consisting of three types of subchains: a subchain of type A monomers is connected
 44 to a subchain of type B monomers, and then connected to a subchain of type C
 45 monomers. Block copolymers can be used as a material in artificial organ technology
 46 and controlled drug delivery.

47 The free energy of triblock copolymers, in the sharp interface model, was derived
 48 by Ren and Wei in [24, 23] as the Γ -limit of Nakazawa and Ohta's diffuse interface
 49 model:

$$50 \quad \mathcal{E}_{\text{triblock}}(\Omega_1, \Omega_2) := \frac{1}{2} \sum_{i=0}^2 \text{Per}(\Omega_i) + \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_i \times \Omega_j} G(x, y) \, dx \, dy.$$

52 Here $\Omega_0 = (\Omega_1 \cup \Omega_2)^c$, the perimeter term is defined by

$$53 \quad \frac{1}{2} \sum_{i=0}^2 \text{Per}(\Omega_i) = \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j),$$

55 and the long-range interaction coefficients γ_{ij} form a 2×2 symmetric matrix. Using
 56 a ‘‘droplet’’ scaling argument, as done by Choksi and Peletier in [4, 5], and by Alama,
 57 Bronsard, the first author, and Wang in [1], it can be shown that the leading order of
 58 the free energy takes the form

$$59 \quad (1.1) \quad E_0(\Omega_1, \Omega_2) = \sum_k e_0(|\Omega_{1,k}|, |\Omega_{2,k}|), \quad \Omega_i = \bigcup_k \Omega_{i,k}, \quad i = 1, 2,$$

61 with

$$62 \quad e_0 : [0, +\infty) \times [0, +\infty) \longrightarrow \mathbb{R},$$

$$63 \quad e_0(m_1, m_2) := \inf \left\{ \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) \right.$$

$$64 \quad \left. + \sum_{i,j=1}^2 \frac{\Gamma_{ij}}{4\pi} \int_{\Omega_i \times \Omega_j} \frac{dx \, dy}{|x - y|} : |\Omega_i| = m_i, \, i = 1, 2 \right\},$$

66 where Γ_{ij} is a suitable scaling of γ_{ij} . That is, E_0 seeks the optimal partition $\Omega_i =$
 67 $\bigcup_k \Omega_{i,k}$, with each couple $(\Omega_{1,k}, \Omega_{2,k})$ minimizing e_0 .

68 Choksi and Peletier showed in [4, Theorem 4.2] that, when the domain is the unit
 69 torus \mathbb{T}^3 , in the small mass volume fraction regime, the first order Γ -limit of the free
 70 energies (see [4, Equation (1.8)])

$$71 \quad E_\eta^{3d}(v) := \begin{cases} \eta \int_{\mathbb{T}^3} |\nabla v| \, dx + \eta \left\| v - \frac{1}{|\mathbb{T}^3|} \int_{\mathbb{T}^3} v \, dx \right\|_{\mathcal{H}^{-1}(\mathbb{T}^3)}^2 & \text{if } v \in BV(\mathbb{T}^3; \{0, \eta^{-3}\}), \\ +\infty & \text{otherwise,} \end{cases}$$

72

73 is of the form

74 perimeter + long range interaction,

75 i.e. (see [4, Equation (4.1)], and more in general [4, Section 4]),

$$76 \quad E_0^{3d}(v) := \begin{cases} \sum_{k=0}^{\infty} e_0(m_k) & \text{if } v = \sum_{k=0}^{\infty} m_k \delta_{x_k}, \sum_{k=0}^{\infty} m_k = M = \text{total mass,} \\ +\infty & \text{otherwise,} \end{cases}$$

77 with

$$79 \quad e_0^{3d}(m) = \inf \left\{ \int_{\mathbb{R}^3} |\nabla z| dx + \|z\|_{H^{-1}(\mathbb{R}^3)}^2 : z \in BV(\mathbb{R}^3; \{0, 1\}), \|z\|_{L^1(\mathbb{R}^3)} = M \right\}.$$

80 The H^{-1} norm can be made explicit:

$$81 \quad \|z\|_{H^{-1}(\mathbb{R}^3)}^2 = \int_{\mathbb{R}^3 \times \mathbb{R}^3} G(|x-y|) z(x) z(y) dx dy,$$

82 where G denotes the Green's function of the Laplacian in \mathbb{R}^3 . That is, the minima
83 seeks the optimal partition, in which each component minimizes the energy e_0^{3d} . An
84 analogous result, but for ternary systems in the two dimensional torus, was obtained
85 in by Alama, Bronsard, the first author, and Wang, in [1, Theorem 3.2].

86 With the same arguments from [4, 1], it is possible to show that, again, with the
87 domain being the unit torus \mathbb{T}^3 , in the small mass volume fraction regime, the first
88 order Γ -limit of the free energies (which are the analogue of [1, Equation (1.8)] for
89 ternary systems in 3D)

$$90 \quad E_{\text{ternary}, \eta}^{3d}(v_{1,\eta}, v_{2,\eta}) := \begin{cases} f_{\eta}(v_{1,\eta}, v_{2,\eta}) & \text{if } v_{1,\eta}, v_{2,\eta} \in BV(\mathbb{T}^3; \{0, \frac{1}{\eta^3}\}), \\ +\infty & \text{otherwise,} \end{cases}$$

$$91 \quad f_{\eta}(v_{1,\eta}, v_{2,\eta}) := \frac{\eta}{2} \sum_{i=0}^2 \int_{\mathbb{T}^3} |\nabla v_{i,\eta}| dx$$

$$92 \quad + \eta^4 \sum_{i,j=1}^2 \gamma_{ij} \int_{\mathbb{T}^3 \times \mathbb{T}^3} G_{\mathbb{T}^3}(|x-y|) v_{i,\eta}(x) v_{j,\eta}(y) dx dy,$$

93 $G_{\mathbb{T}^3} :=$ Green's function of the Laplacian in \mathbb{T}^3 with zero average,

94 can be again written in the form

$$96 \quad (1.2) \quad E_{\text{ternary}, 0}^{3d}(v_1, v_2) := \begin{cases} \sum_{k=0}^{\infty} e_0(m_{1,k}, m_{2,k}) & \text{if } v_i = \sum_{k=0}^{\infty} m_{i,k} \delta_{x_{i,k}}, \sum_{k=0}^{\infty} m_{i,k} = M_i, \\ +\infty & \text{otherwise,} \end{cases}$$

97 $M_i =$ total mass of type i constituent, $i = 1, 2,$

98 where

$$100 \quad e_0(m_1, m_2) = \inf \left\{ \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) + \sum_{i,j=1}^2 \Gamma_{ij} \|z_i\|_{H^{-1}(\mathbb{R}^3)}^2 : \right.$$

$$101 \quad z_i \in BV(\mathbb{R}^3; \{0, 1\}), \|z_i\|_{L^1(\mathbb{R}^3)} = m_i,$$

$$102 \quad \Omega_i = \text{supp } z_i, i = 1, 2, |\Omega_1 \cap \Omega_2| = 0 \left. \right\}, \quad \Omega_0 = (\Omega_1 \cup \Omega_2)^c,$$

103

104 and $\Gamma_{ij}\eta^{-3} = \gamma_{ij} \geq 0$ are coefficients penalizing the Coulomb interaction. Observe
 105 that the problem of minimizing $E_{\text{ternary},0}^{3d}$ is again fully determined once we fix the
 106 total masses M_i and the interaction coefficients Γ_{ij} . Each couple of sets (Ω_1, Ω_2) ,
 107 with the appropriate masses, and minimizing e_0 , is referred to as a “cluster”.

108 Next, we introduce the main energy of this paper: given connected sets Ω_i , with
 109 $\mathbf{1}_{\Omega_i} \in BV(\mathbb{R}^3; \{0, 1\})$, $i = 1, 2$, and $|\Omega_1 \cap \Omega_2| = 0$, define the energy

$$110 \quad (1.3) \quad E(\Omega_1, \Omega_2) := \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) + \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_i \times \Omega_j} |x - y|^{-1} dx dy,$$

112 where $\Omega_0 = (\Omega_1 \cup \Omega_2)^c$. Here γ_{ij} denote the interaction strengths, and are positive,
 113 of order $O(1)$. Note that E is the analogue of e_0 from (1.1), (1.2), and [1], and of e_0^{3d}
 114 from [4], for ternary systems with domain \mathbb{R}^3 . Then, given disjoint unions

$$115 \quad \left(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k} \right),$$

116 with $\Omega_{i,k}$ being the connected components, the total energy of this configuration is
 117 defined by

$$118 \quad \mathcal{E} \left(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k} \right) := \sum_k E(\Omega_{1,k}, \Omega_{2,k}).$$

119 Observe that \mathcal{E} is the analogue of [1, Equation (3.5)] and [4, Equation (4.1)], for 3D
 120 ternary systems. It is also worthy noting that \mathcal{E} is similar to $\mathcal{E}_{\text{liquid}}$, $\mathcal{E}_{\text{diblock}}$, and
 121 $\mathcal{E}_{\text{triblock}}$, as they are all of the form

$$122 \quad \text{perimeter} + \text{long range interaction},$$

123 with the main difference being that \mathcal{E} suppresses the interaction between different
 124 connected components.

125 In the following, when we say “optimal configuration”, unless otherwise specified,
 126 we mean a configuration $(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k})$ minimizing \mathcal{E} .

127 In 2D, due to the fact that the Green’s function is a logarithmic term, the inter-
 128 action was simply the product of the masses, hence it was equivalent to minimize the
 129 perimeter, subject to two mass constraints. It is well known that the double bubble
 130 is the unique such minimizer (see e.g. [8, 18] for the 2D case, and [12] for the 3D case,
 131 and also [22, 7, 16, 17]). In the ternary 3D case, however, such simplification is not
 132 available, and the shape of the minimizers is unclear, even for small masses. This is
 133 a significant hurdle, and studying the shape of minimizers is hindered by the lack of
 134 a quantitative isoperimetric inequality with two mass constraints in 3D

135 Therefore, a priori, it is even unclear whether optimal configurations have finitely
 136 many clusters, as we cannot exclude the presence of infinitely many components with
 137 very small masses. Our main result is to show that this is not the case:

138 **THEOREM 1.** *There exists a computable constant $K = K(M_1, M_2, \gamma_{11}, \gamma_{22})$ such*
 139 *that any optimal configuration has at most K clusters.*

140 **Notation.** Since the position of the clusters is rarely relevant, in this paper we
 141 denote by B_m a ball of mass m .

2. Uniform upper bound on the number of clusters. The proof of Theorem 1 will be split over several lemmas. Throughout the entire section, M_i , $i = 1, 2$, will denote the total masses of type i constituent, and γ_{ij} , $i, j = 1, 2$ will denote the interaction coefficients. These parameters completely determine the minimization problem for \mathcal{E} in 3D. All the M_i and γ_{ij} will assumed to be given, and do not change throughout the section. Our proof will proceed as follows.

1. First, in Lemma 2, we bound from above the number of clusters made purely of one constituent type. Such upper bound will depend only on M_i , γ_{ii} , $i = 1, 2$.
2. Then, in Lemma 3, we show that the total mass of the largest cluster's cannot be too small. Such lower bound on the mass will depend only on M_i , γ_{ii} , $i = 1, 2$.
3. Finally, in Lemmas 4 and 5 we show that the total mass of each cluster is bounded from below by a constant depending only on M_i , γ_{ii} , $i = 1, 2$. Since there is only so much total mass (i.e., $M_1 + M_2$), this allows us to infer Theorem 1.

As we have no information on the shape of optimal configurations, we will often compare their energy against that of a suitable standard double bubble. Further information about the geometry of standard double bubbles are available in the Appendix.

LEMMA 2. Consider an optimal configuration, made of clusters $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq 1$. Then

$$\#\{k : |\Omega_{1,k}| |\Omega_{2,k}| = 0\}$$

is bounded from above by a constant depending only on M_i , γ_{ii} , $i = 1, 2$.

Proof. It is well known (see e.g. [13, 2, 19, 9, 14, 15], and references therein) that there exist $m_{i,B} = m_{i,B}(\gamma_{ii}) > 0$, $i = 1, 2$, such that, for all $m \leq m_{i,B}(\gamma_{ii})$, the minimizer of

$$\inf_{|X|=m} \left\{ \mathcal{H}^2(\partial X) + \gamma_{ii} \int_{X \times X} |x - y|^{-1} dx dy \right\}$$

is given by B_m . Since $\mathcal{H}^2(\partial B_m)$ (resp. $\int_{X \times X} |x - y|^{-1} dx dy$) scales like $m^{2/3}$ (resp. $m^{5/3}$), the perimeter term is dominating for all sufficiently small masses. Thus there exist geometric constants $m_{i,S} = m_{i,S}(\gamma_{ii}) \leq m_{i,B}(\gamma_{ii})$ such that

$$\begin{aligned} & \mathcal{H}^2(\partial B_{m_1}) + \gamma_{ii} \int_{B_{m_1} \times B_{m_1}} |x - y|^{-1} dx dy \\ & + \mathcal{H}^2(\partial B_{m_2}) + \gamma_{ii} \int_{B_{m_2} \times B_{m_2}} |x - y|^{-1} dx dy \\ & > \mathcal{H}^2(\partial B_{m_1+m_2}) + \gamma_{ii} \int_{B_{m_1+m_2} \times B_{m_1+m_2}} |x - y|^{-1} dx dy, \end{aligned}$$

for all $m_1, m_2 \leq m_{i,S}(\gamma_{ii})$, i.e. combining the two balls is energetically favorable whenever $m_1, m_2 \leq m_{i,S}(\gamma_{ii})$. Thus we cannot have two balls of the type i constituent, both with masses less than $m_{i,S}(\gamma_{ii})$. Since the total mass is $M_1 + M_2 < +\infty$, the proof is complete. \square

LEMMA 3. Consider an optimal configuration, made of clusters $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq 1$. Then

$$m_i^+ := \sup_k m_{i,k}, \quad m_{i,k} := |\Omega_{i,k}|, \quad i = 1, 2,$$

183 is bounded from below by

$$184 \quad \min \left\{ \frac{M_i}{2}, \left(\frac{\sqrt[3]{36\pi} M_i}{4 \sum_{i=1}^2 [\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy]} \right)^3 \right\}, \quad i = 1, 2.$$

185 Note that, curiously, this lower bound is independent of γ_{12} . As it will be clear
 186 from the proof, this is due to the fact that an upper bound for the energy of an optimal
 187 configuration is given by the energy of two balls of masses M_1 and M_2 respectively.
 188 Such bound is clearly independent of γ_{12} .

189 *Proof.* The idea is that, for very small masses, the perimeter term is sub-additive
 190 and dominating. Assume $m_i^+ \leq M_i/2$, as otherwise $M_i/2$ is already a lower bound.
 191 Note that

$$192 \quad E(\Omega_{1,k}, \Omega_{2,k}) \geq \mathcal{S}(m_{1,k}, m_{2,k}) \quad \forall k \geq 1,$$

193 where

$$194 \quad (2.1) \quad \mathcal{S}(m_1, m_2) = \text{perimeter of the standard double bubble with masses } m_1 \text{ and } m_2,$$

196 and, by [11, Theorem 4.2] (applied with $v_1 = m_1$, $x = v_2 = m_2$, $n = 3$)

$$197 \quad \mathcal{S}(m_1, m_2) \geq \sum_{i=1}^2 c_i m_i^{2/3}, \quad c_1 = c_2 = \frac{\sqrt[3]{36\pi}}{2}.$$

198 Thus the total energy of our optimal configuration satisfies

$$199 \quad \sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) \geq \sum_{i=1}^2 c_i \sum_{k \geq 1} m_{i,k}^{2/3}.$$

201 By the concavity of the function $t \mapsto t^{2/3}$, the sum $\sum_{k \geq 1} m_{i,k}^{2/3}$ is minimum when
 202 $m_{i,k} \in \{0, m_i^+\}$ for all k . Since $\sum_{k \geq 1} m_{i,k} = M_i$, there are at least $\lfloor \frac{M_i}{m_i^+} \rfloor$ many
 203 clusters containing type i constituents, thus

$$204 \quad \begin{aligned} \sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) &\geq \sum_{i=1}^2 c_i \sum_{k \geq 1} m_{i,k}^{2/3} \geq \sum_{i=1}^2 c_i \left\lfloor \frac{M_i}{m_i^+} \right\rfloor (m_i^+)^{2/3} \\ &\geq \sum_{i=1}^2 c_i \frac{M_i - m_i^+}{(m_i^+)^{1/3}} \geq \sum_{i=1}^2 \frac{c_i}{2} \frac{M_i}{(m_i^+)^{1/3}}. \end{aligned}$$

207 Since our configuration was an optimal one, its energy does not exceed that of two
 208 balls, which we denote by B_{M_1} and B_{M_2} , of masses M_1 and M_2 , respectively. Thus
 209 the above line continues as

$$210 \quad \begin{aligned} \sum_{i=1}^2 \frac{c_i}{2} \frac{M_i}{(m_i^+)^{1/3}} &\leq \sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) \\ &\leq \sum_{i=1}^2 \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy \right], \end{aligned}$$

212

213 hence

$$214 \quad (m_i^+)^{1/3} \geq \frac{c_i M_i}{2 \sum_{i=1}^2 [\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy]},$$

215 and the proof is complete. \square

216 **LEMMA 4.** *Consider an optimal configuration, made of clusters $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq$*
 217 *1. Assume $\sup_k |\Omega_{1,k}|$ and $\sup_k |\Omega_{2,k}|$ are achieved on different clusters, i.e., without*
 218 *loss of generality,*

$$219 \quad |\Omega_{1,1}| = m_1^+ = \sup_k |\Omega_{1,k}|, \quad |\Omega_{2,2}| = n_2^+ = \sup_k |\Omega_{2,k}|.$$

220 Then

$$221 \quad \inf_k \sum_{i=1}^2 |\Omega_{i,k}|$$

222 is bounded from below by a constant depending only on M_i , γ_{ii} , $i = 1, 2$.

Proof. Consider a cluster $(\Omega_{1,k}, \Omega_{2,k})$, with $k \geq 3$, and let

$$m_2 := |\Omega_{2,1}|, \quad n_1 := |\Omega_{1,2}|, \quad \varepsilon_i := |\Omega_{i,k}| > 0, \quad i = 1, 2.$$

223 Note that $m_1^+ \geq n_1$, $n_2^+ \geq m_2$. The construction will be slightly different depending
 224 on the values of $\frac{m_1^+}{m_2}$, $\frac{m_2}{n_2^+}$, and $\frac{\varepsilon_1}{\varepsilon_2}$.

225 *Case 1:* $\frac{m_1^+}{m_2} \geq \frac{\varepsilon_1}{\varepsilon_2}$. Consider the competitor constructed in the following way (see
 226 Figure 1).

- 227 • Move mass ε_1 (resp. rm_2 , with $r := \frac{\varepsilon_1}{m_1^+} \leq 1$) of type I (resp. type II)
 228 constituent from the cluster $(\Omega_{1,k}, \Omega_{2,k})$ to $(\Omega_{1,1}, \Omega_{2,1})$. This is possible since
 229 we are discussing the case $\frac{m_1^+}{m_2} \geq \frac{\varepsilon_1}{\varepsilon_2}$, i.e. $rm_2 = \varepsilon_1 \frac{m_2}{m_1^+} \leq \varepsilon_2$.
- 230 • Replace $(\Omega_{1,k}, \Omega_{2,k})$ and $(\Omega_{1,1}, \Omega_{2,1})$ with $B_{\varepsilon_2 - rm_2}$ (of type II constituent)
 231 and $(\tilde{\Omega}_{1,1}, \tilde{\Omega}_{2,1}) := (1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})$, while every other cluster remains
 232 unaltered.

233 Now we estimate the change in energy. Since our initial configuration was optimal,

$$234 \quad 0 \leq E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) + E(\emptyset, B_{\varepsilon_2 - rm_2})$$

$$235 \quad (2.2) \quad - E(\Omega_{1,1}, \Omega_{2,1}) - E(\Omega_{1,k}, \Omega_{2,k}).$$

237 By a straightforward scaling argument,

$$238 \quad E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1}))$$

$$239 \quad = (1+r)^{2/3} \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}), \quad \Omega_{0,1} := (\Omega_{1,1} \cup \Omega_{2,1})^c,$$

$$240 \quad + (1+r)^{5/3} \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy$$

$$241 \quad \leq (1+r) \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}) + (1+3r) \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy$$

$$242 \quad \leq (1+3r) \left[\underbrace{\sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}) + \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy}_{=E(\Omega_{1,1}, \Omega_{2,1})} \right],$$

$$243$$




FIG. 1. Schematic representation of the construction of the competitor: original clusters (top), and modified clusters (bottom). Though the objects in question are three dimensional, for better clarity, we represented the construction in two dimensions. Only the affected clusters are represented here. The clusters are drawn deliberately deformed, to emphasize the fact that we do not know the clusters' precise shapes.

244 where we used the estimates

$$245 \quad (1+r)^{2/3} \leq 1+r \leq 1+3r, \quad (1+r)^{5/3} \leq (1+r)^2 \stackrel{(r \leq 1)}{\leq} 1+3r.$$

247 Thus, in view of Lemma 3,

$$248 \quad E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) - E(\Omega_{1,1}, \Omega_{2,1}) \\ 249 \quad (2.3) \quad \leq 3rE(\Omega_{1,1}, \Omega_{2,1}) \leq \varepsilon_1 H_1(M_1, M_2, \gamma_{11}, \gamma_{22}), \\ 250 \quad H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) := \sum_{i=1}^2 \frac{3}{m_1^+} \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy \right]. \\ 251$$

252 Now we estimate $E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,k}, \Omega_{2,k})$:

$$253 \quad E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,k}, \Omega_{2,k}) \leq \mathcal{S}(0, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \\ 254 \quad = \mathcal{S}(0, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) + \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \\ 255 \quad \leq -c_1 \varepsilon_1^{2/3}, \quad c_1 := \frac{\sqrt[3]{36\pi}}{2}, \\ 256$$

257 where the last line is due to [11, Theorem 3.2], which gives

$$258 \quad \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \leq 0,$$

259 and [11, Theorem 4.2] (applied with $v_1 = \varepsilon_1$, $x = v_2 = \varepsilon_2 - rm_2$, $n = 3$), which gives

$$\begin{aligned}
260 \quad \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) &\geq \frac{\sqrt[3]{36\pi}}{2} [\varepsilon_1^{2/3} + (\varepsilon_2 - rm_2)^{2/3} + (\varepsilon_1 + \varepsilon_2 - rm_2)^{2/3}] \\
261 &\geq \frac{\sqrt[3]{36\pi}}{2} [\varepsilon_1^{2/3} + 2(\varepsilon_2 - rm_2)^{2/3}] = \frac{\sqrt[3]{36\pi}}{2} \varepsilon_1^{2/3} + \underbrace{\frac{\sqrt[3]{36\pi}}{2} (\varepsilon_2 - rm_2)^{2/3}}_{=S(0, \varepsilon_2 - rm_2)}.
\end{aligned}$$

263 Combining with (2.2) and (2.3) gives the necessary condition

$$\begin{aligned}
264 \quad 0 &\leq E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) + E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,1}, \Omega_{2,1}) - E(\Omega_{1,k}, \Omega_{2,k}) \\
265 \quad (2.4) \quad &\leq \varepsilon_1 H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) - c_1 \varepsilon_1^{2/3},
\end{aligned}$$

267 hence

$$268 \quad \varepsilon_1^{1/3} \geq H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) c_1^{-1},$$

269 thus completing the proof for this case.

270 *Case 2:* $\frac{n_2^+}{n_1} \geq \frac{\varepsilon_2}{\varepsilon_1}$. The competitor constructed in a way similar to the previous
271 case.

- 272 • Move mass ε_2 (resp. rn_1 , with $r := \frac{\varepsilon_2}{n_2^+} \leq 1$) of type II (resp. type I)
273 constituent from the cluster $(\Omega_{1,k}, \Omega_{2,k})$ to $(\Omega_{1,2}, \Omega_{2,2})$. This is possible since
274 we are discussing the case $\frac{n_2^+}{n_1} \geq \frac{\varepsilon_2}{\varepsilon_1}$, i.e. $rn_1 = \varepsilon_2 \frac{n_1}{n_2^+} \leq \varepsilon_1$.
- 275 • Replace $(\Omega_{1,k}, \Omega_{2,k})$ and $(\Omega_{1,2}, \Omega_{2,2})$ with $B_{\varepsilon_1 - rn_1}$ (of type I constituent) and
276 $(1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})$, while every other cluster remains unaltered.

277 Then the proof proceeds like in the previous case. With the same arguments from
278 Case 1, we obtain

$$\begin{aligned}
279 \quad E((1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})) - E(\Omega_{1,2}, \Omega_{2,2}) &\leq 3rE(\Omega_{1,2}, \Omega_{2,2}) \leq \varepsilon_2 H_2(M_1, M_2, \gamma_{11}, \gamma_{22}), \\
280 \quad H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) &:= \sum_{i=1}^2 \frac{3}{n_2^+} \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy \right], \\
281
\end{aligned}$$

282 which is the analogue of (2.3), and

$$\begin{aligned}
283 \quad 0 &\leq E((1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})) + E(\emptyset, B_{\varepsilon_1 - rn_1}) - E(\Omega_{1,2}, \Omega_{2,2}) - E(\Omega_{1,k}, \Omega_{2,k}) \\
284 \quad &\leq \varepsilon_2 H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) - c_2 \varepsilon_2^{2/3}, \\
285
\end{aligned}$$

286 for some computable, purely geometric constant $c_2 > 0$, which is the analogue of
287 (2.4). Thus

$$288 \quad \varepsilon_2^{1/3} \geq H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) c_2^{-1},$$

289 concluding the proof for this case.

290 Finally, note that the above two cases are exhaustive: if Case 1 does not hold,
291 i.e. $\frac{\varepsilon_2}{\varepsilon_1} < \frac{m_2}{m_1^+}$, using $m_1^+ \geq n_1$, $n_2^+ \geq m_2$, we get

$$292 \quad \frac{\varepsilon_2}{\varepsilon_1} < \frac{m_2}{m_1^+} \leq \frac{n_2^+}{n_1},$$

293 i.e. Case 2 holds. The proof is thus complete. \square

294 LEMMA 5. Consider an optimal configuration, made of clusters $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq$
 295 1. Assume $\sup_k |\Omega_{1,k}|$ and $\sup_k |\Omega_{2,k}|$ are achieved on the same clusters, i.e., without
 296 loss of generality,

$$297 \quad |\Omega_{i,1}| = m_i^+ = \sup_k |\Omega_{i,k}|, \quad i = 1, 2.$$

298 Then

$$299 \quad \inf_k \sum_{i=1}^2 |\Omega_{i,k}|$$

300 is again bounded from below by a constant depending only on M_i, γ_{ii} , $i = 1, 2$.

301 *Proof.* We rely on Lemma 4: Consider another cluster $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq 2$. Let
 302 $|\Omega_{1,k}| = \varepsilon_1 > 0$, $|\Omega_{2,k}| = \varepsilon_2 > 0$, and note that one of the following cases must hold.

- 303 1. If $\frac{m_1^+}{m_2^+} \geq \frac{\varepsilon_1}{\varepsilon_2}$, then we can use the construction from Case 1 of Lemma 4.
 304 2. If $\frac{m_1^+}{m_2^+} \leq \frac{\varepsilon_1}{\varepsilon_2}$, i.e. $\frac{m_2^+}{m_1^+} \geq \frac{\varepsilon_2}{\varepsilon_1}$, then we can use the construction from Case 2 of
 305 Lemma 4.

306 The proof is thus complete. \square

307 **3. Appendix: geometry of the standard double bubble.** In [12], it was
 308 shown that the three dimensional standard double bubbles has the least surface area
 309 among all sets enclosing two regions of given volumes.




FIG. 2. The standard double bubble in \mathbb{R}^3 : if the two bubbles that meet have equal volumes, the shared surface between them is a flat disc. But in the case of unequal volumes, the smaller bubble, given its larger internal pressure, will bow slightly into the larger bubble. In either scenario, the two bubbles always meet at angles of 120 degrees. Credit: John M. Sullivan, Technical University of Berlin and University of Illinois at Urbana-Champaign.

310 Geometrically, the standard double bubble is a surface of revolution, with all the
 311 three surfaces being part of spheres, meeting at 120 degrees (see Figures 2 and 3).

312 Below we collect several results, used in the proof of Theorem 1, on the function
 313 \mathcal{S} introduced in (2.1).

314 LEMMA 6. [11, Theorem 3.2] The function \mathcal{S} is strictly concave: given $m_i, n_i \geq 0$,
 315 $i = 1, 2$, it holds

$$316 \quad \mathcal{S}((1-t)m_1 + tn_1, (1-t)m_2 + tn_2) > (1-t)\mathcal{S}(m_1, m_2) + t\mathcal{S}(n_1, n_2)$$




FIG. 3. Cross section of a standard double bubble.

318 for all $t > 0$.

319 COROLLARY 7. [11, Corollary 3.3] The function $\mathcal{S}(m_1, m_2)$ is increasing in both
 320 variables.

321 LEMMA 8. [11, Theorem 4.2] Suppose that in a minimal enclosure of volumes m_1
 322 and m_2 in \mathbb{R}^3 , with the latter having a connected component with volume $x > 0$. Then

$$323 \frac{2\mathcal{S}(m_1, m_2)}{c_1} \geq m_2 x^{-1/3} + m_1^{2/3} + (m_1 + m_2)^{2/3},$$

$$324 c_1 := \sqrt[3]{36\pi} = \text{surface area of the unit ball in } \mathbb{R}^3.$$

326 **Acknowledgments.** XYL acknowledges the support of NSERC. The research
 327 of J. Wei is partially supported by NSERC. We are grateful to Chong Wang for useful
 328 discussions and suggestions.

329

REFERENCES

- 330 [1] S. ALAMA, L. BRONSARD, X. LU, AND C. WANG, *Periodic minimizers of a ternary non-local*
 331 *isoperimetric problem*, Indiana Univ. Math. J., (2021).
- 332 [2] M. BONACINI AND R. CRISTOFERI, *Local and global minimality results for a nonlocal isoperimet-*
 333 *ric problem on R^N* , SIAM Journal on Mathematical Analysis, 46 (2014), pp. 2310–2349.
- 334 [3] R. CHOKSI, C. B. MURATOV, AND I. TOPALOGLU, *An old problem resurfaces nonlocally:*
 335 *Gamow's liquid drops inspire today's research and applications*, Notices of the AMS, 64
 336 (2017), pp. 1275–1283.
- 337 [4] R. CHOKSI AND M. A. PELETIER, *Small volume fraction limit of the diblock copolymer prob-*
 338 *lem: I. sharp-interface functional*, SIAM Journal on Mathematical Analysis, 42 (2010),
 339 pp. 1334–1370.
- 340 [5] ———, *Small volume-fraction limit of the diblock copolymer problem: II. Diffuse-interface*
 341 *functional*, SIAM Journal on Mathematical Analysis, 43 (2011), pp. 739–763.
- 342 [6] R. CHOKSI AND X. REN, *On the derivation of a density functional theory for microphase*
 343 *separation of diblock copolymers*, Journal of Statistical Physics, 113 (2003), pp. 151–176.
- 344 [7] R. DORFF, G. LAWLOR, D. SAMPSON, AND B. WILSON, *Proof of the planar double bubble*
 345 *conjecture using metacalibration methods*, Involve, a Journal of Mathematics, 2 (2010),
 346 pp. 611–628.
- 347 [8] J. FOISY, M. ALFARO GARCIA, J. BROCK, N. HODGES, AND J. ZIMBA, *The standard double soap*
 348 *bubble in R^2 uniquely minimizes perimeter*, Pacific journal of mathematics, 159 (1993),
 349 pp. 47–59.
- 350 [9] R. L. FRANK AND E. H. LIEB, *A compactness lemma and its application to the existence of*
 351 *minimizers for the liquid drop model*, SIAM Journal on Mathematical Analysis, 47 (2015),
 352 pp. 4436–4450.
- 353 [10] G. GAMOW, *Mass defect curve and nuclear constitution*, Proceedings of the Royal Society

- 354 of London. Series A, Containing Papers of a Mathematical and Physical Character, 126
 355 (1930), pp. 632–644.
- 356 [11] M. HUTCHINGS, *The structure of area-minimizing double bubbles*, The Journal of Geometric
 357 Analysis, 7 (1997), pp. 285–304.
- 358 [12] M. HUTCHINGS, F. MORGAN, M. RITORÉ, AND A. ROS, *Proof of the double bubble conjecture*,
 359 Annals of Mathematics, (2002), pp. 459–489.
- 360 [13] H. KNÜPFER AND C. B. MURATOV, *On an isoperimetric problem with a competing nonlocal*
 361 *term I: The planar case*, Communications on Pure and Applied Mathematics, 66 (2013),
 362 pp. 1129–1162.
- 363 [14] H. KNÜPFER, C. B. MURATOV, AND M. NOVAGA, *Low density phases in a uniformly charged*
 364 *liquid*, Communications in Mathematical Physics, 345 (2016), pp. 141–183.
- 365 [15] ———, *Emergence of nontrivial minimizers for the three-dimensional Ohta–Kawasaki energy*,
 366 Pure and Applied Analysis, 2 (2019), pp. 1–21.
- 367 [16] E. MILMAN AND J. NEEMAN, *The Gaussian double-bubble conjecture*, arXiv preprint
 368 arXiv:1801.09296, (2018).
- 369 [17] F. MORGAN AND M. RITORÉ, *Geometric measure theory and the proof of the double bubble*
 370 *conjecture*, in Proc. Clay Research Institution Summer School, MSRI, 2001.
- 371 [18] F. MORGAN AND W. WICHIRAMALA, *The standard double bubble is the unique stable double*
 372 *bubble in R^2* , Proceedings of the American Mathematical Society, 130 (2002), pp. 2745–
 373 2751.
- 374 [19] C. MURATOV AND H. KNÜPFER, *On an isoperimetric problem with a competing nonlocal term II:*
 375 *The general case*, Communications on Pure and Applied Mathematics, 67 (2014), pp. 1974–
 376 1994.
- 377 [20] H. NAKAZAWA AND T. OHTA, *Microphase separation of ABC-type triblock copolymers*, Macro-
 378 molecules, 26 (1993), pp. 5503–5511.
- 379 [21] T. OHTA AND K. KAWASAKI, *Equilibrium morphology of block copolymer melts*, Macromolecules,
 380 19 (1986), pp. 2621–2632.
- 381 [22] B. W. REICHARDT, C. HEILMANN, Y. Y. LAI, AND A. SPIELMAN, *Proof of the double bubble*
 382 *conjecture in R^4 and certain higher dimensional cases*, Pacific journal of mathematics, 208
 383 (2003), pp. 347–366.
- 384 [23] X. REN AND J. WEI, *Triblock copolymer theory: Free energy, disordered phase and weak seg-*
 385 *regation*, Physica D: Nonlinear Phenomena, 178 (2003), pp. 103–117.
- 386 [24] X. REN AND J. WEI, *Triblock copolymer theory: Ordered ABC lamellar phase*, Journal of
 387 Nonlinear Science, 13 (2003), pp. 175–208.