## NOTE

## Polygon Dissections and Standard Young Tableaux

Richard P. Stanley\*

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Communicated by the Managing Editors

Received April 4, 1995

A simple bijection is given between dissections of a convex (n+2)-gon with d diagonals not intersecting in their interiors and standard Young tableaux of shape  $(d+1, d+1, 1^{n-1-d})$ . © 1996 Academic Press, Inc.

For  $0 \le d \le n-1$ , let f(n, d) be the number of ways to draw d diagonals in a convex (n+2)-gon, such that no two diagonals intersect in their interior. For instance, f(n, n-1) is just the Catalan number  $C_n = \binom{2n}{n}/(n+1)$ . A result going back to Kirkman [3], Prouhet [4], and Cayley [1] (with Cayley giving the first complete proof) asserts that

$$f(n, d) = \frac{1}{n+d+2} \binom{n+d+2}{d+1} \binom{n-1}{d}.$$
 (1)

K. O'Hara and A. Zelevinsky observed (unpublished) that the right-hand side of (1) is just the number of standard Young tableaux (as defined, e.g., in [5, p. 66]) of shape  $(d+1, d+1, 1^{n-1-d})$ , where  $1^{n-1-d}$  denotes a sequence of n-1-d 1's. It is natural to ask for a bijection between the polygon dissections and the standard Young tableaux. If one is willing to accept the formula for the number of standard Young tableaux of a fixed shape (either in the original form due to MacMahon or the hook-length formula of Frame-Robinson-Thrall), then one obtains a simple proof

<sup>\*</sup> Partially supported by National Science Foundation Grant DMS-9206374. E-mail address: rstan@math.mit.edu.

of equation (1). In this note we give a simple bijection of the desired type.

First we recall that there is a well-known bijection [2] between dissections D of an (n+2)-gon with d diagonals and integer sequences  $\psi(D) = (a_1, a_2, ..., a_{n+d+1})$  such that (a) either  $a_i = -1$  or  $a_i \ge 1$ , (b) exactly n terms are equal to -1, (c)  $a_1 + a_2 + \cdots + a_i \ge 0$  for all i, and (d)  $a_1 + a_2 + \cdots + a_{n+d+1} = 0$ . This bijection may be defined recursively as follows. Fix an edge e of the dissected polygon D. When we remove e from D, we obtain a sequence of dissected polygons  $D_1, D_2, ..., D_k$  (where k+1 is the number of sides of the region of D to which e belongs), arranged in clockwise order, with  $D_i$  and  $D_{i+1}$  intersecting at a single vertex. If  $D_i$  consists of a single edge, then define  $\psi(D_i) = -1$ , and set recursively  $\psi(D) = (k-1, \psi(D_1)^*, \psi(D_2)^*, ..., \psi(D_{k-1})^*, \psi(D_k))$ , where  $\psi(D_k)^*$  denotes  $\psi(D_k)$  with a -1 appended at the end.

Given a sequence  $(a_1, a_2, ..., a_{n+d+1})$  as above, define a standard Young tableau T of shape  $(d+1, d+1, 1^{n-1-d})$  as follows. We insert the elements 1, 2, ..., n+d+1 successively into T. Once an element is inserted, it remains in place. (There is no "bumping" as in the Robinson-Schensted correspondence.) Suppose that the positive  $a_i$ 's are given by  $b_1, b_2, ..., b_{d+1}$ , in that order. The insertion is then defined by the following three rules:

• If  $a_i > 0$ , then insert *i* at the end of the first row. (We write our tableaux in "English" style, so the longest row is at the top.)

• If  $a_i = -1$  and the number of -1's preceding  $a_i$  is given by  $b_1 + b_2 + \cdots + b_j$  for some  $j \ge 0$ , then insert *i* at the end of the second row.

• If  $a_i = -1$  and the number of -1's preceding  $a_i$  in not of the form  $b_1 + b_2 + \cdots + b_j$ , then insert *i* at the bottom of the first column.

It is an easy exercise to check that the above procedure yields the desired bijection.

EXAMPLE. Let the sequence corresponding to a dissection D (with n = 14, d = 6) be given by

We have  $(b_1, ..., b_7) = (4, 2, 1, 3, 1, 1, 2)$ . We have printed in boldface those -1's that are preceded by  $b_1 + \cdots + b_j - 1$ 's for some *j*. The corresponding standard tableau  $\psi(D)$  is given by

NOTE

1	2	4	7	10	11	18
3	9	13	14	17	19	20
5			-	-	-	-
6						
8						
12						
15						
16						
21						

## REFERENCES

- 1. A. Cayley, On the partitions of a polygon, *Proc. London Math. Soc. (1)* **22** (1890–1891), 237–262.
- I. M. H. Etherington, Some problems of non-associative combinations (1), *Edingburgh Math.* 32 (1940), 1–6.
- 3. T. P. Kirkman, On the *k*-partitions of the *r*-gon and *r*-ace, *Philos. Trans. Roy. Soc. London* 147 (1857), 217–272.
- 4. E. Prouhet, Question 774, Nouvelles Ann. Math. 5 (1866), 384.
- 5. B. E. Sagan, "The Symmetric Group," Wadsworth & Brooks/Cole, Pacific Grove, CA, 1991.