



ELSEVIER

Discrete Mathematics 246 (2002) 57–66

DISCRETE
MATHEMATICS

www.elsevier.com/locate/disc

Shifted quasi-symmetric functions and the Hopf algebra of peak functions

Nantel Bergeron^{a,*}, Stefan Mykytiuk^a, Frank Sottile^{b,2},
Stephanie van Willigenburg^{a,3}

^aDepartment of Mathematics and Statistics, York University, Toronto, Ont. Canada, M3J 1P3

^bDepartment of Mathematics, University of Wisconsin, Van Vleck Hall, 480 Lincoln Drive,
Madison, WI. 53706-1388, USA

Abstract

In his work on P -partitions, Stembridge defined the algebra of peak functions Π , which is both a subalgebra and a retraction of the algebra of quasi-symmetric functions. We show that Π is closed under coproduct, and therefore a Hopf algebra, and describe the kernel of the retraction. Billey and Haiman, in their work on Schubert polynomials, also defined a new class of quasi-symmetric functions—shifted quasi-symmetric functions—and we show that Π is strictly contained in the linear span Ξ of shifted quasi-symmetric functions. We show that Ξ is a coalgebra, and compute the rank of the n th graded component.

Résumé

Dans ses travaux sur les P -partitions, Stembridge définit l'algèbre Π des fonctions de pics. Cette algèbre peut être vue comme une sous-algèbre ou un quotient de l'algèbre des fonctions quasi-symétriques. Nous montrons ici que Π est fermée sous le coproduit, et est donc une algèbre de Hopf. Nous décrivons aussi le noyau du quotient ci-dessus. D'autre part, dans leurs travaux sur les polynômes de Schubert, Billey et Haiman ont défini une nouvelle classe de fonctions quasi-symétriques: les fonctions quasi-symétrique décalé. Nous montrons que Π est strictement contenue dans l'espace linéaire Ξ des fonctions quasi-symétrique gauchis. Puis nous montrons que Ξ est une coalgèbre et calculons les dimensions des composantes de gauchis n .
© 2002 Elsevier Science B.V. All rights reserved.

PACS: 05E15; 05E05; 05A15; 16W30

Keywords: Pieri; Graded operation; Poset; Quasi-symmetric functions

* Corresponding author. Department of Mathematics, York University, 4700 Keele Street, North York, Ontario, Canada M3J 1P6.

E-mail addresses: nantel.bergeron@mathstat.yorku.ca (N. Bergeron),
mykytiuk@pascal.math.yorku.ca (S. Mykytiuk), sottile@math.wisc.edu (F. Sottile),
steph@pascal.math.yorku.ca (S. van Willigenburg).

¹ Supported in part by NSERC. <http://www.math.yorku.ca/bergeron>

² Supported in part by NSF grant DMS-9701755 and NSERC grant OGP0170279. <http://www.math.wisc.edu/~sottile>

³ Supported in part by the LeverhulmeTrust <http://www-theory.dcs.st-and.ac.uk/~stephvw>

1. Introduction

Schur Q functions first arose in the study of projective representations of S_n [7]. Since then they have appeared in variety of contexts including the representations of Lie superalgebras [8] and cohomology classes dual to Schubert cycles in isotropic Grassmanians [4,6]. While studying the duality between skew Schur P and Q functions and their connection to the Schubert calculus of isotropic flag manifolds, we were led to their quasi-symmetric analogues: the *peak functions* of Stembridge [10]. We show that *the linear span of peak functions is a Hopf algebra* (Theorem 2.2). We also show that these peak functions are contained in the strictly larger set of *shifted quasi-symmetric functions* (Theorem 3.6) introduced by Billey and Haiman [1]. We remark that the quasi-symmetric functions here are not any apparent specialization of the quasi-symmetric q -analogues of Hivert [3].

From extensive calculations, we believe that the set of all shifted quasi-symmetric functions form a Hopf algebra, but at present we can only show that:

The set of all shifted quasi-symmetric functions forms a graded coalgebra whose n th graded component has rank π_n , where π_n is given by the recurrence

$$\pi_n = \pi_{n-1} + \pi_{n-2} + \pi_{n-4}$$

with initial conditions $\pi_1 = 1$, $\pi_2 = 1$, $\pi_3 = 2$, $\pi_4 = 4$.

We shall prove this result (Theorems 3.2 and 4.3) and in addition shall establish some other properties of these functions.

A composition $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$ of a positive integer n is an ordered list of positive integers whose sum is n . We denote this by $\alpha \vDash n$. We call the integers α_i the *components* of α , and denote the number of components in α by $k(\alpha)$. There exists a natural one-to-one correspondence between compositions of n and subsets of $[n-1]$. If $A = \{a_1, a_2, \dots, a_{k-1}\} \subset [n-1]$, where $a_1 < a_2 < \dots < a_{k-1}$, then A corresponds to the composition, $\alpha = [a_1 - a_0, a_2 - a_1, \dots, a_k - a_{k-1}]$, where $a_0 = 0$ and $a_k = n$. For ease of notation, we shall denote the set corresponding to a given composition α by $I(\alpha)$. For compositions α and β , we say that α is a *refinement* of β if $I(\beta) \subset I(\alpha)$, and denote this by $\alpha \preceq \beta$.

For any composition $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$ we denote by M_α , the *monomial quasi-symmetric function* [2]:

$$M_\alpha = \sum_{i_1 < i_2 < \dots < i_k} x_{i_1}^{\alpha_1}, \dots, x_{i_k}^{\alpha_k}.$$

We define $M_0 = 1$, where 0 denotes the unique empty composition of 0. We denote by F_α , the *fundamental quasi-symmetric function* [2]:

$$F_\alpha = \sum_{\alpha \preceq \beta} M_\beta.$$

Definition 1.1. For any subset $A \subset [n - 1]$, let $A + 1$ be the subset of $\{2, \dots, n\}$ formed from A by adding 1 to each element of A . Let $\alpha \vDash n$. Then, we define

$$\theta_\alpha = \sum_{\substack{\beta \vDash n \\ I(\alpha) \subset I(\beta) \cup (I(\beta) + 1)}} 2^{k(\beta)} M_\beta.$$

This is the natural extension of the definition of peak functions given in [10].

Example 1.2. We shall often omit the brackets that surround the components of a composition.

If $\alpha = 21$, then $I(\alpha) = \{2\}$, and $I(\alpha) + 1 = \{3\}$. Hence

$$\theta_{21} = 4M_{21} + 4M_{12} + 8M_{111}.$$

Let Σ^n be the \mathbb{Z} -module of quasi-symmetric functions spanned by $\{M_\alpha\}_{\alpha \vDash n}$ and let $\Sigma = \bigoplus_{n \geq 0} \Sigma^n$ be the graded \mathbb{Z} -algebra of quasi-symmetric functions. This is a Hopf algebra [5] with coproduct given by

$$\Delta(M_\alpha) = \sum_{\alpha = \beta \cdot \gamma} M_\beta \otimes M_\gamma,$$

where $\beta \cdot \gamma$ is the concatenation of compositions β and γ .

Example 1.3. $\Delta(M_{32}) = 1 \otimes M_{32} + M_3 \otimes M_2 + M_{32} \otimes 1$.

We compute the coproduct of the functions θ_α .

Lemma 1.4. For any composition $\alpha \vDash n$ we have that

$$\Delta(\theta_\alpha) = 1 \otimes \theta_\alpha + \sum \theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}, \tag{1}$$

where the sum is over all ways of writing α as $\varepsilon \cdot (a + b) \cdot \zeta$, that is, the concatenation of compositions ε and ζ , and a component of α written as the sum of numbers $a > 0$, $b \geq 0$. Also $\phi(b \cdot \zeta) = [1 + \zeta_1, \zeta_2, \dots]$ if $b = 1$ and $b \cdot \zeta$ otherwise.

We shall use this result to show that certain subsets of functions θ_α span coalgebras (Theorems 2.2 and 3.2).

Proof. Definition 1.1 is equivalent to

$$\theta_\alpha = \sum_{\substack{\beta \vDash n \\ \beta^* \leq \alpha}} 2^{k(\beta)} M_\beta,$$

where β^* is the refinement of β obtained by replacing all components $\beta_i > 1$, for $i > 1$, by $[1, \beta_i - 1]$. Thus, the LHS of Eq. (1) is equal to

$$\sum_{\substack{\beta \vDash n \\ \beta^* \preceq \alpha \\ \beta = \gamma \cdot \delta}} 2^{k(\beta)} M_\gamma \otimes M_\delta = \sum_{\substack{\gamma \cdot \delta \vDash n \\ (\gamma \cdot \delta)^* \preceq \alpha}} 2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta. \quad (2)$$

Let $2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta$ be a term of this sum with $\gamma \vDash m$. If $m = 0$, then the term is $1 \otimes 2^{k(\delta)} M_\delta$, where $\delta^* \preceq \alpha$, and it appears in the summand $1 \otimes \theta_\alpha$ on the RHS of Eq. (1). If $m > 0$, then the term can only appear in one summand on the RHS of Eq. (1), namely $\theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$ with $\varepsilon \cdot a \vDash m$. To show that it does indeed appear, we need to prove that $\gamma^* \preceq \varepsilon \cdot a$ and $\delta^* \preceq \phi(b \cdot \zeta)$. Let δ^{**} be the refinement of δ^* obtained by replacing the part δ_1 by $[1, \delta_1 - 1]$, if $\delta_1 > 1$. We have that

$$\gamma^* \cdot \delta^{**} = (\gamma \cdot \delta)^* \preceq \varepsilon \cdot (a + b) \cdot \zeta$$

which implies that $\gamma^* \preceq \varepsilon \cdot a$, and $\delta^{**} \preceq b \cdot \zeta \preceq \phi(b \cdot \zeta)$.

If $\delta \vDash 0$ or $\delta_1 = 1$, then $\delta^* = \delta^{**} \preceq \phi(b \cdot \zeta)$. However, if $\delta_1 > 1$, then there are two possible cases: either $\delta_1 \leq b$, or $b = 1$ and $\delta_1 - 1 \leq \zeta_1$. In the former case $\delta^* \preceq b \cdot \zeta = \phi(b \cdot \zeta)$, while in the latter $\delta_1 \leq 1 + \zeta_1$, whence $\delta^* \preceq [1 + \zeta_1, \zeta_2, \dots] = \phi(b \cdot \zeta)$. Conversely, it is easy to see that all terms belonging to the tensor $1 \otimes \theta_\alpha$ on the RHS of Eq. (1) also appear in Eq. (2). Now let $2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta$ be a term belonging to a tensor $\theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$ on the RHS of Eq. (1). To show that it appears in Eq. (2) we must prove that $(\gamma \cdot \delta)^* \preceq \varepsilon \cdot (a + b) \cdot \zeta$. We have that $\gamma^* \preceq \varepsilon \cdot a$ and $\delta^* \preceq \phi(b \cdot \zeta)$, which imply that

$$(\gamma \cdot \delta)^* = \gamma^* \cdot \delta^{**} \preceq \gamma^* \cdot \delta^* \preceq \varepsilon \cdot a \cdot \phi(b \cdot \zeta).$$

If $b = 0$ or $b > 1$, then

$$(\gamma \cdot \delta)^* \preceq \varepsilon \cdot a \cdot \phi(b \cdot \zeta) = \varepsilon \cdot a \cdot b \cdot \zeta \preceq \varepsilon \cdot (a + b) \cdot \zeta.$$

If $b = 1$, then $\delta^* \preceq \phi(b \cdot \zeta) = [1 + \zeta_1, \zeta_2, \dots]$ implies that

$$\delta^{**} = [1, \dots] \preceq [1, \zeta_1, \dots] = b \cdot \zeta.$$

Therefore,

$$(\gamma \cdot \delta)^* = \gamma^* \cdot \delta^{**} \preceq \varepsilon \cdot a \cdot b \cdot \zeta \preceq \varepsilon \cdot (a + b) \cdot \zeta$$

as desired.

Finally, we note that no term $2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta$ appears more than once in Eq. (2), or more than once in the expansion of the RHS of Eq. (1). The former is clear, while the latter follows from the fact that if $\theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$ and $\theta_{\varepsilon' \cdot a'} \otimes \theta_{\phi(b' \cdot \zeta')}$ are distinct summands on the RHS of Eq. (1), then $\varepsilon \cdot a \vDash k$ and $\varepsilon' \cdot a' \vDash l$ where $k \neq l$. \square

2. The peak Hopf algebra

Definition 2.1. For any composition $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$ we say that θ_α is a peak function if $\alpha_i = 1 \Rightarrow i = k$.

Observe that if θ_α is a peak function and $\alpha \vDash n$, then $I(\alpha) \subset \{2, \dots, n - 1\}$ such that no two i in $I(\alpha)$ are consecutive.

Let Π^n be the \mathbb{Z} -module spanned by all peak functions θ_α , $\alpha \vDash n$, and let $\Pi = \bigoplus_{n \geq 0} \Pi^n$. This was studied by Stembridge [10] who showed that the peak functions are F-positive, are closed under product, and form a basis for Π , and so the rank of Π^n is the n th Fibonacci number. In addition, we also know the following about the algebra of peaks, Π .

Theorem 2.2. Π is closed under coproduct.

Proof. If all components of a composition α , except perhaps the last, are greater than 1, then the same is true for all compositions $\varepsilon \cdot a$ and $\phi(b \cdot \zeta)$ appearing in the RHS of Eq. (1). \square

Let Θ be the \mathbb{Z} -linear map from Σ to Π defined by $\Theta(F_\alpha) = \theta_{\Lambda(\alpha)}$, where $\Lambda(\alpha)$ is the composition formed from $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$ by adding together adjacent components $\alpha_i, \alpha_{i+1}, \dots, \alpha_{i+j}$, where $\alpha_{i+l} = 1$ for $l = 0, \dots, j - 1$, and either $\alpha_{i+j} \neq 1$, or $i + j = k$.

Example 2.3. If $\alpha = 31125111$ then $\Lambda(\alpha) = 3453$.

Stembridge [10] showed that $\Theta : \Sigma \rightarrow \Pi$ is a graded surjective ring homomorphism, and was an analogue of the retraction from the algebra of symmetric functions to Schur Q functions. It is clear from our proof above that this morphism is in fact a Hopf homomorphism. We can describe the kernel of Θ as follows.

Lemma 2.4. The non-zero differences $F_\alpha - F_{\Lambda(\alpha)}$ form a basis of the kernel of Θ .

Proof. Each difference $F_\alpha - F_{\Lambda(\alpha)}$ is in the kernel of Θ as $\Theta(F_\alpha - F_{\Lambda(\alpha)}) = 0$ since $\Lambda(\Lambda(\alpha)) = \Lambda(\alpha)$. In addition, the non-zero differences are linearly independent as they have different leading terms. Letting f_n denote the n th Fibonacci number, there are $2^{n-1} - f_n$ such differences, and since

$$\begin{aligned} \dim \ker \Theta &= \dim \Sigma^n - \dim \Pi^n \\ &= 2^{n-1} - f_n, \end{aligned}$$

our result follows. \square

and $G(3^{(3)})$ is



hence $G(\bigwedge_{i \in S} 3^{(i)})$ is



so $\bigwedge_{i \in S} 3^{(i)}$ is the composition 4.

Definition 3.5. (Billey and Haiman [1]). Let α be a composition of n . Let $\mathcal{A}(I(\alpha))$ denote the set of all sequences $j_1 \leq j_2 \leq \dots \leq j_n$ in \mathbb{N} such that we do not have $j_{i-1} = j_i = j_{i+1}$, for any $i \in I(\alpha)$. The shifted quasi-symmetric function $\theta_\alpha^{\text{BH}}$ is given by

$$\theta_\alpha^{\text{BH}} = \sum_{\substack{J=(j_1, \dots, j_n) \\ j_1 \leq \dots \leq j_n \\ J \in \mathcal{A}(I(\alpha))}} 2^{|j|} x_{j_1} \dots x_{j_n},$$

where $|j|$ denotes the number of distinct values j_i in J .

Theorem 3.6. For any sqs-function θ_α we have that $\theta_\alpha = \theta_\alpha^{\text{BH}}$.

Proof. For each $i \in I(\alpha) \subset [n - 1]$, $j_{i-1} = j_i = j_{i+1}$ is forbidden in any monomial

$$x_{j_1} x_{j_2} \dots x_{j_i} \dots x_{j_n}$$

appearing as a summand of the function $\theta_\alpha^{\text{BH}}$. This is equivalent to saying that M_β is a summand of $\theta_\alpha^{\text{BH}}$ if and only if $G(3^{(i)}) \not\subseteq G(\beta)$ for all $i \in I(\alpha)$. Therefore, at least one of $i - 1$ or i must be the largest label of a vertex in a connected component in $G(\beta)$.

Now, when going from compositions of n to subsets of $[n - 1]$ we can do so using our graphs, G . All we have to do is list the label of the vertex that is the largest in each connected component, not listing n . We call these vertices the *end-points*. We are now in a position to prove the equivalence of Definitions 1.1 and 3.5 for sqs-functions. The powers of 2 agree so we need only show that the indices of summation do too. To see this, take any sqs-function θ_α and let $i \in I(\alpha)$. Then M_β is a summand in $\theta_\alpha^{\text{BH}}$ if at least one of $i - 1$ or i is an end-point in $G(\beta)$. Therefore, i or $i - 1$ belongs to $I(\beta)$, and M_β is a summand of θ_α . Conversely, if M_β is a summand of θ_α , then this implies that for each $i \in I(\alpha)$, we have that $i - 1$ or i belongs to $I(\beta)$, so one of $i - 1$ or i is an end-point in $G(\beta)$, so M_β is a summand of $\theta_\alpha^{\text{BH}}$. \square

4. A basis for Ξ

Definition 4.1. Let θ_α be an sqs-function and $\alpha \vDash n$. We define an internal peak $i \in I(\alpha)$ such that $i - 1, i + 1 \notin I(\alpha)$, and $i \in \{3, \dots, n - 2\}$.

Remark. Observe that the occurrence of an internal peak in the i th position in $I(\alpha) = \{w_1, w_2, \dots\}$, where $w_1 < w_2 < \dots$, is equivalent to having two components of α , say α_i, α_{i+1} such that $\alpha_{i+1} \geq 2$, and $\alpha_i \geq 2$ if $i \neq 1$, or $\alpha_i \geq 3$ if $i = 1$.

We can now describe the basis of Ξ as follows.

Theorem 4.2. *The coalgebra Ξ has a basis consisting of all sqs-functions θ_α where $I(\alpha)$ contains no internal peak.*

We sketch the proof of Theorem 4.2 later.

Theorem 4.3. *The rank of Ξ^n is given by the recurrence*

$$\pi_n = \pi_{n-1} + \pi_{n-2} + \pi_{n-4}$$

with initial conditions $\pi_1 = 1, \pi_2 = 1, \pi_3 = 2, \pi_4 = 4$.

This recurrence was suggested by a superseeker query [9].

Proof. By direct calculation we obtain that $\pi_1 = 1, \pi_2 = 1, \pi_3 = 2$, and $\pi_4 = 4$.

To obtain our recurrence, we observe that for each sqs-function, θ_α where $\alpha \vDash n$, we can encode $I(\alpha)$ as a binary word of length $n - 2$, by placing a 1 in position $i - 1$ if i is contained in $I(\alpha)$, and 0 otherwise. By this one-to-one correspondence, we see that $I(\alpha)$ contains no internal peak if its corresponding binary word does not contain 010 as a subword.

We therefore, count binary words of length n that avoid the subword 010. Appending either 1 or 0 to such a binary word of length $n - 1$ gives one of length n , provided that we have not created the subword 010 in the last three positions. Let a_n, b_n, c_n , and d_n enumerate those binary words of length $n - 2$ that avoid the subword 010 and end in, respectively, 00, 01, 10, and 11. We then obtain the following four simultaneous recursions.

$$a_n = a_{n-1} + c_{n-1}, \quad b_n = a_{n-1} + c_{n-1}, \quad c_n = d_{n-1}, \quad d_n = b_{n-1} + d_{n-1}.$$

Clearly, the number of $I(\alpha)$ s in $[n - 1]$ with no internal peaks is given by

$$\pi_n = a_n + b_n + c_n + d_n.$$

However, by substituting in our recurrences we obtain

$$\begin{aligned}
 \pi_n &= a_n + b_n + c_n + d_n \\
 &= 2a_{n-1} + b_{n-1} + 2c_{n-1} + 2d_{n-1} \\
 &= \pi_{n-1} + a_{n-1} + c_{n-1} + d_{n-1} \\
 &= \pi_{n-1} + a_{n-2} + b_{n-2} + c_{n-2} + 2d_{n-2} \\
 &= \pi_{n-1} + \pi_{n-2} + d_{n-2} \\
 &= \pi_{n-1} + \pi_{n-2} + b_{n-3} + d_{n-3} \\
 &= \pi_{n-1} + \pi_{n-2} + a_{n-4} + b_{n-4} + c_{n-4} + d_{n-4} \\
 &= \pi_{n-1} + \pi_{n-2} + \pi_{n-4}. \quad \square
 \end{aligned}$$

We say that M_β is a maximal term of θ_x if for any γ higher in the partial order of compositions M_γ is not a summand of θ_x . The following lemma is stated without proof.

Lemma 4.4. *Let θ_x be an sqs-function. Consider the collection S of all possible sets derived from $I(x)$ by adding either $i-1$ or $i+1$ to $I(x)$ for all internal peaks $i \in I(x)$. If M_β is a maximal term of θ_x , then β is derived from*

$$\bigwedge_{\substack{i \in I(\tilde{x})^c \\ I(\tilde{x}) \in S}} 3^{(i)}$$

by adding adjacent components equal to 1 together to give a component equal to 2 as often as possible.

Lemma 4.5. *Let θ_x be an sqs-function, and let $I(x)$ have an internal peak in the j th position, then we have the following linear relation:*

$$\begin{aligned}
 \theta_x &= \theta_{[\alpha_1, \dots, \alpha_j - 1, 1, \alpha_{j+1}, \dots, \alpha_k]} + \theta_{[\alpha_1, \dots, \alpha_j, 1, \alpha_{j+1} - 1, \dots, \alpha_k]} \\
 &\quad - \theta_{[\alpha_1, \dots, \alpha_j - 1, 1, 1, \alpha_{j+1} - 1, \dots, \alpha_k]}.
 \end{aligned}$$

Proof. By Definition 3.5 we have that the leading terms of θ_x determine the other summands that belong to θ_x . Hence, by Lemma 4.4 it follows that the summands of θ_x will be the union of the summands of $\theta_{[\alpha_1, \dots, \alpha_j - 1, 1, \alpha_{j+1}, \dots, \alpha_k]}$ and $\theta_{[\alpha_1, \dots, \alpha_j, 1, \alpha_{j+1} - 1, \dots, \alpha_k]}$. However, those summands that appear in both will be duplicated. By definition these will be the summands of $\theta_{[\alpha_1, \dots, \alpha_j - 1, 1, 1, \alpha_{j+1} - 1, \dots, \alpha_k]}$, and the result follows. \square

Proof of Theorem 4.2. From our relation in Lemma 4.5, it follows that any θ_x can be rewritten as a linear combination of functions $\theta_{\tilde{x}}$, where $I(\tilde{x})$ contains no internal peaks. In addition, by Lemma 4.4 and Definition 3.5 we have that the set of all sqs-functions

θ_x where $I(\alpha)$ contains no internal peaks is linearly independent and thus forms a basis for Ξ . \square .

Acknowledgements

The authors are grateful to Bruce Sagan for his helpful comments.

References

- [1] S. Billey, M. Haiman, Schubert polynomials for the classical groups, *J. AMS* 8 (1995) 443–482.
- [2] I. Gessel, Multipartite P -partitions and inner products of skew Schur functions, *Contemp. Math.* 34 (1984) 289–301.
- [3] F. Hivert, Quasi-symmetric functions and Hecke algebra action, *Proceeding of the 10th FPSAC, Toronto, 1998*, pp. 377–388.
- [4] T. Józefiak, Schur Q functions and cohomology of isotropic Grassmanians, *Math. Proc. Cambridge Philos. Soc.* 109 (1991) 471–478.
- [5] C. Malvenuto, C. Reutenauer, Duality between quasi-symmetric functions and the Solomon descent algebra, *J. Algebra* 177 (1995) 967–982.
- [6] P. Pragacz, Algebro-Geometric Applications of Schur S - and Q - Polynomials, *Topics in Invariant Theory, Séminaire d'Algèbre Dubreil–Malliavin 1989–1990*, Springer, Berlin, 1991, pp. 130–191.
- [7] I. Schur, Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen, *J. Reine Angew. Math.* 139 (1911) 155–250.
- [8] A.N. Sergeev, The tensor algebra of the identity representation as a module over the Lie superalgebras $gl(n, m)$ and $Q(n)$, *Math. USSR Sbornik* 51 (1985) 419–427.
- [9] N.J.A. Sloane, An on-line version of the Encyclopedia of integer sequences, *Electron. J. Combin.* 1 (1994) F1. <http://akpublic.research.att.com/~njas/sequences/ol.html>.
- [10] J. Stembridge, Enriched P -partitions, *Trans. AMS* 349 (1997) 763–788.