

Course Outline for Mathematics 257/316 (3 credits) Term 1, Sept.-Dec., 2018

Partial Differential Equations

<u>Prerequisites:</u>	One of Math 215, 255, 265.
<u>Credit:</u>	3 Credits. Credit only given for one of Math 256, 257, 316.
<u>Instructor:</u>	Anthony Peirce, <u>Office:</u> Mathematics Building 108
<u>Home Page:</u>	http://www.math.ubc.ca/~peirce
<u>Office Hours:</u>	Monday: 10-11 am, Wed: 3-3:55 pm, Fri: 10-11 am.
<u>Assessment:</u>	The final grades will be based on homework (10%) (including EXCEL/MATLAB projects), two in class midterm exams (40%) and one final exam (50%). Assignments are to be submitted in hard-copy from at the designated class – no late assignments can be accepted. There will be no make-up midterms. A student must get at least 35% on the final exam to pass this course.
<u>Test Dates:</u>	Wednesday, October 17 th , Wednesday, November 14 th .
<u>Text:</u> (recommended but not required)	Elementary Differential Equations and Boundary Value Problems (10 th Ed), W.E. Boyce & R.C. DiPrima (John Wiley & Sons) 2012
<u>Other References:</u>	1. Applied Partial Differential Equations with Fourier Series and Boundary Value Problems (4 th Ed), R. Haberman, (Pearson), 2004. 3. http://www.math.ubc.ca/~rfroese/notes/Lecs316.pdf , Richard Froese, Partial Differential Equations, UBC M257/316 lecture notes free on the web.

<u>Topics:</u>	<u>Approx Time</u>
1. Review of techniques to solve ODEs	1 hr
2. Series Solutions of variable coefficient ODEs (Chapter 5)	
a. Series solutions at ordinary points (5.1-5.3)	3 hrs
b. Regular singular points (5.4-5.7, 5.8 briefly)	4 hrs
3. Introduction to Partial differential equations (Chapter 10)	
The heat equation (10.5), the wave equation (10.7), Laplace's equation (10.8)	2 hrs
4. Introduction to numerical methods for PDEs using spread sheets	3 hrs
a. First and second derivative approximations using finite differences - errors	
b. Explicit finite difference schemes for the heat equation	
• Stability and derivative boundary conditions	
c. Explicit finite difference schemes for the wave equation	
d. Finite difference approximation of Laplace's Equation – iterative methods	
5. Fourier Series and Separation of Variables (Chapter 10)	
a. The heat equation and Fourier Series (10.1-10.6)	9 hrs
b. The wave equation (10.7)	3 hrs
c. Laplace's equation (10.8)	5 hrs
6. Boundary Value Problems and Sturm-Liouville Theory (Chapter 11)	
a. Eigenfunctions and eigenvalues (11.1)	1 hr
b. Sturm-Liouville boundary value problems (11.2)	1 hr
c. Nonhomogeneous boundary value problems (11.3)	2 hrs
	Tests <u>2 hrs</u>
	36 hrs

Math 257-316 PDE Formula sheet - final exam

Trigonometric and Hyperbolic Function identities

$$\begin{aligned}\sin(\alpha \pm \beta) &= \sin \alpha \cos \beta \pm \sin \beta \cos \alpha & \sin^2 t + \cos^2 t &= 1 \\ \cos(\alpha \pm \beta) &= \cos \alpha \cos \beta \mp \sin \beta \sin \alpha & \sin^2 t &= \frac{1}{2}(1 - \cos(2t)) \\ \sinh(\alpha \pm \beta) &= \sinh \alpha \cosh \beta \pm \sinh \beta \cosh \alpha & \cosh^2 t - \sinh^2 t &= 1 \\ \cosh(\alpha \pm \beta) &= \cosh \alpha \cosh \beta \pm \sinh \beta \sinh \alpha & \sinh^2 t &= \frac{1}{2}(\cosh(2t) - 1)\end{aligned}$$

Basic linear ODE's with real coefficients

	constant coefficients	Euler eq
ODE	$ay'' + by' + cy = 0$	$ax^2y'' + bxy' + cy = 0$
indicial eq.	$ar^2 + br + c = 0$	$ar(r-1) + br + c = 0$
$r_1 \neq r_2$ real	$y = Ae^{r_1x} + Be^{r_2x}$	$y = Ax^{r_1} + Bx^{r_2}$
$r_1 = r_2 = r$	$y = Ae^{rx} + Bxe^{rx}$	$y = Ax^r + Bx^r \ln x $
$r = \lambda \pm i\mu$	$e^{\lambda x}[A \cos(\mu x) + B \sin(\mu x)]$	$x^\lambda[A \cos(\mu \ln x) + B \sin(\mu \ln x)]$

Series solutions for $y'' + p(x)y' + q(x)y = 0$ (*) around $x = x_0$.

Ordinary point x_0 : Two linearly independent solutions of the form:

$$y(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n$$

Regular singular point x_0 : Rearrange (*) as:

$$(x - x_0)^2 y'' + [(x - x_0)p(x)](x - x_0)y' + [(x - x_0)^2 q(x)]y = 0$$

If $r_1 > r_2$ are roots of the indicial equation: $r(r-1) + br + c = 0$ where

$$b = \lim_{x \rightarrow x_0} (x - x_0)p(x) \text{ and } c = \lim_{x \rightarrow x_0} (x - x_0)^2 q(x) \text{ then a solution of (*) is}$$

$$y_1(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^{n+r_1} \text{ where } a_0 = 1.$$

The second linearly independent solution y_2 is of the form:

Case 1: If $r_1 - r_2$ is neither 0 nor a positive integer:

$$y_2(x) = \sum_{n=0}^{\infty} b_n(x - x_0)^{n+r_2} \text{ where } b_0 = 1.$$

Case 2: If $r_1 - r_2 = 0$:

$$y_2(x) = y_1(x) \ln(x - x_0) + \sum_{n=1}^{\infty} b_n(x - x_0)^{n+r_2} \text{ for some } b_1, b_2, \dots$$

Case 3: If $r_1 - r_2$ is a positive integer:

$$y_2(x) = ay_1(x) \ln(x - x_0) + \sum_{n=0}^{\infty} b_n(x - x_0)^{n+r_2} \text{ where } b_0 = 1.$$

Fourier, sine and cosine series

Let $f(x)$ be defined in $[-L, L]$ then its Fourier series $Ff(x)$ is a $2L$ -periodic function on \mathbf{R} : $Ff(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \{a_n \cos(\frac{n\pi x}{L}) + b_n \sin(\frac{n\pi x}{L})\}$

where $a_n = \frac{1}{L} \int_{-L}^L f(x) \cos(\frac{n\pi x}{L}) dx$ and $b_n = \frac{1}{L} \int_{-L}^L f(x) \sin(\frac{n\pi x}{L}) dx$

Theorem (Pointwise convergence) If $f(x)$ and $f'(x)$ are piecewise continuous, then $Ff(x)$ converges for every x to $\frac{1}{2}[f(x-) + f(x+)]$.

Parseval's identity

$$\frac{1}{L} \int_{-L}^L |f(x)|^2 dx = \frac{|a_0|^2}{2} + \sum_{n=1}^{\infty} (|a_n|^2 + |b_n|^2).$$

For $f(x)$ defined in $[0, L]$, its cosine and sine series are

$$Cf(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos(\frac{n\pi x}{L}), \quad a_n = \frac{2}{L} \int_0^L f(x) \cos(\frac{n\pi x}{L}) dx,$$

$$Sf(x) = \sum_{n=1}^{\infty} b_n \sin(\frac{n\pi x}{L}), \quad b_n = \frac{2}{L} \int_0^L f(x) \sin(\frac{n\pi x}{L}) dx.$$

D'Alembert's solution to the wave equation

PDE: $u_{tt} = c^2 u_{xx}$, $-\infty < x < \infty$, $t > 0$ **IC:** $u(x, 0) = f(x)$, $u_t(x, 0) = g(x)$.

SOLUTION: $u(x, t) = \frac{1}{2}[f(x+ct) + f(x-ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds$

Sturm-Liouville Eigenvalue Problems

ODE: $[p(x)y']' - q(x)y + \lambda r(x)y = 0$, $a < x < b$.

BC: $\alpha_1 y(a) + \alpha_2 y'(a) = 0$, $\beta_1 y(b) + \beta_2 y'(b) = 0$.

Hypothesis: p, p', q, r continuous on $[a, b]$. $p(x) > 0$ and $r(x) > 0$ for $x \in [a, b]$. $\alpha_1^2 + \alpha_2^2 > 0$. $\beta_1^2 + \beta_2^2 > 0$.

Properties (1) The differential operator $Ly = [p(x)y']' - q(x)y$ is symmetric in the sense that $(f, Lg) = (Lf, g)$ for all f, g satisfying the BC, where $(f, g) = \int_a^b f(x)g(x) dx$. (2) All eigenvalues are real and can be ordered as $\lambda_1 < \lambda_2 < \dots < \lambda_n < \dots$ with $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$, and each eigenvalue admits a unique (up to a scalar factor) eigenfunction ϕ_n .

(3) **Orthogonality:** $(\phi_m, r\phi_n) = \int_a^b \phi_m(x)\phi_n(x)r(x) dx = 0$ if $\lambda_m \neq \lambda_n$.

(4) **Expansion:** If $f(x) : [a, b] \rightarrow \mathbf{R}$ is square integrable, then

$$f(x) = \sum_{n=1}^{\infty} c_n \phi_n(x), \quad a < x < b, \quad c_n = \frac{\int_a^b f(x)\phi_n(x)r(x) dx}{\int_a^b \phi_n^2(x)r(x) dx}, \quad n = 1, 2, \dots$$