

Planetary Motion

with Corrections from General Relativity

Let $\mathbf{r}(t)$ be the position at time t of a planet (approximated by a point mass, m) in orbit around a sun (also approximated by a point mass, M) whose position is fixed at the origin. According to Newton's law of gravity

$$m\mathbf{r}''(t) = -\frac{GMm}{|\mathbf{r}|^3}\mathbf{r} \quad (1)$$

where G is the usual gravitational constant.

It is possible to considerably simplify (1). The first simplification is a consequence of the fact that (1) is a central force law. That is, the force $-\frac{GMm}{|\mathbf{r}|^3}\mathbf{r}$ is always parallel to the radius vector \mathbf{r} . For all solutions $\mathbf{r}(t)$ of all central force laws, $m\mathbf{r}'' = f(\mathbf{r})\mathbf{r}$, the angular momentum $\mathbf{a}(t) = m\mathbf{r}(t) \times \mathbf{r}'(t)$ is independent of t . To see this, it suffices to observe that the time derivative

$$\frac{d\mathbf{a}}{dt} = \frac{d}{dt}m\mathbf{r} \times \mathbf{r}' = m\mathbf{r}' \times \mathbf{r}' + m\mathbf{r} \times \mathbf{r}'' = m\mathbf{r}' \times \mathbf{r}' + mf(\mathbf{r})\mathbf{r} \times \mathbf{r}$$

is always zero, because $\mathbf{v} \times \mathbf{v} = 0$ for all vectors \mathbf{v} . Consequently, for all t , $\mathbf{r}(t)$ is perpendicular to the fixed vector \mathbf{a} . In other words $\mathbf{r}(t)$ lies in a fixed plane, for all t . We may as well choose our coordinate system so that it is the x - y plane. That is the first simplification.

The second simplification is achieved by switching to polar coordinates and writing

$$\begin{aligned} \mathbf{r}(t) &= r(t)(\cos \theta(t), \sin \theta(t)) \\ \mathbf{r}'(t) &= r'(t)(\cos \theta(t), \sin \theta(t)) + r(t)\theta'(t)(-\sin \theta(t), \cos \theta(t)) \\ \mathbf{r}''(t) &= [r''(t) - r(t)\theta'(t)^2](\cos \theta(t), \sin \theta(t)) + [2r'(t)\theta'(t) + r(t)\theta''(t)](-\sin \theta(t), \cos \theta(t)) \end{aligned} \quad (2)$$

Substituting (2) into (1) gives

$$m[r'' - r\theta'^2](\cos \theta, \sin \theta) + [2r'\theta' + r\theta''](-\sin \theta, \cos \theta) = -\frac{GMm}{r^2}(\cos \theta, \sin \theta)$$

matching coefficients of $(\cos \theta, \sin \theta)$ on the left and right hand sides and then matching coefficients of $(-\sin \theta(t), \cos \theta(t))$ on the left and right hand sides gives

$$m[r'' - r\theta'^2] = -\frac{GMm}{r^2} \quad (3a)$$

$$2r'\theta' + r\theta'' = 0 \quad (3b)$$

In fact (3b) is redundant with conservation of angular momentum. Since $(\cos \theta(t), \sin \theta(t)) \times (\cos \theta(t), \sin \theta(t)) = 0$ and $(\cos \theta(t), \sin \theta(t)) \times (-\sin \theta(t), \cos \theta(t))$ is the unit vector $\hat{\mathbf{k}}$ along the z -axis, the angular momentum

$$\mathbf{a}(t) = m\mathbf{r}(t) \times \mathbf{r}'(t) = mr(t)^2\theta'(t)\hat{\mathbf{k}}$$

and conservation of angular momentum implies that

$$r(t)^2\theta'(t) = \frac{l}{m} \quad (4)$$

where l is the constant magnitude of the angular momentum vector \mathbf{a} . Differentiating (4) with respect to t and multiplying by r gives (3b). We can use (4) to eliminate the θ' in (3a)

$$r'' - \frac{l^2}{m^2 r^3} = -\frac{GM}{r^2} \quad (5)$$

In general relativity (see Misner, Thorne and Wheeler, *Gravitation*, page 656) this is modified to

$$r'' - \frac{l^2}{m^2 r^3} = -\frac{GM}{r^2} \left(1 + \frac{3l^2}{m^2 c^2 |r|^2}\right) \quad (6)$$

assuming that the planet is moving slowly compared to the speed c of light.

The final simplification is another change of variables. Replace r by $u = \frac{1}{r}$ and think of u as being a function of θ , which in turn is a function of t . That is

$$\begin{aligned} r(t) &= \frac{1}{u(\theta(t))} \\ r'(t) &= -\frac{1}{u(\theta(t))^2} \frac{du}{d\theta}(\theta(t))\theta'(t) = -r(t)^2\theta'(t) \frac{du}{d\theta}(\theta(t)) = -\frac{l}{m} \frac{du}{d\theta}(\theta(t)) \quad \text{by (4)} \\ r''(t) &= -\frac{l}{m} \frac{d^2 u}{d\theta^2}(\theta(t))\theta'(t) = -\frac{l}{m} \frac{l}{mr^2} \frac{d^2 u}{d\theta^2} = -\frac{l^2}{m^2} u^2 \frac{d^2 u}{d\theta^2} \end{aligned}$$

Substituting this into (6) gives

$$-\frac{l^2}{m^2} u^2 \frac{d^2 u}{d\theta^2} - \frac{l^2}{m^2} u^3 = -GMu^2 \left(1 + \frac{3l^2}{m^2 c^2} u^2\right)$$

Multiplying through by $-\frac{m^2}{l^2 u^2}$ gives

$$\boxed{\frac{d^2 u}{d\theta^2} + u = \frac{GMm^2}{l^2} + \frac{3GM}{c^2} u^2}$$