


The Variable Length Pendulum

Model a pendulum by a mass m that is connected to a hinge by an idealized rod that is massless and of time-dependent length $\ell(t)$. Denote by θ the angle between the rod and



vertical. At time t , the position and velocity of the mass are

$$\begin{aligned} x(t) &= \ell(t) \sin \theta(t) & \dot{x}(t) &= \dot{\ell}(t) \sin \theta(t) + \ell(t) \dot{\theta}(t) \cos \theta(t) \\ y(t) &= -\ell(t) \cos \theta(t) & \dot{y}(t) &= -\dot{\ell}(t) \cos \theta(t) + \ell(t) \dot{\theta}(t) \sin \theta(t) \end{aligned}$$

So the kinetic and potential energies of the mass are

$$\begin{aligned} T &= \frac{1}{2}m[\dot{x}(t)^2 + \dot{y}(t)^2] = \frac{1}{2}m[\dot{\ell}(t)^2 + \ell(t)^2\dot{\theta}(t)^2] \\ U &= mgy(t) = -mg\ell(t) \cos \theta(t) \end{aligned}$$

The corresponding Lagrangian is

$$\begin{aligned} \mathcal{L}(\theta, \dot{\theta}, t) &= T - U = \frac{1}{2}m[\dot{\ell}(t)^2 + \ell(t)^2\dot{\theta}^2] + mg\ell(t) \cos \theta \\ &= m\left[\frac{1}{2}\ell(t)^2\dot{\theta}^2 + g\ell(t) \cos \theta + \frac{1}{2}\dot{\ell}(t)^2\right] \end{aligned}$$

and Lagrange's equation of motion is

$$\begin{aligned} 0 &= \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}} \right) - \frac{\partial \mathcal{L}}{\partial \theta} \\ &= \frac{d}{dt} m[\ell(t)^2 \dot{\theta}] - m[-g\ell(t) \sin \theta] \\ &= m[\ell(t)^2 \ddot{\theta}(t) + 2\ell(t)\dot{\ell}(t)\dot{\theta}(t) + g\ell(t) \sin \theta(t)] \end{aligned}$$

or

$$\frac{d^2\theta}{dt^2}(t) + 2\frac{\dot{\ell}(t)}{\ell(t)}\dot{\theta}(t) + \frac{g}{\ell(t)} \sin \theta(t) = 0$$

In particular, if $\ell(t) = L + A \cos(\omega t)$, the equation of motion is

$$\frac{d^2\theta}{dt^2}(t) - 2A\omega \frac{\sin(\omega t)}{L + A \cos(\omega t)} \dot{\theta}(t) + \frac{g}{L + A \cos(\omega t)} \sin \theta(t) = 0$$