

Integration of $\sec \theta$, $\csc \theta$, $\sec^3 \theta$ and $\csc^3 \theta$

$\int \sec \theta d\theta$ – by trickery

The standard trick used to integrate $\sec \theta$ is to multiply the integrand by $1 = \frac{\sec \theta + \tan \theta}{\sec \theta + \tan \theta}$ and then substitute $y = \sec \theta + \tan \theta$, $dy = (\sec \theta \tan \theta + \sec^2 \theta) d\theta$.

$$\begin{aligned}\int \sec \theta d\theta &= \int \sec \theta \frac{\sec \theta + \tan \theta}{\sec \theta + \tan \theta} d\theta = \int \frac{\sec^2 \theta + \sec \theta \tan \theta}{\sec \theta + \tan \theta} d\theta = \int \frac{dy}{y} = \ln |y| + C \\ &= \ln |\sec \theta + \tan \theta| + C\end{aligned}$$

$\int \sec \theta d\theta$ – by partial fractions

Another method for integrating $\int \sec \theta d\theta$, that is more tedious, but less dependent on a memorized trick, is to convert $\int \sec \theta d\theta$ into the integral of a rational function using the substitution $y = \sin \theta$, $dy = \cos \theta d\theta$ and then use partial fractions.

$$\begin{aligned}\int \sec \theta d\theta &= \int \frac{1}{\cos \theta} d\theta = \int \frac{\cos \theta}{\cos^2 \theta} d\theta = \int \frac{\cos \theta}{1 - \sin^2 \theta} d\theta = \int \frac{1}{1 - y^2} dy \\ &= \int \frac{1}{2} \left[\frac{1}{1+y} + \frac{1}{1-y} \right] dy = \frac{1}{2} \int \left[\frac{1}{y+1} - \frac{1}{y-1} \right] dy = \frac{1}{2} \left[\ln |y+1| - \ln |y-1| \right] + C \\ &= \frac{1}{2} \ln \left| \frac{y+1}{y-1} \right| + C = \frac{1}{2} \ln \left| \frac{\sin \theta + 1}{\sin \theta - 1} \right| + C\end{aligned}$$

To see that this answer is really the same as the one above, note that

$$\begin{aligned}\frac{\sin \theta + 1}{\sin \theta - 1} &= \frac{(\sin \theta + 1)^2}{\sin^2 \theta - 1} = \frac{(\sin \theta + 1)^2}{-\cos^2 \theta} \\ \Rightarrow \frac{1}{2} \ln \left| \frac{\sin \theta + 1}{\sin \theta - 1} \right| &= \frac{1}{2} \ln \left| \frac{(\sin \theta + 1)^2}{-\cos^2 \theta} \right| = \frac{1}{2} \ln \left| \frac{(\sin \theta + 1)^2}{\cos^2 \theta} \right| = \ln \left| \frac{\sin \theta + 1}{\cos \theta} \right| = \ln |\tan \theta + \sec \theta|\end{aligned}$$


$\int \csc \theta d\theta$ – by the $x = \tan \frac{\theta}{2}$ substitution

The integral $\int \csc \theta d\theta$ may also be evaluated by both the methods above. The answers are

$$\int \csc \theta d\theta = \ln |\cot \theta - \csc \theta| + C = -\frac{1}{2} \ln \left| \frac{\cos \theta + 1}{\cos \theta - 1} \right| + C$$

Since $\csc \theta$ is a rational function of $\sin \theta$ and $\cos \theta$, the substitution

$$\begin{aligned}x &= \tan \frac{\theta}{2} & \theta &= 2 \tan^{-1} x \\ d\theta &= \frac{2}{1+x^2} dx \\ \sin \theta &= 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} = 2 \frac{x}{\sqrt{1+x^2}} \frac{1}{\sqrt{1+x^2}} = \frac{2x}{1+x^2} \\ \cos \theta &= \cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2} = \frac{1}{1+x^2} - \frac{x^2}{1+x^2} = \frac{1-x^2}{1+x^2}\end{aligned}$$



converts $\int \csc \theta d\theta$ into the integral of a rational function.

$$\int \csc \theta d\theta = \int \frac{1}{\sin \theta} d\theta = \int \frac{1+x^2}{2x} \frac{2}{1+x^2} dx = \int \frac{1}{x} dx = \ln |x| + C = \ln \left| \tan \frac{\theta}{2} \right| + C$$

To see that this answer is really the same as the one above, note that

$$\cot \theta - \csc \theta = \frac{\cos \theta - 1}{\sin \theta} = \frac{-2 \sin^2(\theta/2)}{2 \sin(\theta/2) \cos(\theta/2)} = -\tan \frac{\theta}{2}$$

$\int \sec^3 x dx$ – **by trickery**

The standard trick used to evaluate $\int \sec^3 x dx$ is integration by parts with $u = \sec x$, $dv = \sec^2 x dx$, $du = \sec x \tan x dx$, $v = \tan x$.

$$\int \sec^3 x dx = \int \sec x \sec^2 x dx = \sec x \tan x - \int \tan x \sec x \tan x dx$$

Since $\tan^2 x + 1 = \sec^2 x$, we have $\tan^2 x = \sec^2 x - 1$ and

$$\int \sec^3 x dx = \sec x \tan x - \int [\sec^3 x - \sec x] dx = \sec x \tan x + \ln |\sec x + \tan x| + C - \int \sec^3 x dx$$

where we used $\int \sec x dx = \ln |\sec x + \tan x| + C$. Now moving the $\int \sec^3 x dx$ from the right hand side to the left hand side

$$\begin{aligned} 2 \int \sec^3 x dx &= \sec x \tan x + \ln |\sec x + \tan x| + C \\ \Rightarrow \int \sec^3 x dx &= \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C \end{aligned}$$

for a new arbitrary constant C .

$\int \sec^3 x dx$ – **by partial fractions**

Another method for integrating $\int \sec^3 x dx$, that is more tedious, but less dependent on trickery, is to convert $\int \sec^3 x dx$ into the integral of a rational function using the substitution $y = \sin x$, $dy = \cos x dx$ and then use partial fractions.

$$\begin{aligned} \int \sec^3 x dx &= \int \frac{1}{\cos^3 x} dx = \int \frac{\cos x}{\cos^4 x} dx = \int \frac{\cos x}{[1 - \sin^2 x]^2} dx = \int \frac{1}{[1 - y^2]^2} dy = \int \frac{1}{[y^2 - 1]^2} dy \\ &= \int \left[\frac{1}{2} \left(\frac{1}{y-1} - \frac{1}{y+1} \right) \right]^2 dy = \frac{1}{4} \int \left[\frac{1}{(y-1)^2} - \frac{2}{(y-1)(y+1)} + \frac{1}{(y+1)^2} \right] dy \\ &= \frac{1}{4} \int \left[\frac{1}{(y-1)^2} - \frac{1}{y-1} + \frac{1}{y+1} + \frac{1}{(y+1)^2} \right] dy \\ &= \frac{1}{4} \left[-\frac{1}{y-1} - \ln |y-1| + \ln |y+1| - \frac{1}{y+1} \right] + C = -\frac{1}{4} \frac{2y}{y^2-1} + \frac{1}{4} \ln \left| \frac{y+1}{y-1} \right| + C \\ &= \frac{1}{2} \frac{y}{1-y^2} + \frac{1}{4} \ln \left| \frac{y+1}{y-1} \right| + C = \frac{1}{2} \frac{\sin x}{\cos^2 x} + \frac{1}{2} \ln \left| \frac{\sin x + 1}{\sin x - 1} \right| + C \end{aligned}$$

$\int \csc^3 \theta \, d\theta$ – by the $x = \tan \frac{\theta}{2}$ substitution

As another example of the $x = \tan \frac{\theta}{2}$, $d\theta = \frac{2}{1+x^2} dx$, $\sin \theta = \frac{2x}{1+x^2}$, $\cos \theta = \frac{1-x^2}{1+x^2}$ substitution, we evaluate

$$\begin{aligned} \int \csc^3 \theta \, d\theta &= \int \frac{1}{\sin^3 \theta} \, d\theta = \int \left(\frac{1+x^2}{2x}\right)^3 \frac{2}{1+x^2} dx = \frac{1}{4} \int \frac{1+2x^2+x^4}{x^3} dx \\ &= \frac{1}{4} \left[\frac{x^{-2}}{-2} + 2 \ln |x| + \frac{x^2}{2} \right] + C = \frac{1}{8} \left[-\cot^2 \frac{\theta}{2} + 4 \ln \left| \tan \frac{\theta}{2} \right| + \tan^2 \frac{\theta}{2} \right] + C \end{aligned}$$

By the usual double angle formulae

$$\begin{aligned} \tan^2 \frac{\theta}{2} - \cot^2 \frac{\theta}{2} &= \frac{\sin^2 \frac{\theta}{2}}{\cos^2 \frac{\theta}{2}} - \frac{\cos^2 \frac{\theta}{2}}{\sin^2 \frac{\theta}{2}} = \frac{\sin^4 \frac{\theta}{2} - \cos^4 \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} \cos^2 \frac{\theta}{2}} \\ &= \frac{\sin^2 \frac{\theta}{2} - \cos^2 \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} \cos^2 \frac{\theta}{2}} \quad \text{since } \sin^2 \frac{\theta}{2} + \cos^2 \frac{\theta}{2} = 1 \\ &= \frac{-\cos \theta}{\frac{1}{4} \sin^2 \theta} \end{aligned}$$

so we may also write

$$\int \csc^3 \theta \, d\theta = -\frac{1}{2} \cot \theta \csc \theta + \frac{1}{2} \ln |\cot \theta - \csc \theta| + C$$