

MATHEMATICS 317 April 2004 Final Exam

- [15] 1) At time $t = 0$, NASA launches a rocket which follows a trajectory so that its position at any time t is

$$x = \frac{4\sqrt{2}}{3}t^{3/2}, \quad y = \frac{4\sqrt{2}}{3}t^{3/2}, \quad z = t(2 - t)$$

- (i) Assuming that the flight ends when $z = 0$, find out how far the rocket travels.
- (ii) Find the unit tangent and unit normal to the trajectory at its highest point.
- (iii) The material of part (iii) was not covered this year.

- [10] 2) The material of this question was not covered this year.

- [10] 3) True or false (reasons must be given):

- (i) If a smooth vector field on \mathbb{R}^3 is curl free and divergence free, then its potential is harmonic.
- (ii) If \vec{F} is a smooth conservative vector field on \mathbb{R}^3 , then its flux through any smooth closed surface is zero.

- [15] 4) A physicist studies a vector field \vec{F} in her lab. She knows from theoretical considerations that \vec{F} must be of the form $\vec{F} = \nabla \times \vec{G}$, for some smooth vector field \vec{G} . Experiments also show that \vec{F} must be of the form

$$\vec{F}(x, y, z) = (xz + xy)\hat{i} + \alpha(yz - xy)\hat{j} + \beta(yz + xz)\hat{k}$$

where α and β are constant.

- (i) Determine α and β .
- (ii) Further experiments show that $\vec{G} = xyz\hat{i} - xyz\hat{j} + g(x, y, z)\hat{k}$. Find the unknown function $g(x, y, z)$.

- [10] 5) Recall that if S is a smooth closed surface with outer normal field \hat{n} , then for any smooth function $p(x, y, z)$ on \mathbb{R}^3 , we have

$$\iint_S p\hat{n} \, dS = \iiint_E \nabla p \, dV$$

where E is the solid bounded by S . Show that as a consequence, the total force exerted on the surface of a solid body contained in a gas of constant pressure is zero. (Recall that the pressure acts in the direction normal to the surface.)

- [15] 6) Use Green's theorem to establish that if C is a simple closed curve in the plane, then the area A enclosed by C is given by

$$A = \frac{1}{2} \oint_C x \, dy - y \, dx$$

Use this to calculate the area inside the curve $x^{2/3} + y^{2/3} = 1$. Hint: note the trig identities $\sin 2\alpha = 2\sin\alpha \cos\alpha$ and $2\sin^2\alpha = 1 - \cos 2\alpha$.]

- [15] 7) Consider the vector field $\vec{F}(x, y, z) = -2xy\hat{i} + (y^2 + \sin(xz))\hat{j} + (x^2 + y^2)\hat{k}$.

- (i) Calculate $\nabla \cdot \vec{F}$.
- (ii) Find the flux of \vec{F} through the surface S defined by

$$x^2 + y^2 + (z - 12)^2 = 13^2, \quad z \geq 0$$

using the outward normal to S .

- [10] 8) Let C be the curve given by the parametric equations:

$$x = \cos t, \quad y = \sqrt{2} \sin t, \quad z = \cos t, \quad 0 \leq t \leq 2\pi$$

and let

$$\vec{F} = z\hat{i} + x\hat{j} + y^3z^3\hat{k}$$

Use Stokes' theorem to evaluate

$$\oint \vec{F} \cdot d\vec{r}$$