

Euler Angles

Euler angles are three angles that provide coordinates on $SO(3)$. Denote by $R_1(\theta)$ and $R_3(\theta)$ rotations about the x - and z - axes, respectively, by an angle θ . That is

$$R_1(\theta) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix} \quad R_3(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

For each $(\varphi, \theta, \psi) \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$, the matrix $R_3(\varphi)R_1(\theta)R_3(\psi)$ is certainly in $SO(3)$. The following theorem says that each $R \in SO(3)$ has a representation of the form $R_3(\varphi)R_1(\theta)R_3(\psi)$ with $(\varphi, \theta, \psi) \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$ and that this representation is unique unless θ is 0 or π (in which case $R_3(\varphi)R_1(\theta)R_3(\psi)$ depends only on $\varphi + \psi$ or $\varphi - \psi$, respectively).

Theorem.

(a) (Existence) For each $R \in SO(3)$, there an $(\varphi, \theta, \psi) \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$ such that $R = R_3(\varphi)R_1(\theta)R_3(\psi)$.

(b) (Uniqueness) Let $(\varphi, \theta, \psi), (\varphi', \theta', \psi') \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$ and assume that $R_3(\varphi)R_1(\theta)R_3(\psi) = R_3(\varphi')R_1(\theta')R_3(\psi')$. Then $\theta = \theta'$. If $\theta \neq 0, \pi$, then $\varphi = \varphi'$ and $\psi = \psi'$.


(c) If $R = [\hat{i}_1 \ \hat{i}_2 \ \hat{i}_3]$, then

- θ is the angle between \hat{i}_3 and the z -axis and
- if $\theta \notin \{0, \pi\}$, then φ is the angle between the projection of \hat{i}_3 on the xy -plane and the x -axis, plus $\frac{\pi}{2}$

Proof: (a) Write $R = [\hat{i}_1 \ \hat{i}_2 \ \hat{i}_3]$ where \hat{i}_1, \hat{i}_2 and \hat{i}_3 are three mutually perpendicular unit vectors that form a right handed triple and use

$$\hat{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad \hat{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \hat{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

to denote the standard basis for \mathbb{R}^3 .




Set

$$\hat{n} = \begin{cases} \frac{\hat{e}_3 \times \hat{i}_3}{|\hat{e}_3 \times \hat{i}_3|} & \text{if } \hat{i}_3 \neq \pm \hat{e}_3 \\ \hat{i}_1 & \text{if } \hat{i}_3 = \hat{e}_3 \\ -\hat{i}_1 & \text{if } \hat{i}_3 = -\hat{e}_3 \end{cases}$$


Since \hat{e}_1 and \hat{n} are both unit vectors that are perpendicular to \hat{e}_3 , there is an angle $\varphi \in [0, 2\pi)$ such that rotation about \hat{e}_3 by φ maps \hat{e}_1 to \hat{n} . This rotation leaves \hat{e}_3 invariant. Thus

$$\hat{n} = R_3(\varphi)\hat{e}_1 \quad \hat{e}_3 = R_3(\varphi)\hat{e}_3$$



Now \hat{e}_3 and \hat{i}_3 are both unit vectors that are perpendicular to \hat{n} . So there is an angle $\theta \in [0, 2\pi)$ such that rotation about \hat{n} by θ maps \hat{e}_3 to \hat{i}_3 . In fact $\theta \in [0, \pi]$ because⁽¹⁾ \hat{n} has direction $\hat{e}_3 \times \hat{i}_3$. The rotation about \hat{n} by θ is implemented by $R_3(\varphi)R_1(\theta)R_3(-\varphi)$ (since $\frac{d}{d\theta}R_3(\varphi)R_1(\theta)R_3(-\varphi)\vec{x} = R_3(\varphi)(\hat{e}_1 \times (R_3(-\varphi)\vec{x})) = (R_3(\varphi)\hat{e}_1) \times \vec{x} = \hat{n} \times \vec{x}$). So


$$\begin{aligned} \hat{i}_3 &= R_3(\varphi)R_1(\theta)R_3(-\varphi)\hat{e}_3 = R_3(\varphi)R_1(\theta)\hat{e}_3 \\ \hat{n} &= R_3(\varphi)R_1(\theta)R_3(-\varphi)\hat{n} = R_3(\varphi)R_1(\theta)\hat{e}_1 \end{aligned}$$



Finally, \hat{i}_1 and \hat{n} are both unit vectors that are perpendicular to \hat{i}_3 . So there is an angle $\psi \in [0, 2\pi)$ such that rotation about \hat{i}_3 by ψ maps \hat{n} to \hat{i}_1 . The rotation about \hat{i}_3 by ψ is implemented by $R_3(\varphi)R_1(\theta)R_3(\psi)R_1(-\theta)R_3(-\varphi)$. (Note that $R_3(\varphi)R_1(\theta)R_3(\psi)R_1(-\theta)R_3(-\varphi)\hat{i}_3 = \hat{i}_3$.) So

⁽¹⁾ Use a coordinate system with \hat{e}_3 on the positive z -axis and \hat{n} on the positive y -axis. If $\hat{i}_3 = a\hat{e}_1 + b\hat{e}_2 + c\hat{e}_3$, then $\hat{e}_3 \times \hat{i}_3 = a\hat{e}_2 - b\hat{e}_1$. For this to be a positive multiple of \hat{e}_2 , we need $b = 0$ and $a > 0$. So \hat{i}_3 must be in the xz -plane with positive x -component.

$$\begin{aligned}\hat{i}_1 &= R_3(\varphi)R_1(\theta)R_3(\psi)R_1(-\theta)R_3(-\varphi)\hat{n} = R_3(\varphi)R_1(\theta)R_3(\psi)\hat{e}_1 \\ \hat{i}_3 &= R_3(\varphi)R_1(\theta)R_3(\psi)R_1(-\theta)R_3(-\varphi)\hat{i}_3 = R_3(\varphi)R_1(\theta)R_3(\psi)\hat{e}_3\end{aligned}$$



In other words, the two matrices R and $R_3(\varphi)R_1(\theta)R_3(\psi)$ have the same first column (namely \hat{i}_1) and the same third column (namely \hat{i}_3). The columns of any matrix in $SO(3)$ form a right handed triple of mutually perpendicular unit vectors. As both R and $R_3(\varphi)R_1(\theta)R_3(\psi)$ are in $SO(3)$, they both must have the same middle column too (namely \hat{i}_2).

Uniqueness: Let $(\varphi, \theta, \psi), (\varphi', \theta', \psi') \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$ and assume that

$$R_3(\varphi)R_1(\theta)R_3(\psi) = R_3(\varphi')R_1(\theta')R_3(\psi')$$

Multiplying on the left by $R_3(-\varphi')$ and on the right by $R_3(-\psi')$ gives

$$R_3(\varphi - \varphi')R_1(\theta)R_3(\psi - \psi') = R_1(\theta')$$

In particular $R_3(\varphi - \varphi')R_1(\theta)R_3(\psi - \psi')\hat{e}_3 = R_1(\theta')\hat{e}_3$. As

$$R_1(\theta')\hat{e}_3 = \begin{bmatrix} 0 \\ -\sin \theta' \\ \cos \theta' \end{bmatrix}$$

and


$$\begin{aligned}R_3(\varphi - \varphi')R_1(\theta)R_3(\psi - \psi')\hat{e}_3 &= R_3(\varphi - \varphi')R_1(\theta)\hat{e}_3 \\ &= \begin{bmatrix} \cos(\varphi - \varphi') & -\sin(\varphi - \varphi') & 0 \\ \sin(\varphi - \varphi') & \cos(\varphi - \varphi') & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -\sin \theta \\ \cos \theta \end{bmatrix} \\ &= \begin{bmatrix} \sin(\varphi - \varphi') \sin \theta \\ -\cos(\varphi - \varphi') \sin \theta \\ \cos \theta \end{bmatrix}\end{aligned}$$

we have

$$\begin{bmatrix} 0 \\ -\sin \theta' \\ \cos \theta' \end{bmatrix} = \begin{bmatrix} \sin(\varphi - \varphi') \sin \theta \\ -\cos(\varphi - \varphi') \sin \theta \\ \cos \theta \end{bmatrix}$$

As $0 \leq \theta, \theta' \in [0, \pi]$ and \cos is 1-1 on $[0, \pi]$, we now have that $\theta = \theta'$. As long as $\theta \neq 0, \pi$, we also have $\sin(\varphi - \varphi') = 0$ and $\cos(\varphi - \varphi') = 1$, which forces $\varphi - \varphi'$ to be an integer multiple of 2π . Since $0 \leq \varphi, \varphi' < 2\pi$, this in turn forces $\varphi = \varphi'$. Finally, $R_3(\varphi)R_1(\theta)R_3(\psi) = R_3(\varphi')R_1(\theta')R_3(\psi')$ has now reduced to $R_3(\varphi)R_1(\theta)R_3(\psi) = R_3(\varphi)R_1(\theta)R_3(\psi')$, which is equivalent to $R_3(\psi) = R_3(\psi')$. Since $0 \leq \psi, \psi' < 2\pi$, this forces $\psi = \psi'$.

(c) Assume that $\theta \neq 0$. Then $\hat{n} = \frac{\hat{e}_3 \times \hat{i}_3}{|\hat{e}_3 \times \hat{i}_3|}$ is perpendicular to both \hat{i}_3 and \hat{e}_3 . We defined θ to be the angle of rotation about \hat{n} that maps \hat{e}_3 to \hat{i}_3 . By definition, this is the angle between \hat{e}_3 and \hat{i}_3 . Let P_{xy} denote the xy -plane and P_3 denote the plane containing \hat{e}_3 and \hat{i}_3 . Then \hat{n} lies in P_{xy} (because it is perpendicular to \hat{e}_3) and is perpendicular to P_3 . The projection of \hat{i}_3 onto the xy -plane (denoted $\vec{\pi}_3$ in the figure below) lies in both P_3 and P_{xy} and so is an angle $\frac{\pi}{2}$ from \hat{n} . We defined φ to be the angle between \hat{e}_1 and \hat{n} .



■