


# Derivation of the Wave Equation

In these notes we apply Newton's law to an elastic string, concluding that small amplitude transverse vibrations of the string obey the wave equation. Consider a tiny element of the string.



The basic notation is

$u(x, t)$  = vertical displacement of the string from the  $x$  axis at position  $x$  and time  $t$

$\theta(x, t)$  = angle between the string and a horizontal line at position  $x$  and time  $t$

$T(x, t)$  = tension in the string at position  $x$  and time  $t$

$\rho(x)$  = mass density of the string at position  $x$

The forces acting on the tiny element of string are

- (a) tension pulling to the right, which has magnitude  $T(x + \Delta x, t)$  and acts at an angle  $\theta(x + \Delta x, t)$  above horizontal
- (b) tension pulling to the left, which has magnitude  $T(x, t)$  and acts at an angle  $\theta(x, t)$  below horizontal and, possibly,
- (c) various external forces, like gravity. We shall assume that all of the external forces act vertically and we shall denote by  $F(x, t)\Delta x$  the net magnitude of the external force acting on the element of string.

The mass of the element of string is essentially  $\rho(x)\sqrt{\Delta x^2 + \Delta u^2}$  so the vertical component of Newton's law says that

$$\rho(x)\sqrt{\Delta x^2 + \Delta u^2} \frac{\partial^2 u}{\partial t^2}(x, t) = T(x + \Delta x, t) \sin \theta(x + \Delta x, t) - T(x, t) \sin \theta(x, t) + F(x, t)\Delta x$$


Dividing by  $\Delta x$  and taking the limit as  $\Delta x \rightarrow 0$  gives

$$\begin{aligned} \rho(x)\sqrt{1 + \left(\frac{\partial u}{\partial x}\right)^2} \frac{\partial^2 u}{\partial t^2}(x, t) &= \frac{\partial}{\partial x} [T(x, t) \sin \theta(x, t)] + F(x, t) \\ &= \frac{\partial T}{\partial x}(x, t) \sin \theta(x, t) + T(x, t) \cos \theta(x, t) \frac{\partial \theta}{\partial x}(x, t) + F(x, t) \end{aligned} \tag{1}$$

We can dispose of all the  $\theta$ 's by observing from the figure that

$$\tan \theta(x, t) = \lim_{\Delta x \rightarrow 0} \frac{\Delta u}{\Delta x} = \frac{\partial u}{\partial x}(x, t)$$

which implies, using the figure on the right below, that

$$\begin{aligned} \sin \theta(x, t) &= \frac{\frac{\partial u}{\partial x}(x, t)}{\sqrt{1 + \left(\frac{\partial u}{\partial x}(x, t)\right)^2}} & \cos \theta(x, t) &= \frac{1}{\sqrt{1 + \left(\frac{\partial u}{\partial x}(x, t)\right)^2}} & \begin{array}{c} \sqrt{1 + \tan^2 \theta} \\ \tan \theta \\ 1 \end{array} \\ \theta(x, t) &= \tan^{-1} \frac{\partial u}{\partial x}(x, t) & \frac{\partial \theta}{\partial x}(x, t) &= \frac{\frac{\partial^2 u}{\partial x^2}(x, t)}{1 + \left(\frac{\partial u}{\partial x}(x, t)\right)^2} \end{aligned}$$


Substituting these formulae into (1) give a horrendous mess. However, we can get considerable simplification by looking only at small vibrations. By a small vibration, we mean that  $|\theta(x, t)| \ll 1$  for all  $x$  and  $t$ . This implies that  $|\tan \theta(x, t)| \ll 1$ , hence that  $|\frac{\partial u}{\partial x}(x, t)| \ll 1$  and hence that

$$\sqrt{1 + \left(\frac{\partial u}{\partial x}\right)^2} \approx 1 \quad \sin \theta(x, t) \approx \frac{\partial u}{\partial x}(x, t) \quad \cos \theta(x, t) \approx 1 \quad \frac{\partial \theta}{\partial x}(x, t) \approx \frac{\partial^2 u}{\partial x^2}(x, t) \quad (2)$$

Substituting these into equation (1) give

$$\rho(x) \frac{\partial^2 u}{\partial t^2}(x, t) = \frac{\partial T}{\partial x}(x, t) \frac{\partial u}{\partial x}(x, t) + T(x, t) \frac{\partial^2 u}{\partial x^2}(x, t) + F(x, t) \quad (3)$$

which is indeed relatively simple, but still exhibits a problem. This is one equation in the two unknowns  $u$  and  $T$ .

Fortunately there is a second equation lurking in the background, that we haven't used. Namely, the horizontal component of Newton's law of motion. As a second simplification, we assume that there are only transverse vibrations. Our tiny string element moves only vertically. Then the net horizontal force on it must be zero. That is,

$$T(x + \Delta x, t) \cos \theta(x + \Delta x, t) - T(x, t) \cos \theta(x, t) = 0$$

Dividing by  $\Delta x$  and taking the limit as  $\Delta x$  tends to zero gives

$$\frac{\partial}{\partial x} [T(x, t) \cos \theta(x, t)] = 0$$

For small amplitude vibrations,  $\cos \theta$  is very close to one and  $\frac{\partial T}{\partial x}(x, t)$  is very close to zero. In other words  $T$  is a function of  $t$  only, which is determined by how hard you are pulling on the ends of the string at time  $t$ . So for small, transverse vibrations, (3) simplifies further to

$$\rho(x) \frac{\partial^2 u}{\partial t^2}(x, t) = T(t) \frac{\partial^2 u}{\partial x^2}(x, t) + F(x, t) \quad (4)$$

In the event that the string density  $\rho$  is a constant, independent of  $x$ , the string tension  $T(t)$  is a constant independent of  $t$  (in other words you are not continually playing with the tuning pegs) and there are no external forces  $F$  we end up with

$$\boxed{\frac{\partial^2 u}{\partial t^2}(x, t) = c^2 \frac{\partial^2 u}{\partial x^2}(x, t)}$$

where

$$c = \sqrt{\frac{T}{\rho}}$$