

Orbifold Euler characteristics and the number of commuting  $m$ -tuples in the symmetric groups

By Jim Bryan\* and Jason Fulman

University of California at Berkeley and Dartmouth College

Direct Correspondence to:

Jason Fulman

Dartmouth College

Department of Mathematics

6188 Bradley Hall

Hanover, NH 03755

email:jason.e.fulman@dartmouth.edu

\* Supported in part by a grant from the Ford Foundation.

## Abstract

Generating functions for the number of commuting  $m$ -tuples in the symmetric groups are obtained. We define a natural sequence of “orbifold Euler characteristics” for a finite group  $G$  acting on a manifold  $X$ . Our definition generalizes the ordinary Euler characteristic of  $X/G$  and the string-theoretic orbifold Euler characteristic. Our formulae for commuting  $m$ -tuples underlie formulas that generalize the results of Macdonald and Hirzebruch-Höfer concerning the ordinary and string-theoretic Euler characteristics of symmetric products.

## 1 Introduction

Let  $X$  be a manifold with the action of a finite group  $G$ . The Euler characteristic of the quotient space  $X/G$  can be computed by the Lefschetz fixed point formula:

$$\chi(X/G) = \frac{1}{|G|} \sum_{g \in G} \chi(X^g)$$

where  $X^g$  is the fixed point set of  $g$ . Motivated by string theory, physicists have defined an “orbifold characteristic” by

$$\chi(X, G) = \frac{1}{|G|} \sum_{gh=hg} \chi(X^{(g,h)})$$

where the sum runs over commuting pairs and  $X^{(g,h)}$  denotes the common fixed point set of  $g$  and  $h$ .

We introduce a natural sequence of orbifold Euler characteristics  $\chi_m(X, G)$  for  $m = 1, 2, \dots$  so that  $\chi(X/G)$  and  $\chi(X, G)$  appear as the first two terms. Namely, if we denote by  $Com(G, m)$  the set of mutually commuting  $m$ -tuples  $(g_1, \dots, g_m)$  and by  $X^{(g_1, \dots, g_m)}$  the simultaneous fixed point set, then we define the  $m$ -th orbifold characteristic to be

$$\chi_m(X, G) = \frac{1}{|G|} \sum_{Com(G, m)} \chi(X^{(g_1, \dots, g_m)}). \quad (1)$$

In the case of a symmetric product, *i.e.*  $X$  is the  $n$ -fold product  $M^n$  and  $G$  is the symmetric group  $S_n$ , there are combinatorial formulas for  $\chi_1$  and  $\chi_2$  due to Macdonald [5] and Hirzebruch-Höfer [3] respectively. The main result of this note (Theorem 1) is a generalization of those formulas to  $\chi_m$  for arbitrary  $m$ . In the case where  $M$  has (ordinary) Euler characteristic 1, our formulas specialize to generating functions for  $|Com(S_n, m)|$ , the number of commuting  $m$ -tuples in  $S_n$ .

Finally, we remark that the first two terms in our sequence  $\chi_m(X, G)$  of orbifold Euler characteristics are the Euler characteristics of the cohomology theories  $H_G^*(X; \mathbf{Q})$  and  $K_G^*(X; \mathbf{Q})$  respectively. This was observed by Segal, [1] who was led to speculate that the heirarchy of generalized cohomology theories investigated by Hopkins and Kuhn [4] may have something to do with the sequence of Euler characteristics defined in this paper (our definition is implicitly suggested in [1]). We hope that our combinatorial formulas will provide clues to the nature of these theories.

## 2 Formulae

In this section we specialize to the case of symmetric products so that  $X = M^n$  and  $G = S_n$ . For  $(\pi_1, \dots, \pi_m) \in Com(S_n, m)$ , let  $\#(\pi_1, \dots, \pi_m)$  be the number of connected components in the graph on vertex set  $\{1, \dots, n\}$  defined by connecting the vertices according to the permutations  $\pi_1, \dots, \pi_m$ . For instance,  $\#(\pi_1)$  is the number of cycles of  $\pi$ . The main result of this note is the following theorem.

**Theorem 1** *Let  $\chi$  denote the (ordinary) Euler characteristic of  $M$ . The generating function for the orbifold Euler characteristic  $\chi_m(M^n, S_n)$  satisfies the following formulas:*

$$\sum_{n=0}^{\infty} \chi_m(M^n, S_n) u^n = \sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{\pi_1, \dots, \pi_m \in Com(S_n, m)} \chi^{\#(\pi_1, \dots, \pi_m)} \quad (2)$$

$$= \left( \sum_{n=0}^{\infty} |Com(S_n, m)| \frac{u^n}{n!} \right)^\chi \quad (3)$$

$$= \prod_{i_1, \dots, i_{m-1}=1}^{\infty} (1 - u^{i_1 \dots i_{m-1}})^{-\chi i_1^{m-2} i_2^{m-3} \dots i_{m-2}}. \quad (4)$$

**Remarks:** We will show that Equation 2 follows directly from the definitions and a straightforward geometric argument. Equation 3 is proved in Lemma 1 and shows that it suffices to prove Equation 4 in the case  $\chi = 1$ . Our main result then should be regarded as Equation 4 which in light of Equation 3 gives a generating function for the number of commuting  $m$ -tuples in  $S_n$ . Note also that for  $m = 1$  Equation 4 is Macdonald's formula  $(1 - u)^{-\chi}$  for the Euler characteristic of a symmetric product and for  $m = 2$  Equation 4 is Hirzebruch and Höfer's formula for the string-theoretic orbifold Euler characteristic of a symmetric product.

To prove Equation 2 it suffices to see that

$$\chi(M^{(\pi_1, \dots, \pi_m)}) = (\chi(M))^{\#(\pi_1, \dots, \pi_m)}.$$

Partition  $\{1, \dots, n\}$  into disjoint subsets  $I_1, \dots, I_{\#(\pi_1, \dots, \pi_m)}$  according to the components of the graph associated to  $(\pi_1, \dots, \pi_m)$ . Then the small diagonal in the product  $\prod_{i \in I_j} M_i$  is fixed by  $(\pi_1, \dots, \pi_m)$  and is homeomorphic to  $M$ . The full fixed set of  $(\pi_1, \dots, \pi_m)$  is then the product of all the small diagonals in the subproducts associated to the  $I_j$ 's. By the multiplicative properties of Euler characteristic we see that  $\chi(M^{(\pi_1, \dots, \pi_m)}) = (\chi(M))^{\#(\pi_1, \dots, \pi_m)}$ .

**Lemma 1** *For  $\chi$  a natural number,*

$$\sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{\pi_1, \dots, \pi_m \in \text{Com}(S_n, m)} \chi^{\#(\pi_1, \dots, \pi_m)} = \left( \sum_{n=0}^{\infty} \frac{u^n |\text{Com}(S_n, m)|}{n!} \right)^{\chi}$$

PROOF: It suffices to show that an ordered  $m$ -tuple  $(\pi_1, \dots, \pi_m)$  of mutually commuting elements of  $S_n$  contributes equally to the coefficient of  $\frac{u^n}{n!}$  on both sides of the equation. The contribution to this coefficient on the left-hand side is  $\chi^{\#(\pi_1, \dots, \pi_m)}$ .

The right hand side can be rewritten as

$$\sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{n_1, \dots, n_\chi: \sum n_i = n} \binom{n}{n_1, \dots, n_\chi} |Com(S_{n_1}, m)| \cdots |Com(S_{n_\chi}, m)|.$$

Observe that  $\binom{n}{n_1, \dots, n_\chi} |Com(S_{n_1}, m)| \cdots |Com(S_{n_\chi}, m)|$  is the number of ways of decomposing the vertex set  $\{1, \dots, n\}$  into  $\chi$  ordered subsets  $S_1, \dots, S_\chi$  of sizes  $n_1, \dots, n_\chi$  and defining an ordered  $m$ -tuple of mutually commuting elements of  $S_{n_i}$  on each subset. Gluing these together defines an ordered  $m$ -tuple of mutually commuting elements of  $S_n$ . Note that the  $m$ -tuple  $(\pi_1, \dots, \pi_m)$  arises in  $\chi^{\#(\pi_1, \dots, \pi_m)}$  ways, because each of the  $\#(\pi_1, \dots, \pi_m)$  connected components of the graph corresponding to  $(\pi_1, \dots, \pi_m)$  could have come from any of the  $\chi$  subsets  $S_1, \dots, S_\chi$ .  $\square$

Let us now recall some facts about wreath products of groups. All of this can be found in Sections 4.1 and 4.2 of James and Kerber [2]. Given a group  $G$ , the wreath product  $GWrS_n$  is defined as a set by  $(g_1, \dots, g_n; \pi)$  where  $g_i \in G$  and  $\pi \in S_n$ . Letting permutations act on the right, the group multiplication is defined by:

$$(g_1, \dots, g_n; \pi)(h_1, \dots, h_n; \tau) = (g_1 h_{(1)\pi^{-1}}, \dots, g_n h_{(n)\pi^{-1}}; \pi\tau)$$

Furthermore, the conjugacy classes of  $GWrS_n$  are parameterized as follows. Let  $Cl_1, \dots, Cl_i$  be the conjugacy classes of  $G$ . Then the conjugacy classes of  $GWrS_n$  correspond to arrays  $(M_{j,k})$  satisfying the properties:

1.  $M_{j,k} = 0$  if  $j > i$
2.  $\sum_{j,k} k M_{j,k} = n$

The correspondence can be made explicit. For  $(g_1, \dots, g_n; \pi) \in GWrS_n$ , let  $M_{j,k}$  be the number of  $k$ -cycles of  $\pi$  such that multiplying the  $k$   $g_i$  whose subscripts lies in the  $k$ -cycle gives an element of  $G$  belonging to the conjugacy class  $Cl_j$  of  $G$ . The matrix so-defined clearly satisfies the above two conditions.

Lemma 2 is a key ingredient of this paper. It says that centralizers of elements of wreath products can be expressed in terms of wreath products; this will lead to an inductive proof of Theorem 1.

**Lemma 2** *Let  $C_i$  denote a cyclic group of order  $i$ . Then the centralizer in  $C_iWrS_n$  of an element in the conjugacy class corresponding to the data  $M_{j,k}$  is isomorphic to the direct product*

$$\prod_{j,k} C_{i_k}WrS_{M_{j,k}}$$

PROOF: To start, let us construct an element  $(g_1, \dots, g_n; \pi)$  of  $C_iWrS_n$  with conjugacy class data  $M_{j,k}$ . This can be done as follows:

1. Pick  $\pi$  to be any permutation with  $\sum_j M_{j,k}$   $k$ -cycles
2. For each  $j$  choose  $M_{j,k}$  of the  $k$ -cycles of  $\pi$  and think of them as  $k$ -cycles of type  $j$
3. Assign (in any order) to the  $g_i$  whose subscripts are contained in a  $k$ -cycle of type  $j$  of  $\pi$  the values  $(c_j, 1, \dots, 1)$  where  $c_j$  is an element in the  $j$ th conjugacy class of the group  $C_i$

To describe the centralizer of this element  $(g_1, \dots, g_n; \pi)$ , note that conjugation in  $GW_rS_n$  works as

$$\begin{aligned} & (h_1, \dots, h_n; \tau)(g_1, \dots, g_n; \pi)(h_{(1)\tau}^{-1}, \dots, h_{(n)\tau}^{-1}; \tau^{-1}) \\ &= (h_1 g_{(1)\tau^{-1}} h_{(1)\tau\pi^{-1}\tau^{-1}}^{-1}, \dots; \tau\pi\tau^{-1}) \end{aligned}$$

It is easy to see that if  $(h_1, \dots, h_n; \tau)$  commutes with  $(g_1, \dots, g_n; \pi)$ , then  $\tau$  operates on the  $M_{j,k}$   $k$ -cycles of  $\pi$  of type  $j$  by first permuting these cycles amongst themselves and then performing some power of a cyclic shift within each cycle. Further, among the  $h_i$  whose subscripts lie in a

$k$ -cycle of  $\pi$  of type  $j$  exactly one can be chosen arbitrarily in  $C_i$ —the other  $h$ 's with subscripts in that  $k$ -cycle then have determined values.

The direct product assertion of the theorem is then easily checked; the only non-trivial part is to see the copy of  $C_{ik}WrS_{M_{j,k}}$ . Here the  $S_{M_{j,k}}$  permutes the  $M_{j,k}$   $k$ -cycles of type  $j$ , and the generator of the  $C_{ik}$  corresponds to having  $\tau$  cyclically permuting within the  $k$  cycle and having the  $h$ 's with subscripts in that  $k$ -cycle equal to  $\{c_j, 1, \dots, 1\}$ , where  $c_j$  is a generator of  $C_i$ .  $\square$

With these preliminaries in hand, induction can be used to prove the following result. Note that by Lemma 1, only the  $i = 1$  case of Theorem 2 is needed to prove the main result of this paper, Theorem 1. However, the stronger statement (general  $i$ ) in Theorem 2 makes the induction work by making the induction hypothesis stronger.

**Theorem 2** For  $m \geq 2$ ,

$$\sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, m)|}{|C_i Wr S_n|} = \prod_{i_1, \dots, i_{m-1}=1}^{\infty} \left( \frac{1}{1 - u^{i_1 \dots i_{m-1}}} \right)^{i_1^{m-1} i_1^{m-2} i_2^{m-3} \dots i_{m-2}}$$

PROOF: The proof proceeds by induction on  $m$ . We use the notation that if  $\lambda$  denotes a conjugacy class of a group  $G$ , then  $C_G(\lambda)$  is the centralizer in  $G$  of some element of  $\lambda$  (hence  $C_G(\lambda)$  is well defined up to isomorphism). For the base case  $m = 2$  observe that

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, 2)|}{|C_i Wr S_n|} \\ &= \sum_{n=0}^{\infty} \frac{u^n}{|C_i Wr S_n|} \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} \frac{|C_i Wr S_n|}{|C_{C_i Wr S_n}(M_{j,k})|} |C_{C_i Wr S_n}(M_{j,k})| \\ &= \sum_{n=0}^{\infty} u^n \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} 1 \\ &= \prod_{i_1=1}^{\infty} \left( \frac{1}{1 - u^{i_1}} \right)^{i_1} \end{aligned}$$

For the induction step, the parameterization of conjugacy classes of wreath products and Lemma

2 imply that

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, m)|}{|C_i Wr S_n|} \\
= & \sum_{n=0}^{\infty} \frac{u^n}{|C_i Wr S_n|} \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} \frac{|C_i Wr S_n|}{|C_{C_i Wr S_n}(M_{j,k})|} |Com(C_{C_i Wr S_n}(M_{j,k}), m-1)| \\
= & \left[ \prod_{k=1}^{\infty} \sum_{a=0}^{\infty} \frac{u^{ka} |Com(C_{ik} Wr S_a, m-1)|}{|C_{ik} Wr S_a|} \right]^i \\
= & \left[ \prod_{k=1}^{\infty} \prod_{i_2, \dots, i_{m-1}=1}^{\infty} \left( \frac{1}{1 - u^{ki_2 \dots i_{m-1}}} \right)^{(ik)^{m-2} i_2^{m-3} \dots i_{m-2}} \right]^i \\
= & \prod_{i_1, \dots, i_{m-1}=1}^{\infty} \left( \frac{1}{1 - u^{i_1 \dots i_{m-1}}} \right)^{i_1^{m-1} i_2^{m-2} i_3^{m-3} \dots i_{m-2}}
\end{aligned}$$

□

## References

- [1] Atiyah, M. and Segal, G., On equivariant Euler characteristics. J. Geom. Phys. **6**, no. 4, 671-677 (1989).
- [2] James, G. and Kerber, A., The representation theory of the symmetric group. Encyclopedia of Mathematics and its Applications. Volume 16, (1981).
- [3] Hirzebruch, F. and Höfer, T., On the Euler number of an orbifold. Mathematische Annalen **286**, 255–260 (1990).
- [4] Hopkins, M., Kuhn, N., and Ravenel, D., Morava  $K$ -theories of classifying spaces and generalized characters for finite groups. Algebraic topology (San Feliu de Guixols), Lecture Notes in Math. **1509**, 186-209 (1990).

- [5] Macdonald, I.G., The Poincaré polynomial of a symmetric product. Proc. Camb. Phil. Soc.  
58, 563-568 (1962).

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF CALIFORNIA

BERKELEY, CA 94720

DEPARTMENT OF MATHEMATICS

DARTMOUTH COLLEGE

HANOVER, NH 03755

`jbryan@math.berkeley.edu`

`jason.e.fulman@dartmouth.edu`