

Orbifold Euler characteristics and the number of commuting m -tuples in the symmetric groups

By Jim Bryan* and Jason Fulman

University of California at Berkeley and Dartmouth College

Direct Correspondence to:

Jason Fulman

Dartmouth College

Department of Mathematics

6188 Bradley Hall

Hanover, NH 03755

email:jason.e.fulman@dartmouth.edu

* Supported in part by a grant from the Ford Foundation.

Abstract

Generating functions for the number of commuting m -tuples in the symmetric groups are obtained. We define a natural sequence of “orbifold Euler characteristics” for a finite group G acting on a manifold X . Our definition generalizes the ordinary Euler characteristic of X/G and the string-theoretic orbifold Euler characteristic. Our formulae for commuting m -tuples underlie formulas that generalize the results of Macdonald and Hirzebruch-Höfer concerning the ordinary and string-theoretic Euler characteristics of symmetric products.

1 Introduction

Let X be a manifold with the action of a finite group G . The Euler characteristic of the quotient space X/G can be computed by the Lefschetz fixed point formula:

$$\chi(X/G) = \frac{1}{|G|} \sum_{g \in G} \chi(X^g)$$

where X^g is the fixed point set of g . Motivated by string theory, physicists have defined an “orbifold characteristic” by

$$\chi(X, G) = \frac{1}{|G|} \sum_{gh=hg} \chi(X^{(g,h)})$$

where the sum runs over commuting pairs and $X^{(g,h)}$ denotes the common fixed point set of g and h .

We introduce a natural sequence of orbifold Euler characteristics $\chi_m(X, G)$ for $m = 1, 2, \dots$ so that $\chi(X/G)$ and $\chi(X, G)$ appear as the first two terms. Namely, if we denote by $Com(G, m)$ the set of mutually commuting m -tuples (g_1, \dots, g_m) and by $X^{(g_1, \dots, g_m)}$ the simultaneous fixed point set, then we define the m -th orbifold characteristic to be

$$\chi_m(X, G) = \frac{1}{|G|} \sum_{Com(G, m)} \chi(X^{(g_1, \dots, g_m)}). \quad (1)$$

In the case of a symmetric product, *i.e.* X is the n -fold product M^n and G is the symmetric group S_n , there are combinatorial formulas for χ_1 and χ_2 due to Macdonald [5] and Hirzebruch-Höfer [3] respectively. The main result of this note (Theorem 1) is a generalization of those formulas to χ_m for arbitrary m . In the case where M has (ordinary) Euler characteristic 1, our formulas specialize to generating functions for $|Com(S_n, m)|$, the number of commuting m -tuples in S_n .

Finally, we remark that the first two terms in our sequence $\chi_m(X, G)$ of orbifold Euler characteristics are the Euler characteristics of the cohomology theories $H_G^*(X; \mathbf{Q})$ and $K_G^*(X; \mathbf{Q})$ respectively. This was observed by Segal, [1] who was led to speculate that the heirarchy of generalized cohomology theories investigated by Hopkins and Kuhn [4] may have something to do with the sequence of Euler characteristics defined in this paper (our definition is implicitly suggested in [1]). We hope that our combinatorial formulas will provide clues to the nature of these theories.

2 Formulae

In this section we specialize to the case of symmetric products so that $X = M^n$ and $G = S_n$. For $(\pi_1, \dots, \pi_m) \in Com(S_n, m)$, let $\#(\pi_1, \dots, \pi_m)$ be the number of connected components in the graph on vertex set $\{1, \dots, n\}$ defined by connecting the vertices according to the permutations π_1, \dots, π_m . For instance, $\#(\pi_1)$ is the number of cycles of π . The main result of this note is the following theorem.

Theorem 1 *Let χ denote the (ordinary) Euler characteristic of M . The generating function for the orbifold Euler characteristic $\chi_m(M^n, S_n)$ satisfies the following formulas:*

$$\sum_{n=0}^{\infty} \chi_m(M^n, S_n) u^n = \sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{\pi_1, \dots, \pi_m \in Com(S_n, m)} \chi^{\#(\pi_1, \dots, \pi_m)} \quad (2)$$

$$= \left(\sum_{n=0}^{\infty} |Com(S_n, m)| \frac{u^n}{n!} \right)^\chi \quad (3)$$

$$= \prod_{i_1, \dots, i_{m-1}=1}^{\infty} (1 - u^{i_1 \dots i_{m-1}})^{-\chi i_1^{m-2} i_2^{m-3} \dots i_{m-2}}. \quad (4)$$

Remarks: We will show that Equation 2 follows directly from the definitions and a straightforward geometric argument. Equation 3 is proved in Lemma 1 and shows that it suffices to prove Equation 4 in the case $\chi = 1$. Our main result then should be regarded as Equation 4 which in light of Equation 3 gives a generating function for the number of commuting m -tuples in S_n . Note also that for $m = 1$ Equation 4 is Macdonald's formula $(1 - u)^{-\chi}$ for the Euler characteristic of a symmetric product and for $m = 2$ Equation 4 is Hirzebruch and Höfer's formula for the string-theoretic orbifold Euler characteristic of a symmetric product.

To prove Equation 2 it suffices to see that

$$\chi(M^{(\pi_1, \dots, \pi_m)}) = (\chi(M))^{\#(\pi_1, \dots, \pi_m)}.$$

Partition $\{1, \dots, n\}$ into disjoint subsets $I_1, \dots, I_{\#(\pi_1, \dots, \pi_m)}$ according to the components of the graph associated to (π_1, \dots, π_m) . Then the small diagonal in the product $\prod_{i \in I_j} M_i$ is fixed by (π_1, \dots, π_m) and is homeomorphic to M . The full fixed set of (π_1, \dots, π_m) is then the product of all the small diagonals in the subproducts associated to the I_j 's. By the multiplicative properties of Euler characteristic we see that $\chi(M^{(\pi_1, \dots, \pi_m)}) = (\chi(M))^{\#(\pi_1, \dots, \pi_m)}$.

Lemma 1 For χ a natural number,

$$\sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{\pi_1, \dots, \pi_m \in \text{Com}(S_n, m)} \chi^{\#(\pi_1, \dots, \pi_m)} = \left(\sum_{n=0}^{\infty} \frac{u^n |\text{Com}(S_n, m)|}{n!} \right)^{\chi}$$

PROOF: It suffices to show that an ordered m -tuple (π_1, \dots, π_m) of mutually commuting elements of S_n contributes equally to the coefficient of $\frac{u^n}{n!}$ on both sides of the equation. The contribution to this coefficient on the left-hand side is $\chi^{\#(\pi_1, \dots, \pi_m)}$.

The right hand side can be rewritten as

$$\sum_{n=0}^{\infty} \frac{u^n}{n!} \sum_{n_1, \dots, n_\chi: \sum n_i = n} \binom{n}{n_1, \dots, n_\chi} |Com(S_{n_1}, m)| \cdots |Com(S_{n_\chi}, m)|.$$

Observe that $\binom{n}{n_1, \dots, n_\chi} |Com(S_{n_1}, m)| \cdots |Com(S_{n_\chi}, m)|$ is the number of ways of decomposing the vertex set $\{1, \dots, n\}$ into χ ordered subsets S_1, \dots, S_χ of sizes n_1, \dots, n_χ and defining an ordered m -tuple of mutually commuting elements of S_{n_i} on each subset. Gluing these together defines an ordered m -tuple of mutually commuting elements of S_n . Note that the m -tuple (π_1, \dots, π_m) arises in $\chi^{\#(\pi_1, \dots, \pi_m)}$ ways, because each of the $\#(\pi_1, \dots, \pi_m)$ connected components of the graph corresponding to (π_1, \dots, π_m) could have come from any of the χ subsets S_1, \dots, S_χ . \square

Let us now recall some facts about wreath products of groups. All of this can be found in Sections 4.1 and 4.2 of James and Kerber [2]. Given a group G , the wreath product $GWrS_n$ is defined as a set by $(g_1, \dots, g_n; \pi)$ where $g_i \in G$ and $\pi \in S_n$. Letting permutations act on the right, the group multiplication is defined by:

$$(g_1, \dots, g_n; \pi)(h_1, \dots, h_n; \tau) = (g_1 h_{(1)\pi^{-1}}, \dots, g_n h_{(n)\pi^{-1}}; \pi\tau)$$

Furthermore, the conjugacy classes of $GWrS_n$ are parameterized as follows. Let Cl_1, \dots, Cl_i be the conjugacy classes of G . Then the conjugacy classes of $GWrS_n$ correspond to arrays $(M_{j,k})$ satisfying the properties:

1. $M_{j,k} = 0$ if $j > i$
2. $\sum_{j,k} k M_{j,k} = n$

The correspondence can be made explicit. For $(g_1, \dots, g_n; \pi) \in GWrS_n$, let $M_{j,k}$ be the number of k -cycles of π such that multiplying the k g_i whose subscripts lies in the k -cycle gives an element of G belonging to the conjugacy class Cl_j of G . The matrix so-defined clearly satisfies the above two conditions.

Lemma 2 is a key ingredient of this paper. It says that centralizers of elements of wreath products can be expressed in terms of wreath products; this will lead to an inductive proof of Theorem 1.

Lemma 2 *Let C_i denote a cyclic group of order i . Then the centralizer in C_iWrS_n of an element in the conjugacy class corresponding to the data $M_{j,k}$ is isomorphic to the direct product*

$$\prod_{j,k} C_{i_k}WrS_{M_{j,k}}$$

PROOF: To start, let us construct an element $(g_1, \dots, g_n; \pi)$ of C_iWrS_n with conjugacy class data $M_{j,k}$. This can be done as follows:

1. Pick π to be any permutation with $\sum_j M_{j,k}$ k -cycles
2. For each j choose $M_{j,k}$ of the k -cycles of π and think of them as k -cycles of type j
3. Assign (in any order) to the g_i whose subscripts are contained in a k -cycle of type j of π the values $(c_j, 1, \dots, 1)$ where c_j is an element in the j th conjugacy class of the group C_i

To describe the centralizer of this element $(g_1, \dots, g_n; \pi)$, note that conjugation in GW_rS_n works as

$$\begin{aligned} & (h_1, \dots, h_n; \tau)(g_1, \dots, g_n; \pi)(h_{(1)\tau}^{-1}, \dots, h_{(n)\tau}^{-1}; \tau^{-1}) \\ &= (h_1 g_{(1)\tau^{-1}} h_{(1)\tau\pi^{-1}\tau^{-1}}^{-1}, \dots; \tau\pi\tau^{-1}) \end{aligned}$$

It is easy to see that if $(h_1, \dots, h_n; \tau)$ commutes with $(g_1, \dots, g_n; \pi)$, then τ operates on the $M_{j,k}$ k -cycles of π of type j by first permuting these cycles amongst themselves and then performing some power of a cyclic shift within each cycle. Further, among the h_i whose subscripts lie in a

k -cycle of π of type j exactly one can be chosen arbitrarily in C_i —the other h 's with subscripts in that k -cycle then have determined values.

The direct product assertion of the theorem is then easily checked; the only non-trivial part is to see the copy of $C_{ik}WrS_{M_{j,k}}$. Here the $S_{M_{j,k}}$ permutes the $M_{j,k}$ k -cycles of type j , and the generator of the C_{ik} corresponds to having τ cyclically permuting within the k cycle and having the h 's with subscripts in that k -cycle equal to $\{c_j, 1, \dots, 1\}$, where c_j is a generator of C_i . \square

With these preliminaries in hand, induction can be used to prove the following result. Note that by Lemma 1, only the $i = 1$ case of Theorem 2 is needed to prove the main result of this paper, Theorem 1. However, the stronger statement (general i) in Theorem 2 makes the induction work by making the induction hypothesis stronger.

Theorem 2 For $m \geq 2$,

$$\sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, m)|}{|C_i Wr S_n|} = \prod_{i_1, \dots, i_{m-1}=1}^{\infty} \left(\frac{1}{1 - u^{i_1 \dots i_{m-1}}} \right)^{i_1^{m-1} i_1^{m-2} i_2^{m-3} \dots i_{m-2}}$$

PROOF: The proof proceeds by induction on m . We use the notation that if λ denotes a conjugacy class of a group G , then $C_G(\lambda)$ is the centralizer in G of some element of λ (hence $C_G(\lambda)$ is well defined up to isomorphism). For the base case $m = 2$ observe that

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, 2)|}{|C_i Wr S_n|} \\ &= \sum_{n=0}^{\infty} \frac{u^n}{|C_i Wr S_n|} \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} \frac{|C_i Wr S_n|}{|C_{C_i Wr S_n}(M_{j,k})|} |C_{C_i Wr S_n}(M_{j,k})| \\ &= \sum_{n=0}^{\infty} u^n \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} 1 \\ &= \prod_{i_1=1}^{\infty} \left(\frac{1}{1 - u^{i_1}} \right)^{i_1} \end{aligned}$$

For the induction step, the parameterization of conjugacy classes of wreath products and Lemma 2 imply that

$$\begin{aligned}
& \sum_{n=0}^{\infty} \frac{u^n |Com(C_i Wr S_n, m)|}{|C_i Wr S_n|} \\
= & \sum_{n=0}^{\infty} \frac{u^n}{|C_i Wr S_n|} \sum_{\substack{(M_{j,k}): 1 \leq j \leq i \\ \sum_{j,k} k M_{j,k} = n}} \frac{|C_i Wr S_n|}{|C_{C_i Wr S_n}(M_{j,k})|} |Com(C_{C_i Wr S_n}(M_{j,k}), m-1)| \\
= & \left[\prod_{k=1}^{\infty} \sum_{a=0}^{\infty} \frac{u^{ka} |Com(C_{ik} Wr S_a, m-1)|}{|C_{ik} Wr S_a|} \right]^i \\
= & \left[\prod_{k=1}^{\infty} \prod_{i_2, \dots, i_{m-1}=1}^{\infty} \left(\frac{1}{1 - u^{ki_2 \dots i_{m-1}}} \right)^{(ik)^{m-2} i_2^{m-3} \dots i_{m-2}} \right]^i \\
= & \prod_{i_1, \dots, i_{m-1}=1}^{\infty} \left(\frac{1}{1 - u^{i_1 \dots i_{m-1}}} \right)^{i_1^{m-1} i_2^{m-2} i_3^{m-3} \dots i_{m-2}}
\end{aligned}$$

□

References

- [1] Atiyah, M. and Segal, G., On equivariant Euler characteristics. J. Geom. Phys. **6**, no. 4, 671-677 (1989).
- [2] James, G. and Kerber, A., The representation theory of the symmetric group. Encyclopedia of Mathematics and its Applications. Volume 16, (1981).
- [3] Hirzebruch, F. and Höfer, T., On the Euler number of an orbifold. Mathematische Annalen **286**, 255–260 (1990).
- [4] Hopkins, M., Kuhn, N., and Ravenel, D., Morava K -theories of classifying spaces and generalized characters for finite groups. Algebraic topology (San Feliu de Guixols), Lecture Notes in Math. **1509**, 186-209 (1990).

- [5] Macdonald, I.G., The Poincaré polynomial of a symmetric product. Proc. Camb. Phil. Soc.
58, 563-568 (1962).

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF CALIFORNIA

BERKELEY, CA 94720

DEPARTMENT OF MATHEMATICS

DARTMOUTH COLLEGE

HANOVER, NH 03755

`jbryan@math.berkeley.edu`

`jason.e.fulman@dartmouth.edu`