

# NON-DEGENERACY OF GAUSS CURVATURE EQUATION WITH NEGATIVE CONIC SINGULARITY

JUNCHENG WEI AND LEI ZHANG

**ABSTRACT.** We study the Gauss curvature equation with negative singularities. For a local mean field type equation with only one negative index we prove a uniqueness property. For a global equation with one or two negative indexes we prove the non-degeneracy of the linearize equations.

## 1. INTRODUCTION

In this article we study two closely related equations defined locally and globally in  $\mathbb{R}^2$ , respectively. The first equation is define in  $\Omega \subset \mathbb{R}^2$ , which is simply connected, open and bounded. Throughout the whole article we shall always assume that the boundary of  $\Omega$ , denoted as  $\partial\Omega$ , is a rectifiable Jordan curve. We say  $\Omega$  is regular from now on. Let  $p_0, p_1, \dots, p_m \in \Omega$  be a finite set in  $\Omega$ . Then we consider  $v$  as a solution of

$$(1.1) \quad \begin{cases} \Delta v + \lambda \frac{e^v}{\int_{\Omega} e^v} = -4\pi\alpha_0\delta_{p_0} + \sum_{i=1}^m 4\pi\alpha_i\delta_{p_i}, & \text{in } \Omega, \\ v = 0, & \text{on } \partial\Omega. \end{cases}$$

where  $\alpha_0 \in (0, 1)$ ,  $\alpha_1, \dots, \alpha_m > 0$  and  $\lambda \in \mathbb{R}$ .

The second equation is concerned with the stability of the following global equation: Suppose  $u$  is a solution of

$$(1.2) \quad \Delta u + e^u = \sum_{i=1}^N 4\pi\beta_i\delta_{p_i}, \quad \text{in } \mathbb{R}^2$$

where  $\beta_1, \dots, \beta_n$  are constants greater than  $-1$ ,  $p_1, \dots, p_n$  are the location of singular sources in  $\mathbb{R}^2$ . For this equation we shall prove that under some restrictions of  $\beta_i$ , any bounded solution of the linearized equation has to be the trivial solution.

The background of both equations is incredibly rich not only in mathematics but also in Physics. In particular, the study of (1.1) reveals core information on the configuration of vortices in the electroweak theory of Glashow-Salam-Weinberg [21] and Self-Dual Chern-Simons theories [18, 19, 20]. Also in statistical mechanics the behavior of solutions in (1.1) is closely related to Onsager's model of two-dimensional turbulence with vortex sources [8, 9]. Most of the motivation and application of both equations come from their

---

*Date:* June 8, 2017.

connection with conformal geometry. The singular sources represent conic singularities on a surface with constant curvature. There is a large amount of interesting works that discuss the qualitative properties of solutions of such equations. The readers may read into the following works and the references therein [2, 4, 5, 6, 7, 9, 10, 11, 12, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 35, 37, 38]. It is important to observe that it seems there are very few works to discuss singularities with negative strength and even fewer about the comparison between the negative indexes and positive ones. In this article, using an improved version of the Alexander-Bol's inequality we discuss the uniqueness property and the non-degeneracy for a local equation and a global equation. Our proof is based on techniques developed in a number of works of Bartolucci, Lin, Chang-Chen-Lin, etc.

To state the main result on the local equation, we first rewrite (1.1) using the following Green's function.

For  $p \in \Omega$ , let  $G_\alpha(x, p)$  satisfy

$$\begin{cases} \Delta G_\alpha(x, p) = 4\pi\alpha\delta_p, & \text{in } \Omega, \\ G_\alpha(x, p) = 0, & x \in \partial\Omega. \end{cases}$$

and

$$u = v - \sum_{j=0}^m G_{\alpha_j}(x, p_j).$$

Then  $u$  satisfies

$$(1.3) \quad \begin{cases} \Delta u + \lambda \frac{He^u}{\int_\Omega He^u} = 0, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

where

$$(1.4) \quad H(x) = \exp\left(\sum_{j=0}^m G_{\alpha_j}(x, p)\right) = e^{h(x)} |x - p_0|^{-2\alpha_0} \prod_{i=1}^m |x - p_i|^{2\alpha_i},$$

where  $h$  is harmonic in  $\Omega$  and is continuous up to the boundary.

The first main result is

**Theorem 1.1.** *Let  $u$  be a solution of (1.3) and  $H$  be defined by (1.4). Assume that  $\Omega$  is regular, then for any  $\lambda \leq 8\pi(1 - \alpha_0)$  there exists at most one solution to (1.1).*

Here we note that for  $\lambda < 8\pi(1 - \alpha_0)$ , the existence result has been established by Bartolucci-Malchiodi [6]. The existence result for  $\lambda = 8\pi(1 - \alpha_0)$  will be discussed in a separate work.

The second main result of this article is to consider the nondegeneracy of (1.2) when there are exactly two negative indexes:

$$(1.5) \quad \begin{cases} \Delta u + e^u = -4\pi\alpha_1\delta_{p_1} - 4\pi\alpha_2\delta_{p_2} + \sum_{i=3}^n 4\pi\beta_i\delta_{p_i} & \text{in } \mathbb{R}^2, \\ u(x) = -4\log|x| + \text{a bounded function near } \infty. \end{cases}$$

where  $\alpha_1, \alpha_2 \in (0, 1)$  and  $\beta_i > 0$  for  $i = 3, \dots, n$  and we assume that  $n \geq 3$ . The assumption of  $u$  at infinity says that  $\infty$  is not a singularity of  $u$  when  $\mathbb{R}^2$  is identified with  $\mathbb{S}^2$ .

Let

$$(1.6) \quad u_1(x) = u(x) + \sum_{i=1}^2 2\alpha_i \log|x - p_i| - 2 \sum_{i=3}^n \beta_i \log|x - p_i|$$

then clearly  $u_1$  satisfies

$$\begin{cases} \Delta u_1 + H_1 e^{u_1} = 0, & \text{in } \mathbb{R}^2, \\ u_1(x) = (-4 - 2\alpha_1 - 2\alpha_2 + 2 \sum_{i=3}^n \beta_i) \log|x| + O(1), & \text{for } |x| > 1, \end{cases}$$

where

$$(1.7) \quad H_1(x) = \prod_{i=1}^2 |x - p_i|^{-2\alpha_i} \prod_{i=3}^n |x - p_i|^{2\beta_i}, \quad \text{for } x \in \mathbb{R}^2.$$

Our second main result is

**Theorem 1.2.** *Let  $u$ ,  $u_1$  and  $H_1$  be defined as in (1.5), (1.6) and (1.7), respectively. Suppose  $\phi$  be a classical solution of*

$$(1.8) \quad \Delta\phi + H_1(x)e^{u_1}\phi = 0, \quad \text{in } \mathbb{R}^2.$$

*If  $\lim_{x \rightarrow \infty} |\phi(x)|/\log|x| = 0$  and  $\alpha_1, \alpha_2, \beta_i$  satisfy the following condition:*

$$(1.9) \quad -\max\{\alpha_1, \alpha_2\} + \min\{\alpha_1, \alpha_2\} + \sum_{i=3}^n \beta_i \leq 0,$$

*then  $\phi \equiv 0$ .*

Here we recall that the total angles at singularities are  $2\pi(1 - \alpha_1)$ ,  $2\pi(1 - \alpha_2)$ ,  $2\pi(1 + \beta_i)$  ( $i = 3, \dots, n$ ). For a surface  $S$  with conic singularities, let

$$\chi(S, \theta) = \chi(S) + \sum_i \left( \frac{\theta_i}{2\pi} - 1 \right)$$

where  $\theta_i$  is the total angle at a conic singularity.  $\chi(S, \theta)$  is to characterize three cases:

- (i) subcritical case if  $\chi(S, \theta) < \min_i \{2, \theta_i/\pi\}$ ,
- (ii) critical case if  $\chi(S, \theta) = \min_i \{2, \theta_i/\pi\}$ ,
- (iii) supercritical case if  $\chi(S, \theta) > \min_i \{2, \theta_i/\pi\}$ .

In our case  $\chi(S) = 2$  because  $S$  is the standard sphere. It is easy to see that (1.9) refers to the super-critical case. For the subcritical case Troyanov's

well known result [34] states that every conic singular metric is pointwise conformal to a metric with constant curvature.

Finally if there is only one negative singular source a similar result still holds: Let  $u$  satisfy

$$(1.10) \quad \begin{cases} \Delta u + e^u = -4\pi\alpha\delta_{p_1} + \sum_{i=2}^n 4\pi\beta_i\delta_{p_i} & \text{in } \mathbb{R}^2, \\ u(x) = -4\log|x| + \text{a bounded function near } \infty. \end{cases}$$

where  $\alpha \in (0, 1)$  and  $\beta_i > 0$  for  $i = 2, \dots, n$  and we assume that  $n \geq 3$ .

Let

$$u_1(x) = u(x) + 2\alpha \log|x - p_1| - 2 \sum_{i=2}^n \beta_i \log|x - p_i|$$

then clearly  $u_1$  satisfies

$$(1.11) \quad \begin{cases} \Delta u_1 + H_2 e^{u_1} = 0, & \text{in } \mathbb{R}^2, \\ u_1(x) = (-4 - 2\alpha + 2 \sum_{i=2}^n \beta_i) \log|x| + O(1), & \text{for } |x| > 1, \end{cases}$$

where

$$(1.12) \quad H_2(x) = |x - p_1|^{-2\alpha} \prod_{i=2}^n |x - p_i|^{2\beta_i}, \quad \text{for } x \in \mathbb{R}^2.$$

Our third main result is

**Theorem 1.3.** *Let  $u_1$  be a solution of (1.11) with  $H_2$  defined in (1.12). Let  $\phi$  be a classical solution of*

$$(1.13) \quad \Delta\phi + H_2(x)e^{u_1}\phi = 0, \quad \text{in } \mathbb{R}^2.$$

*If  $\lim_{x \rightarrow \infty} |\phi(x)|/\log|x| = 0$  and  $\alpha, \beta_i$  satisfy the following condition:*

$$(1.14) \quad -\alpha + \sum_{i=2}^n \beta_i \leq 0,$$

*then  $\phi \equiv 0$ .*

The organization of this article is as follows. In section two we derive a Bol's inequality with one negative singular source. Then in section three the first two eigenvalues of the linearized local equation is discussed. The proofs of major theorems are arranged in sections 4 and 5. The main approach of this article follows closely from previous works of Bartolucci, Chang, Chen and Lin, etc.

## 2. ON THE BOL'S INEQUALITY AND THE FIRST EIGENVALUES OF THE LOCAL EQUATION

One of the major tools we shall use is the following Bol's inequality:

**Proposition 2.1.** *Let  $\Omega \subset \subset \mathbb{R}^2$  be a simply connected, open and bounded domain in  $\mathbb{R}^2$ . Let  $u$  be a solution of*

$$\Delta u + V e^u = 0, \quad \text{in } \Omega$$

for

$$(2.1) \quad V = |x - p_1|^{-2\alpha_0} \prod_{i=2}^n |x - p_i|^{2\beta_i} e^g$$

and  $\Delta g \geq 0$  in  $\Omega$ . Here  $p_1, \dots, p_n$  ( $n \geq 2$ ) are distinct points in  $\Omega$ . Let  $\omega \subset \Omega$  be an open subset of  $\Omega$  such that  $\partial\omega$  is a finite union of rectifiable Jordan curves. Let

$$L_{\alpha_0}(\partial\omega) = \int_{\partial\omega} (Ve^u)^{1/2} ds, \quad M_{\alpha_0}(\omega) = \int_{\omega} Ve^u dx.$$

Then

$$(2.2) \quad 2L_{\alpha_0}^2(\partial\omega) \geq (8\pi(1 - \alpha_0) - M_{\alpha_0}(\omega))M_{\alpha_0}(\omega).$$

The strict inequality holds if  $\omega$  contains more than one singular source or is multiple connected.

Our proof of Proposition 2.1 is motivated by the argument in Bartolucci-Castorina [3] and Bartolucci-Lin [4, 5]. In fact if  $\alpha_0 = 0$  it was established by Bartolucci-Lin [4], if  $V$  has only singular source at 0, it was established by Bartolucci-Castorina. It all starts from an inequality of Huber [13]:

*Theorem A (Huber): Let  $\omega$  be an open, bounded, simply connected domain with  $\partial\omega$  being a rectifiable Jordan curve,  $\tilde{V} = |x|^{-2\alpha_0} e^g$  for some  $\Delta g \geq 0$  in  $\omega$ . Then*

$$\begin{aligned} \left( \int_{\partial\omega} \tilde{V}^{1/2} ds \right)^2 &\geq 4\pi(1 - \alpha_0) \int_{\omega} \tilde{V} dx, \quad \text{if } 0 \in \omega, \\ \left( \int_{\partial\omega} \tilde{V}^{1/2} ds \right)^2 &\geq 4\pi \int_{\omega} \tilde{V} dx, \quad \text{if } 0 \notin \omega, \end{aligned}$$

Huber's theorem can be adjusted to the following version

*Theorem B (Bartolucci-Castorina): Let  $\omega \subset \mathbb{R}^2$  be an open bounded domain such that  $\partial\omega$  is a rectifiable Jordan curve. Suppose  $\bar{\omega}_B$  is the closure of possibly disconnected bounded component of  $\mathbb{R}^2 \setminus \omega$  and  $\omega_B$  be the interior of  $\bar{\omega}_B$ . Let  $\tilde{V} = |x|^{-2\alpha_0} e^g$  for some  $g$  satisfying  $\Delta g \geq 0$  in the interior of  $\bar{\omega} \cup \bar{\omega}_B$ . Then*

$$\left( \int_{\partial\omega} \tilde{V}^{1/2} ds \right)^2 \geq 4\pi(1 - \alpha_0) \int_{\omega} \tilde{V} dx,$$

if 0 is in the interior of  $\bar{\omega} \cup \bar{\omega}_B$ .

$$\left( \int_{\partial\omega} \tilde{V}^{1/2} ds \right)^2 \geq 4\pi \int_{\omega} \tilde{V} dx,$$

if 0 is not in the interior of  $\bar{\omega} \cup \bar{\omega}_B$ .

**Proof of Proposition 2.1:** We shall only consider the first case mentioned in Theorem B because the other case corresponds to  $\alpha_0 = 0$ . Find

$$\begin{cases} \Delta q = 0, & \text{in } \omega, \\ q = u, & \text{on } \partial\omega. \end{cases}$$

and let  $\eta = u - q$ . Then the equation for  $\eta$  is

$$(2.3) \quad \begin{cases} \Delta\eta + Ve^q e^\eta = 0, & \text{in } \omega, \\ \eta = 0, & \text{on } \partial\omega, \end{cases}$$

and we use

$$t_m = \max_{\bar{\omega}} \eta.$$

Then we set

$$\Omega(t) = \{x \in \omega; \eta(x) > t\}, \quad \Gamma(t) = \partial\Omega(t), \quad \mu(t) = \int_{\Omega(t)} Ve^q dx.$$

Clearly  $\Omega(0) = \omega$ ,  $\mu(0) = \int_{\omega} Ve^q dx$ ,  $\mu(t_m) = \lim_{t \rightarrow t_m^-} \mu(t) = 0$ . Since  $\mu$  is continuous and strictly decreasing, it is easy to see that

$$(2.4) \quad \frac{d\mu(t)}{dt} = - \int_{\Gamma(t)} \frac{Ve^q}{|\nabla\eta|} ds, \quad a.e. \quad t \in [0, t_m].$$

For all  $s \in [0, \mu(0)]$ , set

$$\eta^*(s) = |\{t \in [0, t_m], \mu(t) > s\}|$$

where  $|E|$  is the Lebesgue measure of the measurable set  $E \in \mathbb{R}$ . It is easy to see that  $\eta^*$  is the inverse of  $\mu$  on  $[0, t_m]$  and is continuous, strictly monotone and differentiable almost everywhere. By (2.4) we have, for almost all  $s \in [0, \mu(0)]$ , that

$$(2.5) \quad \frac{d\eta^*}{ds} = - \left( \int_{\Gamma(\eta^*(s))} \frac{Ve^q}{|\nabla\eta|} dt \right)^{-1}.$$

Let

$$F(s) = \int_{\Omega(\eta^*(s))} e^\eta Ve^q dx, \quad a.e. \quad s \in [0, \mu(0)].$$

Then by the definition of  $\Omega(t)$  we see that

$$F(s) = \int_{\eta^*(s)}^{t_m} e^t \left( \int_{\Gamma_t} \frac{Ve^q}{|\nabla\eta|} ds \right) dt$$

Using  $\beta = \mu(t)$  we further have

$$(2.6) \quad F(s) = \int_0^s e^{\eta^*(\beta)} d\beta$$

where  $\eta^* = \mu^{-1}$  and (2.4) are used. The definition of  $F$  also gives

$$F(0) = \int_{\Omega(\eta^*(0))} e^\eta Ve^q = \int_{\Omega(t_m)} e^\eta Ve^q = 0$$

and  $F(\mu(0)) = \int_{\omega} e^\eta d\tau = M(\omega)$ . Consequently from (2.6) we obtain

$$(2.7) \quad \frac{dF}{ds} = e^{\eta^*(s)}, \quad \frac{d^2F}{ds^2} = \frac{d\eta^*}{ds} e^{\eta^*(s)} = \frac{d\eta^*}{ds} \frac{dF}{ds}. \quad a.e.s.$$

Here we use Bartolucci-Castorina's argument in [3] to show that  $\eta^*$  is locally lipschitz in  $(0, \mu(0))$ :

**Lemma 2.1.** *For any  $0 < \bar{a} \leq a < b \leq \bar{b} < \bar{u}(0)$ , there exists  $C(\bar{a}, \bar{b}, \beta_1, \dots, \beta_k) > 0$  such that*

$$\eta^*(a) - \eta^*(b) \leq C(b - a).$$

**Proof of Lemma 2.1:** First we find  $\Omega_{a,b}$  that satisfies

$$\{x \in \omega; \quad \eta^*(b) \leq \eta(x) \leq \eta^*(a)\} \subset \subset \Omega_{a,b} \subset \subset \omega.$$

Using Green's representation formula we have

$$|\nabla \eta(x)| \leq C + C \int_{\Omega_{a,b}} \frac{1}{|x - y|} |y - p_1|^{-2\alpha_0} dy.$$

Standard estimate gives

$$(2.8) \quad |\nabla \eta(x)| \leq C + C|x - p_0|^{1-2\alpha_0}.$$

Recall that  $d\eta = Ve^q dx$ . Thus

$$\begin{aligned} b - a &= \mu(\eta^*(b)) - \mu(\eta^*(a)) \\ &= \int_{\eta > \eta^*(b)} d\tau - \int_{\eta > \eta^*(a)} d\tau \geq \int_{\eta^*(b) < \eta < \eta^*(a)} d\tau \\ &= \int_{\eta^*(b)}^{\eta^*(a)} \left( \int_{\Gamma(t)} \frac{Ve^q}{|\nabla \eta|} ds \right) dt \end{aligned}$$

Using the expression of  $V$  in (2.1) and (2.8) we further have

$$\begin{aligned} b - a &\geq \frac{1}{C} \int_{\eta^*(b)}^{\eta^*(a)} \left( \int_{\Gamma(t)} \frac{1}{|x - p_0|^{2\alpha_0} + |x - p_0|} \right) dt \\ &\geq \frac{1}{C} \int_{\eta^*(b)}^{\eta^*(a)} L_1(\Gamma(t)) dt \\ &\geq \min_{\eta^*(b) \leq t \leq \eta^*(a)} L_1(\Gamma(t)) \int_{\eta^*(b)}^{\eta^*(a)} dt \\ &\geq C(\eta^*(a) - \eta^*(b)) \end{aligned}$$

where the estimate of  $\nabla \eta$  was used,  $L_1(\Gamma(t))$  stands for the Lebesgue measure of  $\Gamma$  and in the last inequality the following standard iso-perimetric inequality  $L_1(\Gamma(t)) \geq 4\pi|\Omega(t)| \geq 4\pi|\Omega(\eta^*(\bar{a}))| > 0$  is used. Lemma 2.1 is established.  $\square$

Now we go back to the proof of Proposition 2.1. By Cauchy's inequality

$$\begin{aligned} ( \int_{\Gamma(\eta^*(s))} (Ve^q)^{1/2} ds )^2 &\leq ( \int_{\Gamma(\eta^*(s))} \frac{Ve^q}{|\nabla \eta|} ds ) ( \int_{\Gamma(\eta^*(s))} |\nabla \eta| ds ) \\ (2.9) \quad &= (-\frac{d\eta^*}{ds})^{-1} ( \int_{\Gamma(\eta^*(s))} (-\frac{\partial \eta}{\partial \nu}) ds ), \quad \text{for a.e. } s \in [0, \mu(0)] \end{aligned}$$

where  $\nu = \nabla\eta/|\nabla\eta|$ . Moreover from (2.3)

$$(2.10) \quad \int_{\Gamma(\eta^*(s))} \left(-\frac{\partial\eta}{\partial\nu}\right) ds = \int_{\Omega(\eta^*(s))} V e^q e^\eta dx = F(s), \quad a.e.s \in [0, \mu(0)]$$

By Theorem A the following inequality holds for almost all  $s \in [0, \mu(0)]$ :

$$(2.11) \quad \left(\int_{\Gamma(\eta^*(s))} (V e^q)^{\frac{1}{2}} ds\right)^2 \geq 4\pi(1 - \alpha_0)\mu(\eta^*(s)) = 4\pi(1 - \alpha_0)s.$$

Putting (2.10) in (2.9) yields

$$(2.12) \quad \left(\int_{\Gamma(\eta^*(s))} (V e^q)^{\frac{1}{2}} ds\right)^2 \leq \left(-\frac{d\eta^*}{ds}\right)^{-1} F(s),$$

Using (2.11) in (2.12) we have

$$4\pi(1 - \alpha_0)s \leq \left(-\frac{d\eta^*}{ds}\right)^{-1} F(s), \quad a.e.s \in [0, \mu(0)],$$

which is equivalent to

$$(2.13) \quad 4\pi(1 - \alpha_0)s \frac{d\eta^*}{ds} + F(s) \geq 0, \quad a.e.s \in [0, \mu(0)].$$

By (2.7) and (2.13), we obtain

$$\frac{d}{ds} [4\pi(1 - \alpha_0)(s \frac{dF}{ds} - F(s)) + \frac{1}{2} F^2(s)] \geq 0, \quad a.e.s \in [0, \mu(0)].$$

Let  $P(s)$  denote the function in the brackets, then  $P$  is well defined, continuous, nondecreasing on  $[0, \mu(0)]$ . By the Lipschitz property of  $\eta^*$ ,  $P$  is absolutely continuous on  $[0, \mu(0)]$ ,

$$P(\mu(0)) - P(0) = \lim_{b \rightarrow \mu(0)^-} \lim_{a \rightarrow 0^+} \int_a^b \frac{dP}{ds} ds.$$

Using  $F(0) = 0$ ,  $F(\mu(0)) = M(\omega)$ ,  $\frac{dF}{ds}|_{s=\mu(0)} = e^0 = 1$ , we have

$$8\pi(1 - \alpha_0)(\mu(0) - M(\omega)) + M(\omega)^2 \geq 0.$$

Then Huber's inequality and  $\Gamma(0) = \partial\omega$  further yield

$$\begin{aligned} 2l^2(\partial\omega) &= 2\left(\int_{\partial\omega} (V e^v)^{1/2} ds\right)^2 \\ &= 2\left(\int_{\partial\omega} (V e^q)^{\frac{1}{2}} ds\right)^2 \\ &\geq 8\pi(1 - \alpha_0)\mu(0) \\ &\geq M(\omega)(8\pi(1 - \alpha_0) - M(\omega)). \end{aligned}$$

where we have used the fact that  $v = q$  on  $\partial\omega$ . The Bol's inequality is established. The equality holds if  $V e^q = |x - p_0|^{-2\alpha_0} |\Phi'_t|^2 e^k$  on  $\Omega(t)$  for almost all  $t \in (0, t_m)$  where  $k$  is a constant. In particular for  $t = 0$   $\Phi_0$  maps  $\Omega$  to a ball. In this case  $g$  must be harmonic. On the other hand from the equality of Cauchy's inequality we have

$$V e^q = c_t |\nabla\eta|^2, \quad \text{on } \Gamma(t), \quad a.e.t \in (0, t_m),$$

for some  $c_t > 0$ . Put  $w = \Phi_0(z)$  and  $\xi(w) = \eta(\Phi_0^{-1}(w)) + k$ , we see that  $\xi$  satisfies

$$\Delta\xi + |x|^{-2\alpha_0}e^\xi = 0,$$

and  $\xi$  is radial. This  $\xi$  is a scaling of

$$\log \frac{8(1-\alpha_0)^2}{1+|x|^{2(1-\alpha_0)})^2}.$$

Thus we have strict inequality in Bol's inequality if at least one of the following situations occurs:

- (1)  $p_1 \notin \omega$ ,
- (2)  $\omega$  has at least two singular sources
- (3)  $\omega$  is not simply connected.

□

### 3. THE FIRST EIGENVALUES OF THE LINEARIZED LOCAL EQUATION

**Proposition 3.1.** *Let  $\Omega$  be an open, bounded domain of  $\mathbb{R}^2$  with rectifiable boundary  $\partial\Omega$ ,  $V = |x|^{-\alpha_0} \prod_{i=1}^k |x-p_i|^{2\beta_i} e^g$  for some subharmonic and smooth function  $g$ ,  $\alpha_0 \in (0, 1)$ ,  $\beta_1, \dots, \beta_k > 0$ , and we assume all the singularities  $0, p_1, \dots, p_k$  are in  $\Omega$ . Let  $w$  be a classical solution of*

$$\Delta w + V e^w = 0, \quad \text{in } \Omega.$$

Suppose  $\hat{\nu}_1$  is the first eigenvalue of

$$(3.1) \quad \begin{cases} -\Delta\phi - V e^w \phi = \hat{\nu}_1 V e^w \phi, & \text{in } \Omega, \\ \phi = 0, & \text{on } \partial\Omega. \end{cases}$$

Then if  $\int_\Omega V e^w \leq 4\pi(1-\alpha_0)$  we have  $\hat{\nu}_1 > 0$ . Moreover if  $\int_\Omega V e^w \leq 8\pi(1-\alpha_0)$  we have  $\hat{\nu}_2 > 0$

**Proof:** Let  $\nu_1 = \hat{\nu}_1 + 1$  and  $\phi$  the eigenfunction corresponding to  $\hat{\nu}_1$ , then we have  $\phi > 0$  and

$$\begin{cases} -\Delta\phi = \nu_1 V e^w \phi, & \text{in } \Omega, \\ \phi = 0, & \text{on } \partial\Omega. \end{cases}$$

Let

$$U_0(x) = (-2) \log(1 + |x|^{2(1-\alpha_0)}) + \log(8(1-\alpha_0)^2).$$

Then clearly  $U_0$  solves

$$\Delta U_0 + |x|^{-2\alpha_0} e^{U_0} = 0, \quad \text{in } \mathbb{R}^2.$$

For  $t \in (0, t_+)$  where  $t_+ = \max_{\bar{\Omega}} \phi$ , we set  $\Omega(t) = \{x \in \Omega, \phi(x) > t\}$  and we set  $R(t)$  to satisfy

$$\int_{\Omega(t)} V e^w = \int_{B_{R(t)}} e^{U_0} |x|^{-2\alpha_0}.$$

Clearly  $\Omega(0) = \Omega$ ,  $R_0 = \lim_{t \rightarrow 0+} R(t)$ ,  $\lim_{t \rightarrow t_+ -} R(t) = 0$ . Let  $\phi^*$  be a radial function from  $B_{R_0} \rightarrow \mathbb{R}$ . For  $y \in B_{R_0}$  and  $|y| = r$ , set

$$\phi^*(r) = \sup\{t \in (0, t_+) \mid R(t) > r\}.$$

Then  $\phi^*(R_0) = \lim_{r \rightarrow R_0-} \phi^*(r) = 0$ , and the definition implies

$$B_{R(t)} = \{y \in \mathbb{R}^2, \phi^*(y) > t\}.$$

$$\int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0} = \int_{\Omega(t)} V e^w, \quad t \in [0, t_+].$$

$$\int_{B_{R_0}} |x|^{-2\alpha_0} e^{U_0} |\phi^*|^2 = \int_{\Omega} V e^w \phi^2.$$

Then for almost all  $t$

$$\begin{aligned} (3.2) \quad & -\frac{d}{dt} \int_{\Omega(t)} |\nabla \phi|^2 = \int_{\phi=t} |\nabla \phi| \\ & \geq \left( \int_{\phi=t} (V e^w)^{1/2} ds \right)^2 \left( \int_{\phi=t} \frac{V e^w}{|\nabla \phi|} ds \right)^{-1}, \\ & = \left( -\frac{d}{dt} \int_{\Omega(t)} V e^w \right)^{-1} \left( \int_{\phi=t} (V e^w)^{1/2} ds \right)^2 \\ & \geq \frac{1}{2} (8\pi(1-\alpha_0) - \int_{\Omega(t)} V e^w) \left( \int_{\Omega_t} V e^w \right) \left( -\frac{d}{dt} \int_{\Omega(t)} V e^w \right)^{-1}, \\ & = \frac{1}{2} (8\pi(1-\alpha_0) - \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0}) \left( \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0} \right) \\ & \quad \cdot \left( -\frac{d}{dt} \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0} \right)^{-1}. \end{aligned}$$

Applying the same computation to  $\phi^*$  we see that for almost all  $t$ , since  $\phi^*$  is radial, we have

$$\begin{aligned} & -\frac{d}{dt} \int_{\Omega(t)} |\nabla \phi^*|^2 = \int_{\phi^*=t} |\nabla \phi^*| \\ & = \left( \int_{\phi^*=t} |x|^{-\alpha_0} e^{U_0/2} ds \right)^2 \left( \int_{\phi^*=t} \frac{|x|^{-2\alpha_0} e^{U_0}}{|\nabla \phi^*|} ds \right)^{-1} \\ & = \left( -\frac{d}{dt} \int_{\Omega(t)} |x|^{-2\alpha_0} e^{U_0} \right)^{-1} \left( \int_{\phi^*=t} |x|^{-\alpha_0} e^{U_0/2} ds \right)^2. \end{aligned}$$

Direct computation on  $U_0$  gives

$$\left( -\frac{d}{dt} \int_{\Omega(t)} |x|^{-2\alpha_0} e^{U_0} \right)^{-1} = \frac{1}{2} (8\pi(1-\alpha_0) - \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0}) \left( \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0} \right).$$

Thus the combination of the two equations above gives

$$(3.3) \quad \begin{aligned} & -\frac{d}{dt} \int_{\Omega(t)} |\nabla \phi^*|^2 \\ &= \frac{1}{2} (8\pi(1-\alpha_0) - \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0}) (\int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0}) (-\frac{d}{dt} \int_{\phi^* > t} e^{U_0} |x|^{-2\alpha_0})^{-1}. \end{aligned}$$

for almost all  $t \in (0, t_+)$ .

Integrating (3.2) and (3.3) for  $t \in (0, t_+)$  we have

$$\int_{B_{R_0}} |\nabla \phi^*|^2 \leq \int_{\Omega} |\nabla \phi|^2.$$

If  $\nu_1 \leq 1$ , we obtain from (3.1) that

$$\begin{aligned} 0 &\geq (\nu_1 - 1) \int_{\Omega} V e^w |\phi|^2 = \int_{\Omega} |\nabla \phi|^2 - \int_{\Omega} V e^w |\phi|^2 \\ &\geq \int_{B_{R_0}} |\nabla \phi^*|^2 - \int_{B_{R_0}} e^{U_0} |x|^{-2\alpha_0} |\phi^*|^2. \end{aligned}$$

Thus the first eigenvalue of

$$-\Delta - |x|^{-2\alpha_0} e^{U_0}$$

on  $B_{R_0}$  with Dirichlet boundary condition is non-positive. Since

$$\psi = 2(1-\alpha_0) \frac{1-|x|^{2(1-\alpha_0)}}{1+|x|^{2(1-\alpha_0)}}$$

satisfies

$$-\Delta \psi - |x|^{-2\alpha_0} e^{U_0} \psi = 0 \quad \text{in } \mathbb{R}^2,$$

we see that  $R_0 \geq 1$ . But

$$\int_{B_1} |x|^{-2\alpha_0} e^{U_0} = 4\pi(1-\alpha_0),$$

we clearly have  $\hat{\nu} \geq 0$ . From the proof of the Bol's inequality we see that the strictly inequality holds because  $\Omega$  has more than one singular points in its interior.

The proof of  $\hat{\nu}_2 > 0$  for a higher thresh-hold of  $\int_{\Omega} V e^w$  is very similar. If we consider  $\Omega_+$  and  $\Omega_-$ , which are the set of points where  $\phi$  is positive or negative, respectively. Then the integral of  $V e^w$  on at least one of them is less than or equal to  $4\pi(1-\alpha_0)$ . The argument of re-distribution of mass can be applied to at least one of them. Then we see that either one of them has the integral of  $V e^w$  strictly less than  $4\pi(1-\alpha_0)$ , which leads to a contradiction, or both regions have their integral equal to  $4\pi(1-\alpha_0)$ . In the latter case the equality cannot hold because 0 can only be in the interior of at most one region. Then at least one region either does not contain 0 in its interior, or is not simply connected. The strictly inequality holds in at least one region. Thus  $\hat{\nu}_2 > 0$  if  $\int_{\Omega} V e^w \leq 8\pi(1-\alpha_0)$ .  $\square$

## 4. PROOF OF THEOREM 1.2

First we claim that  $\phi$  in the linearized equation is actually bounded. Recall that  $u_1$  satisfies

$$\begin{aligned} \Delta u_1 + H_1 e^{u_1} &= 0, \quad \text{in } \mathbb{R}^2, \\ u_1(x) &= (-4 + 2\alpha_1 + 2\alpha_2 - 2 \sum_{i=3}^n \beta_i) \log |x| + O(1), \quad \text{at } \infty. \end{aligned}$$

By the equation for  $\phi$  and the mild growth rate of  $\phi$  at infinity, we have

$$\phi(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \log|x-y| H_1(y) e^{u_1(y)} \phi(y) dy + c, \quad x \in \mathbb{R}^2$$

for some  $c \in \mathbb{R}$ .

Differentiating the equation above, we have

$$\partial_i \phi(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{x_i - y_i}{|x-y|^2} H_1 e^{u_1} \phi(y) dy, \quad i = 1, 2, \quad x \in \mathbb{R}^2.$$

By standard estimates in different regions of  $\mathbb{R}^2$ , it is easy to see that

$$\partial_i \phi(x) = A \frac{x_i}{|x|^2} + O(|x|^{-1-\delta}), \quad |x| > 1, \quad i = 1, 2.$$

for  $A = \frac{1}{2\pi} \int_{\mathbb{R}^2} H_1 e^{u_1} \phi$  and some  $\delta > 0$ . Thus the assumption  $\lim_{|x| \rightarrow \infty} \phi(x)/\log|x| = 0$  actually implies

$$(4.1) \quad \int_{\mathbb{R}^2} H_1 e^{u_1} \phi = 0.$$

and

$$(4.2) \quad \phi(x) = C + O(|x|^{-\delta}), \quad |x| > 1$$

for some  $\delta > 0$ .

Next we make a transformation on the equation for  $u_1$ . Without loss of generality we assume  $p_1 = 0$  and we write  $H_1$  as

$$H_1(x) = |x|^{-2\alpha_1} V_1.$$

Let

$$u_2(x) = u_1\left(\frac{x}{|x|^2}\right) - (4 - 2\alpha_1) \log|x|,$$

then direct computation shows that

$$\Delta u_2 + V_2 e^{u_2} = 0, \quad \text{in } \mathbb{R}^2$$

and

$$u_2(x) = (-4 + 2\alpha_1) \log|x| + O(1), \quad \text{at } \infty.$$

where  $V_2(x) = V_1(x/|x|^2)$ . It is also easy to verify that

$$(4.3) \quad \int_{\mathbb{R}^2} H_1 e^{u_1} = \int_{\mathbb{R}^2} V_2 e^{u_2}.$$

Setting  $\phi_1(x) = \phi(x/|x|^2)$ , we see that

$$\Delta \phi_1 + V_2 e^{u_2} \phi_1 = 0, \quad \text{in } \mathbb{R}^2.$$

Here we note that by the bound of  $\phi_1$  near the origin the equation above holds in the whole  $\mathbb{R}^2$ .

First by the asymptotic behavior of  $u_1$  at infinity, integration of the equation for  $u_1$  gives

$$(4.4) \quad \frac{1}{2\pi} \int_{\mathbb{R}^2} H_1 e^{u_1} = 4 - 2(\alpha_1 + \alpha_2) + 2 \sum_{i=3}^n \beta_i \leq 4(1 - \alpha_2).$$

From the definition of  $\phi$  we have  $\phi_1(x) \rightarrow c_0$  as  $x \rightarrow \infty$  for some  $c_0 \in \mathbb{R}$ . Without loss of generality we assume  $c_0 \leq 0$ . By the same estimate for  $\phi$  we have

$$(4.5) \quad \int_{\mathbb{R}^2} V_2 e^{u_2} \phi_1 = 0.$$

By (4.3) and (4.4) we have

$$\int_{\mathbb{R}^2} V_2 e^{u_2} \leq 8\pi(1 - \alpha_2).$$

Let  $\phi_2$  be an eigenfunction corresponding to eigenvalue  $\hat{\nu}$ :

$$\begin{cases} -\Delta \phi_2 - V_2 e^{u_2} \phi_2 = \hat{\nu} V_2 e^{u_2} \phi_2, & \text{in } \mathbb{R}^2, \\ \lim_{x \rightarrow \infty} \phi_2(x) = c_0 \leq 0, \\ \int_{\mathbb{R}^2} V_2 e^{u_2} \phi_2 = 0. \end{cases}$$

We claim that  $\hat{\nu} > 0$ .

By way of contradiction we assume that  $\hat{\nu} \leq 0$ . By setting  $\nu = 1 + \hat{\nu}$  we clearly have  $\nu \leq 1$  and

$$\Delta \phi_2 + \nu V_2 e^{u_2} \phi_2 = 0, \quad \text{in } \mathbb{R}^2.$$

Let  $\Omega^+ = \{x; \phi_2(x) > c_0\}$  then by the same argument as in the proof of the previous proposition we must have

$$\int_{\Omega^+} V_2 e^{u_2} = c_2(c_0) \geq 4\pi(1 - \alpha_2)$$

and if the equality holds, we have  $c_0 = 0$ , there is one singular source with negative index  $-4\pi\alpha_2$  in the interior of  $\Omega_+$ , which has to be simply connected at the same time. All other singular sources (which have positive indexes) are not in the interior of  $\Omega_+$ .

Let  $\phi^*$  be the rearrangement of  $\phi_2$  in  $\Omega_+$ . By the previous argument we have

$$\int_{\Omega_+} |\nabla \phi_2|^2 \leq \int_{B_{R_1}} |\nabla \phi^*|^2$$

and  $c_2(c_0) = \int_{B_{R_1}} |x|^{-2\alpha_2} e^{U_0}$ . Let

$$c_1 = \min_{\mathbb{R}^2} \phi_2$$

and we set  $R_2$  to make

$$\int_{B_{R_2} \setminus B_{R_1}} |x|^{-2\alpha_2} e^{U_0} = \int_{\mathbb{R}^2 \setminus \Omega_+} V_1 e^{u_2}.$$

Note that  $R_2$  could be  $\infty$ . Then we define a radial function  $\phi^{**}$  from  $B_{R_2} \setminus B_{R_1} \rightarrow \mathbb{R}$ : for any  $y \in B_{R_2} \setminus B_{R_1}$ ,  $|y| = r$ ,

$$\phi^{**}(r) = \inf\{t \in (c_1, c_0) \mid R^{(-)}(t) < r\},$$

where  $R^{(-)}(t)$  is defined by

$$\int_{B_{R_2} \setminus B_{R^{(-)}(t)}} |x|^{-2\alpha_2} e^{U_0} = \int_{\phi_2 < t} V_2 e^{u_2}, \quad \forall t \in (c_1, c_0).$$

The definition of  $\phi^{**}$  implies

$$\int_{B_{R_2} \setminus B_{R_1}} |x|^{-2\alpha_2} e^{U_0} |\phi^{**}|^2 = \int_{\Omega^-} V_2 e^{u_2} |\phi_2|^2, \quad \Omega^- = \mathbb{R}^2 \setminus \Omega_+,$$

and

$$\int_{B_{R_2} \setminus B_{R_1}} |x|^{-2\alpha_2} e^{U_0} \phi^{**} = \int_{\Omega^-} V_2 e^{u_2} \phi_2, \quad \Omega^- = \mathbb{R}^2 \setminus \Omega_+.$$

The symmetrization also gives

$$\int_{B_{R_2} \setminus B_{R_1}} |\nabla \phi^{**}|^2 \leq \int_{\Omega^-} |\nabla \phi_2|^2.$$

Now we set

$$\phi_* : B_{R_2} \rightarrow \mathbb{R}, \quad \phi_* \text{ radial } \phi_*(r) = \begin{cases} \phi^*(r), & r \in [0, R_1], \\ \phi^{**}(r), & r \in [R_1, R_2]. \end{cases}$$

Since  $\phi_*$  is continuous, monotone, we have

$$\int_{B_{R_2}} |\nabla \phi_*|^2 \leq \int_{\mathbb{R}^2} |\nabla \phi_2|^2 = \int_{\mathbb{R}^2} V_2 e^{u_2} |\phi_2|^2 = \int_{B_{R_2}} |x|^{-2\alpha_2} e^{U_0} |\phi_*|^2.$$

From the definition of  $\phi_*$  we also have

$$\int_{B_{R_2}} |x|^{-2\alpha_2} e^{U_0} \phi_* = 0.$$

Let

$$\begin{aligned} K^* &= \inf\left\{\int_{\mathbb{R}^2} |\nabla \psi|^2 dx, \quad \psi \text{ is radial,}\right. \\ &\quad \left.\int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \psi = 0, \quad \int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \psi^2 = 1\right\}. \end{aligned}$$

By Hölder's inequality we have

$$\left| \int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \psi dx \right| \leq \left( \int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \psi^2 \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \right)^{1/2}.$$

Which implies that the minimizer (say  $\psi^*$ ) also satisfies

$$\int_{\mathbb{R}^2} |x|^{-2\alpha_2} e^{U_0} \psi^* = 0.$$

Clearly the minimizer  $\psi^*$  satisfies

$$\Delta \psi^* + K^* |x|^{-2\alpha_2} e^{U_0} \psi^* = 0, \quad \text{in } \mathbb{R}^2.$$

From  $\phi_*$  and the definition of  $K^*$  we already know  $K^* \in (0, 1)$ . Our goal is to show that  $K^* = 1$  by an argument of Chang-Chen-Lin [2].  $\psi^*$  should only change sign once. Let  $\xi_0$  be the zero of  $\psi^*$ .

Integrating the equation for  $\psi^*$ , we have

$$r \frac{d}{dr} \psi^*(r) = -K^* \int_0^r |s|^{1-2\alpha_2} e^{U_0(s)} \psi^*(s) ds = K^* \int_r^\infty s^{1-2\alpha_2} e^{U_0} \psi^*(s) ds < 0$$

for  $r > \xi_0$ . Thus  $\psi^*$  is decreasing for  $r \geq \xi_0$  and  $r \frac{d}{dr} \psi^*(r) \rightarrow 0$  as  $r \rightarrow \infty$ . The equation for  $\psi^*$  also gives

$$|r \frac{d}{dr} \psi^*(r)| \leq K^* \left( \int_r^\infty |s|^{1-2\alpha_2} e^{U_0} (\psi^*(s))^2 ds \right)^{1/2} \left( \int_r^\infty s^{1-2\alpha_2} e^{U_0(s)} ds \right)^{1/2} \leq C r^{-1}$$

for large  $r$ . Therefore  $\lim_{r \rightarrow \infty} \psi^*(r)$  exists and is a negative constant.

Let

$$\psi(r) = 2(1 - \alpha_2) \frac{1 - r^{2(1-\alpha_2)}}{1 + r^{2(1-\alpha_2)}}.$$

Then  $\psi$  satisfies

$$\Delta \psi + r^{-2\alpha_2} e^{U_0} \psi = 0, \quad \text{in } \mathbb{R}^2.$$

It is easy to obtain from the equation for  $\psi$  and the one for  $\psi^*$  the following:

$$r \left( \frac{\psi^*}{\psi(r)} \right)' = \frac{1 - K^*}{\psi^2(r)} \int_0^r s^{1-2\alpha_2} e^{U_0(s)} \psi^*(s) \psi(s) ds.$$

If  $\xi_0 < 1$ ,  $\frac{\psi^*(r)}{\psi(r)}$  is increasing from  $r \in (0, \xi_0]$ . Clearly this is not possible because otherwise this could happen:

$$0 < \frac{\psi^*(0)}{\psi(0)} < \frac{\psi^*(\xi_0)}{\psi(\xi_0)} = 0.$$

On other hand we observe that it is also absurd to have  $\xi_0 > 1$ , indeed, had this happened, we would start from

$$\lim_{R \rightarrow \infty} R \left( \frac{\psi^*}{\psi} \right)'(R) \psi^2(R) - r \left( \frac{\psi^*}{\psi} \right)'(r) \psi^2(r) = (1 - K^*) \int_r^\infty s^{1-2\alpha_2} e^{U_0} \psi^*(s) \psi(s) ds.$$

Since

$$\lim_{R \rightarrow \infty} R \left( \frac{d}{dr} \psi^*(R) \psi(R) - \psi'(R) \psi^*(R) \right) = 0,$$

we have

$$-r \left( \frac{\psi^*}{\psi} \right)' \psi^2(r) = (1 - K^*) \int_r^\infty s^{1-2\alpha_2} e^{U_0(s)} \psi^*(s) \psi(s) ds.$$

If  $\xi_0 > 1$ ,  $\frac{\psi^*(r)}{\psi(r)}$  is decreasing for  $r > 1$ , which yields

$$0 = \frac{\psi^*(\xi_0)}{\psi(\xi_0)} > \lim_{r \rightarrow \infty} \frac{\psi^*(r)}{\psi(r)} = -\frac{1}{2(1-\alpha_2)} \lim_{r \rightarrow \infty} \psi^*(r) > 0.$$

This contradiction proves that  $\xi_0 = 1$  and  $\psi^*(r)\psi(r) > 0$  for all  $r \neq 1$ . Furthermore

$$\begin{aligned} 0 &= \lim_{r \rightarrow \infty} \left( \frac{d}{dr} \psi^*(r)\psi(r) - \frac{d}{dr} \psi(r)\psi^*(r) \right) r \\ &= (1 - K^*) \int_0^\infty s^{1-2\alpha_2} e^{U_0} \psi^*(s)\psi(s) ds. \end{aligned}$$

Thus we have proved that  $K^* = 1$  and the desired contradiction. Theorem 1.2 is established.  $\square$

The proof of Theorem 1.3 is very similar, we just use Kelvin transformation to move the negative singularity to infinity, then use the same argument with the standard Bol's inequality for non-negative indexes.

## 5. THE PROOF OF THEOREM 1.1.

Our argument follows from a previous result of Bartolucci-Lin [5] for non-negative indexed singularities. We prove by way of contradiction. Suppose  $u$  is a solution of (1.3) and a nonzero function  $\phi \in H_0^1(\Omega)$  is a solution of

$$\begin{cases} -\Delta \tilde{\phi} - \lambda \frac{He^u}{\int_\Omega He^u dx} \tilde{\phi} + \lambda (\int_\Omega He^u \tilde{\phi}) \frac{He^u}{(\int_\Omega He^u)^2} = 0, & \text{in } \Omega, \\ \tilde{\phi} = 0, & \text{on } \partial\Omega. \end{cases}$$

Let  $w = u + \log \lambda - \log(\int_\Omega He^u dx)$  and

$$\phi = \tilde{\phi} - \frac{\int_\Omega He^u \tilde{\phi}}{\int_\Omega He^u},$$

we have

$$(5.1) \quad \begin{cases} \Delta \phi + He^w \phi = 0, & \text{in } \Omega, \\ \phi = c_0, & \text{on } \partial\Omega, \\ \int_\Omega He^w \phi = 0, \\ \lambda = \int_\Omega He^w \leq 8\pi(1-\alpha_0). \end{cases}$$

Without loss of generality we assume  $c_0 \leq 0$ . Our goal is to show that  $\phi \equiv c_0$ , which further leads to  $c_0 = 0$  obviously. If  $c_0 = 0$ ,  $\phi$  must change sign if not identically equal to 0. But this situation is ruled out by Proposition 3.1 that  $\nu_2 > 0$ . So we only consider  $c_0 < 0$ . Let

$$\Omega_+ = \{x \in \Omega, \phi(x) > 0\}, \quad \Omega_- = \{x \in \Omega, \phi(x) < 0\}.$$

Clearly  $\text{dist}(\Omega_+, \partial\Omega) > 0$ . Then if  $\int_{\Omega_+} He^w \leq 4\pi(1-\alpha_0)$  there is no way for  $\phi$  to satisfy (5.1) on  $\Omega_+$  without being identically zero. Then using the

same rearrangement argument as in the proof of Theorem 1.2 we can also reach the following conclusion: If  $\phi_2$  is a solution of

$$\begin{cases} -\Delta\phi_2 - \lambda e^u w\phi_2 = \nu e^u w\phi_2, & \text{in } \Omega, \\ \phi_2 = c_0, & \text{on } \partial\Omega, \end{cases}$$

then  $\nu > 0$ . The remaining part of the proof of Theorem 1.1 follows by standard argument in [2] and [5]. We include it with necessary modification.

If we use  $L_\lambda$  to denote the linearized operator of (1.3), we know that all eigenvalues of  $L_\lambda$  are strictly positive for  $\lambda \in [0, 8\pi(1 - \alpha_0)]$ . By using the improved Moser-Trudinger inequality [25] one can easily find a solution of (1.3) by direct minimization method. By the uniform estimate of the linearized equation and standard elliptic estimate we have: for any  $\epsilon \in (0, 8\pi(1 - \alpha_0))$ ,

$$(5.2) \quad \|u_\lambda\|_\infty \leq \lambda C_\epsilon$$

for some  $C_\epsilon > 0$ ,  $\lambda \in [0, 8\pi(1 - \alpha_0)]$  and  $u_\lambda$  as solution of (1.3). Let  $S_\lambda$  be the solution's branch for (1.3) bifurcating from  $(u, \lambda) = (0, 0)$ . The standard bifurcation theory of Crandall-Rabinowitz [16] gives that  $S_\lambda$  is a simple branch near  $\lambda = 0$ . Which means for  $\lambda > 0$  small there exists one and only solution for (1.3) and  $S_\lambda$  is smooth in  $C^2(\Omega) \times \mathbb{R}$ . By the implicit function theorem (because  $L_\lambda$  has positive first eigenvalue)  $S_\lambda$  can be extended uniquely for  $\lambda \in (0, 8\pi(1 - \alpha_0))$ . If for any given  $\lambda \in (0, 8\pi(1 - \alpha_0))$  there is another solution, it implies another solution's branch, which does not bend in  $[0, 8\pi(1 - \alpha_0))$ . By the uniform estimate (5.2) this second branch intersects  $S_\lambda$  at  $(u, \lambda) = (0, 0)$ . This contradiction proves the uniqueness for  $\lambda \in [0, 8\pi(1 - \alpha_0))$ . If a solution exists for  $\lambda = 8\pi(1 - \alpha_0)$ , the implicit function theorem and the uniqueness result can be combined to prove the uniqueness in this case as well. Theorem 1.1 is established.  $\square$

## REFERENCES

- [1] Bandle , C. ( 1976 ). On a differential inequality and its applications to geometry . Math. Zeit. 147 : 253 – 261 .
- [2] Chang, Sun-Yung A.; Chen, Chiun-Chuan; Lin, Chang-Shou Extremal functions for a mean field equation in two dimension. Lectures on partial differential equations, 6193, New Stud. Adv. Math., 2, Int. Press, Somerville, MA, 2003.
- [3] Bartolucci, Daniele; Castorina, Daniele Self-gravitating cosmic strings and the Alexandrov's inequality for Liouville-type equations. Commun. Contemp. Math. 18 (2016), no. 4, 1550068, 26 pp.
- [4] Bartolucci, Daniele; Lin, Chang-Shou Existence and uniqueness for mean field equations on multiply connected domains at the critical parameter. Math. Ann. 359 (2014), no. 1-2, 144.
- [5] Bartolucci, D.; Lin, C. S. Uniqueness results for mean field equations with singular data. Comm. Partial Differential Equations 34 (2009), no. 7-9, 676702.
- [6] Bartolucci, D; Malchiodi, A. An improved geometric inequality via vanishing moments, with applications to singular Liouville equations, Commun. Math. Phys. 322, 415-452 (2013).

- [7] Bartolucci D., Tarantello G.: Liouville type equations with singular data and their application to periodic multivortices for the electroweak theory. *Commun. Math. Phys.* 229, 347 (2002)
- [8] Caglioti E., Lions P.L., Marchioro C., Pulvirenti M.: A special class of stationary flows for two dimensional Euler equations: a statistical mechanics description. II. *Commun. Math. Phys.* 174, 229260 (1995)
- [9] Chanillo S., Kiessling M.: Rotational symmetry of solutions of some nonlinear problems in statistical mechanics and in geometry. *Commun. Math. Phys.* 160, 217238 (1994)
- [10] Chen C.C., Lin C.S., Wang G.: Concentration phenomena of two-vortex solutions in a Chern-Simons model. *Ann. Scuola Norm. Sup. Pisa III*, 367397 (2004).
- [11] Chen C.C., Lin C.S.: Mean field equations of liouville type with singular data: sharper estimates. *Discr. Cont. Dyn. Syst.* 28(3), 12371272 (2010)
- [12] Chen, C. C, Lin, C. S. Mean field equation of Liouville type with singular data: topological degree. *Comm. Pure Appl. Math.* 68 (2015), no. 6, 887947.
- [13] Huber, Alfred On the isoperimetric inequality on surfaces of variable Gaussian curvature. *Ann. of Math.* (2) 60, (1954). 237247.
- [14] Chen W., Li C.: Qualitative properties of solutions of some nonlinear elliptic equations in R 2. *Duke Math. J.* 71(2), 427439 (1993)
- [15] Chen, Wen Xiong; Li, Congming What kinds of singular surfaces can admit constant curvature? *Duke Math. J.* 78 (1995), no. 2, 437451.
- [16] Crandall, Michael G.; Rabinowitz, Paul H. Some continuation and variational methods for positive solutions of nonlinear elliptic eigenvalue problems. *Arch. Rational Mech. Anal.* 58 (1975), no. 3, 207218.
- [17] Ding W., Jost J., Li J., Wang G.: An analysis of the two-vortex case in the Chern-Simons-Higgs model. *Calc. Var. P.D.E.* 7, 8797 (1998)
- [18] Dunne, G.: Self-dual Chern-Simons Theories. Lecture Notes in Physics 36, Berlin-Heidelberg-New York: Springer, 1995.
- [19] Hong J., Kim Y., Pac P.Y.: Multivortex Solutions of the Abelian Chern-Simons Theory. *Phys. Rev. Lett.* 64, 22302233 (1990)
- [20] Jackiw R., Weinberg E.J.: Selfdual Chern-Simons vortices. *Phys. Rev. Lett.* 64, 22342237 (1990)
- [21] Lai, C.H. (ed.): Selected Papers on Gauge Theory of Weak and Electromagnetic Interactions. Singapore: World Scientific, 1981
- [22] Li, Y. Y. Harnack type inequality: The method of moving planes. *Commun. Math. Phys.* 200(2), 421444 (1999).
- [23] Lin C.S., Wei J.C., Ye D.: Classification and nondegeneracy of  $SU(n + 1)$  Toda system with singular sources. *Invent. Math.* 190(1), 169207 (2012).
- [24] Luo, F; Tian, G; Liouville equation and spherical convex polytopes. *Proc. Amer. Math. Soc.* 116 (1992), no. 4, 11191129.
- [25] Malchiodi A., Ruiz D.: New improved Moser-Trudinger inequalities and singular Liouville equations on compact surfaces. *G.A.F.A.* 21(5), 11961217 (2011).
- [26] Malchiodi A., Ruiz D.: A variational analysis of the toda system on compact surfaces. *Comm. Pure Appl. Math.* 66(3), 322371 (2013)
- [27] Nolasco M., Tarantello G.: Vortex condensates for the  $SU(3)$  Chern-Simons theory. *Commun. Math. Phys.* 213, 599639 (2000).
- [28] Ohtsuka H., Suzuki T.: Blow-up analysis for  $SU(3)$  Toda system. *J. Diff. Eq.* 232(2), 419440 (2007).
- [29] Spruck J., Yang Y.: On Multivortices in the Electroweak Theory I:Existence of Periodic Solutions. *Commun. Math. Phys.* 144, 116 (1992).
- [30] Struwe M., Tarantello G.: On multivortex solutions in Chern-Simons gauge theory. *Boll. Unione Mat. Ital., Sez. B, Artic. Ric. Mat.* 8(1), 109121 (1998)

- [31] Suzuki, Takashi Global analysis for a two-dimensional elliptic eigenvalue problem with the exponential nonlinearity. Ann. Inst. H. Poincar Anal. Non Linéaire 9 (1992), no. 4, 367–397.
- [32] Tarantello G.: Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. Disc. Cont. Dyn. Syst. 28(3), 931–973 (2010).
- [33] Tarantello, G; Blow-up analysis for a cosmic strings equation. J. Funct. Anal. 272 (2017), no. 1, 255–338.
- [34] Troyanov, Marc; Prescribing curvature on compact surfaces with conical singularities. Trans. Amer. Math. Soc. 324 (1991), no. 2, 793–821.
- [35] Troyanov, M.: Metrics of constant curvature on a sphere with two conical singularities. Proc. Third Int. Symp. on Diff. Geom. (Peniscola 1988), Lect. Notes in Math. 1410, Berlin: Springer-Verlag, 1991, pp. 296–306
- [36] Yang, Y.: Solitons in Field Theory and Nonlinear Analysis. Springer Monographs in Mathematics, New York: Springer, 2001
- [37] Zhang, L.: Blowup solutions for some nonlinear elliptic equations involving exponential nonlinearities, Comm. Math. Phys., 268, (2006) no 1: 105–133.
- [38] Zhang L.: Asymptotic behavior of blowup solutions for elliptic equations with exponential nonlinearity and singular data. Comm. Contemp. Math. 11(3), 395–411 (2009)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER,  
B.C., CANADA V6T 1Z2  
*E-mail address:* jcwei@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 358 LITTLE HALL P.O.Box  
118105, GAINESVILLE FL 32611-8105  
*E-mail address:* leizhang@ufl.edu