

1 CURVE SKETCHING NOTES

1.1 Tools. Let f be differentiable as needed on (a, b) .

Fact (First derivative). (1) If $f'(x) > 0$ for all $x \in (a, b)$ then f is strictly increasing there.
 (2) If $f'(x) < 0$ for all $x \in (a, b)$ then f is strictly decreasing there.

Every change involves either a *critical point* (f' vanishes) or a *singularity* (f' undefined).

Fact (Second derivative). (1) If $f''(x) > 0$ for all $x \in (a, b)$ then f is concave up there.
 (2) If $f''(x) < 0$ for all $x \in (a, b)$ then f is concave down there.

Definition. A change in concavity is called an *inflection point*.

Theorem. (Tests for minima and maxima) Let $x_0 \in (a, b)$ be a critical or singular number for f , and suppose f is continuous at x_0 , differentiable near it.

- (1) Either of the following is sufficient to show that f has a local minimum at x_0 :
 - (a) $f''(x_0) > 0$ or;
 - (b) $f'(x)$ is negative to the left of x_0 , positive to its right.
- (2) Either of the following shows that f has a local maximum at x_0 :
 - (a) $f''(x_0) < 0$ or;
 - (b) $f'(x)$ is positive to the left of x_0 , negative to its right.

1.2 Curve sketching protocol. Given a function f .

0th derivative stuff:

- (a) The domain and the domain of continuity.
- (b) Domains where $f > 0$, $f < 0$.
- (c) Anchor points: x - and y -intercepts.
- (d) Horizontal and vertical asymptotes.

1st derivative stuff:

- (a) Evaluate $f'(x)$ [**high stakes: error here loses a lot of points down the line**]
 Using this, determine:
- (b) Domains where $f' > 0$, $f' < 0$
- (c) Critical and singular points.

2nd derivative stuff:

- (a) Domains where $f'' > 0$, $f'' < 0$
- (b) Points where $f''(x) = 0$, inflection points.

2 CURVE SKETCHING EXAMPLES

$$2.1 \quad f(x) = \frac{x^2-9}{x^2+3}.$$

$$\bullet \quad f'(x) \stackrel{\text{quot}}{=} \frac{2x(x^2+3)-2x(x^2-9)}{(x^2+3)^2} = \frac{24x}{(x^2+3)^2}.$$

$$\bullet \quad f''(x) = 24 \frac{1}{(x^2+3)^2} - 24 \frac{x \cdot 2 \cdot 2x}{(x^2+3)^3} = 24 \frac{(x^3+3)-4x^2}{(x^2+3)^3} = 72 \frac{1-x^2}{(x^2+3)^3}.$$

Thus

(1) f defined on \mathbb{R} , cts everywhere (defined by formula; denominator everywhere nonzero).

Moreover

(a) $f(0) = -3$, $f(x) = \frac{(x-3)(x+3)}{x^2+3}$ so vanishes at $x = \pm 3$, negative between them, positive otherwise.


(b) $\lim_{x \rightarrow \pm\infty} f(x) = \lim_{x \rightarrow \pm\infty} \frac{1-9/x^2}{1+3/x^2} = 1$.

(2) $f'(x)$ is negative for $x < 0$, zero at $x = 0$, positive at $x > 0$ (hence at the critical number $x = 0$ we have a local minimum)

(3) $f''(x)$ has the same sign as $(1-x)^2 = (1-x)(1+x)$ so it is negative if $x < -1$ or $x > 1$, positive if $-1 < x < +1$ and zero at $x = \pm 1$ which are therefore inflection points.

The "special" points were: $-3, -1, 0, 1, 3$ so we break up the domain of f at those points:

x	$(-\infty, -3)$	-3	$(-3, -1)$	-1	$(-1, 0)$	0	$(0, 1)$	1	$(1, 3)$	3	$(3, \infty)$
f	+	0	-	-2	-	-3	-	-2	-	0	+
f'	+	+	+	+	+	0	+	+	+	+	+
f''	-	-	-	0	-	-	-	0	-	-	-



2.2 $f(x) = x^{2/3}(x-1)$.

$$\begin{aligned} \bullet f'(x) &= \frac{2}{3}x^{-1/3}(x-1) + x^{2/3} = \frac{2(x-1)+3x}{3x^{1/3}} = \frac{5x-2}{3x^{1/3}} \\ \bullet f''(x) &= \frac{5}{3x^{1/3}} - \frac{5x-2}{9x^{4/3}} = \frac{15x-(5x-2)}{9x^{4/3}} = \frac{10x+2}{9x^{4/3}} \end{aligned}$$

Thus (note: $x^{2/3}$ and $x^{4/3}$ are always non-negative; $x^{1/3}$ has the same sign as x)

- (1) f defined on \mathbb{R} ($x^{1/3}$ defined everywhere), continuous there (defined by formula).
 - (a) $f(0) = 0$, $f(1) = 0$ and f is positive if $x < 1$ negative if $x > 1$ ($x^{2/3} \geq 0$ for all x)
 - (b) $\lim_{x \rightarrow \pm\infty} |f(x)| = \infty$ so no horizontal asymptotes.
- (2) The critical numbers are 0 (f' undefined) and $\frac{2}{5}$ ($f' = 0$). Otherwise $f' > 0$ if $x < 0$, $f' < 0$ if $0 < x < \frac{2}{5}$ and $f' > 0$ if $x > \frac{2}{5}$.
- (3) . Thus f'' is undefined at 0, vanishes at $-\frac{1}{5}$, and is negative if $x < -\frac{1}{5}$, positive if $-\frac{1}{5} < x < 0$ or $x > 0$, so only $-\frac{1}{5}$ is an inflection point.

Summary table:

x	$(-\infty, -\frac{1}{5})$	$-\frac{1}{5}$	$(-\frac{1}{5}, 0)$	0	$(0, \frac{2}{5})$	$\frac{2}{5}$	$(\frac{2}{5}, 1)$	1	$(1, \infty)$
f	-	$-\frac{6}{5^{5/3}}$	-	0	-	$-\frac{3 \cdot 4^{1/3}}{5^{5/3}}$	-	0	+
f'	+	+	+	undef	-	0	+	+	+
f''	-	0	+	undef	+	+	+	+	+

