

Lior Silberman's Math 412: Problem Set 9 (due 24/3/2023)

Practice: Norms

P1. Call two norms $\|\cdot\|_1, \|\cdot\|_2$ on V *equivalent* if there are constants $c, C > 0$ such that for all $\underline{v} \in V$,

$$c \|\underline{v}\|_1 \leq \|\underline{v}\|_2 \leq C \|\underline{v}\|_1.$$

- (a) Show that this is an equivalence relation.
- (b) Suppose the two norms are equivalent and that $\lim_{n \rightarrow \infty} \|\underline{v}_n\|_1 = 0$ (that is, that $\underline{v}_n \xrightarrow[n \rightarrow \infty]{\|\cdot\|_1} \underline{0}$). Show that $\lim_{n \rightarrow \infty} \|\underline{v}_n\|_2 = 0$ (that is, that $\underline{v}_n \xrightarrow[n \rightarrow \infty]{\|\cdot\|_2} \underline{0}$).
- (*c) Show the converse of (b) also holds. In other words, two norms are equivalent iff they determine the same notion of convergence.

Norms

1. Let $f(x) = x^2$ on $[-1, 1]$.
 - (a) For $1 \leq p < \infty$. Calculate $\|f\|_{L^p} = \left(\int_{-1}^1 |f(x)|^p dx \right)^{1/p}$.
 - (b) Calculate $\|f\|_{L^\infty} = \sup \{|f(x)| : -1 \leq x \leq 1\}$. Check that $\lim_{p \rightarrow \infty} \|f\|_{L^p} = \|f\|_{L^\infty}$.
 - (c) Calculate $\|f\|_{H^2} = \left(\|f\|_{L^2}^2 + \|f'\|_{L^2}^2 + \|f''\|_{L^2}^2 \right)^{1/2}$.

SUPP Show that the H^2 norm is equivalent to the norm $\left(\|f\|_{L^2}^2 + \|f''\|_{L^2}^2 \right)^{1/2}$.
2. Let $A \in M_n(\mathbb{R})$. Write $\|A\|_p$ for its $\ell^p \rightarrow \ell^p$ operator norm.
 - (a) Show $\|A\|_1 = \max_j \sum_{i=1}^n |a_{ij}|$.
 - (b) Show that $\|A\|_\infty = \max_i \sum_{j=1}^n |a_{ij}|$.

RMK See below on *duality*.
3. The *spectral radius* of $A \in M_n(\mathbb{C})$ is the magnitude of its largest eigenvalue: $\rho(A) = \max \{|\lambda| \mid \lambda \in \text{Spec}(A)\}$.
 - (a) Show that for any norm $\|\cdot\|$ on \mathbb{C}^n and any $A \in M_n(\mathbb{C})$, $\rho(A) \leq \|A\|$.
 - (b) Suppose that A is diagonalizable. Show that there is a norm on \mathbb{C}^n such that $\|A\| = \rho(A)$.
 - (*c) Show that if A is Hermitian then $\|A\|_2 = \rho(A)$.
 - (d) Show that if A, B are similar, and $\|\cdot\|$ is any norm in \mathbb{C}^n , then $\lim_{m \rightarrow \infty} \|A^m\|^{1/m} = \lim_{m \rightarrow \infty} \|B^m\|^{1/m}$ (in the sense that, if one limit exists, then so does the other, and they are equal).
 - **e) Show that for any norm on \mathbb{C}^n and any $A \in M_n(\mathbb{C})$, we have $\lim_{m \rightarrow \infty} \|A^m\|^{1/m} = \rho(A)$.

Supplement: one more norm

- A. The *Hilbert-Schmidt* norm on $M_n(\mathbb{C})$ is $\|A\|_{\text{HS}} = \left(\sum_{i,j=1}^n |a_{ij}|^2 \right)^{1/2}$.
- PRAC Verify that $\|A\|_{\text{HS}} = (\text{Tr}(A^\dagger A))^{1/2}$.
- (a) Show that $\|\cdot\|_{\text{HS}}$ is, indeed, a norm.
 - (b) Show that $\|A\|_2 \leq \|A\|_{\text{HS}}$.

Supplement: Norms and constructions

B. (Direct sum) Let $\{(V_i, \|\cdot\|_i)\}_{i=1}^n$ be normed spaces, and let $1 \leq p \leq \infty$. For $\underline{v} = (v_i) \in \bigoplus_{i=1}^n V_i$ define

$$\|\underline{v}\| = \left(\sum_{i=1}^n \|v_i\|_i^p \right)^{1/p}.$$

Show that this defines a norm on $\bigoplus_{i=1}^n V_i$.

DEF This operation is called the L^p -sum of the normed spaces.

- C. (Quotient) Let $(V, \|\cdot\|)$ be a normed space, and let $W \subset V$ be a subspace. For $\underline{v} + W \in V/W$ set $\|\underline{v} + W\|_{V/W} = \inf \{\|\underline{v} + \underline{w}\| : \underline{w} \in W\}$.
- (a) Show that $\|\cdot\|_{V/W}$ is 1-homogenous and satisfies the triangle inequality (a “seminorm”).
- (b) Show that $\|\underline{v} + W\|_{V/W} = 0$ iff \underline{v} is in the closure of W , so that $\|\cdot\|_{V/W}$ is a norm iff W is closed in V .

For duality in norms see problems A, B. Norming tensor product spaces is complicated.

- D. Let V be a normed space. The operator norm on $V^* = \text{Hom}_{\text{cts}}(V, \mathbb{R})$ is called the *dual norm*.
- (a) Let $V = \mathbb{R}^n$ and identify V^* with \mathbb{R}^n via the usual pairing. Show that the norm on V^* dual to the ℓ^1 -norm is the ℓ^∞ norm and vice versa. Show that the ℓ^2 -norm is self-dual.
- (b) Use A(a),(b) to show that the dual to the ℓ^p norm on \mathbb{R}^n is the ℓ^q norm where $\frac{1}{p} + \frac{1}{q} = 1$.
- (c) Let U be another normed space and let $T: U \rightarrow V$ be bounded. Let $T': V' \rightarrow U'$ be the algebraic dual map as defined in this course. Show that for every $\underline{v}^* \in V^* \subset V'$, $T'\underline{v}^* \in U^*$ (that is, it is continuous). We write $T^*: V^* \rightarrow U^*$ for the dual map restricted to continuous functionals.
- (d) Show that T^* is itself bounded, in that $\|T^*\|_{V^* \rightarrow U^*} \leq \|T\|_{U \rightarrow V}$.
- E. A *seminorm* on a vector space V is a map $V \rightarrow \mathbb{R}_{\geq 0}$ that satisfies all the conditions of a norm except that it can be zero for non-zero vectors.
- (a) Show that for any $f \in V'$, $\varphi(\underline{v}) = |f(\underline{v})|$ is a seminorm.
- (b) Construct a seminorm on \mathbb{R}^2 not of this form.
- (c) Let Φ be a family of seminorms on V which is pointwise bounded. Show that $\bar{\varphi}(\underline{v}) = \sup \{\varphi(\underline{v}) \mid \varphi \in \Phi\}$ is again a seminorm.

Supplementary problem: Continuity

- F. Let V, W be normed vector spaces, equipped with the metric topology coming from the norm. Let $T \in \text{Hom}_F(V, W)$. Show that the following are equivalent:
- (1) T is continuous.
 - (2) T is continuous at zero.
 - (3) T is *bounded*: $\|T\|_{V \rightarrow W} < \infty$, that is: for some $C > 0$ and all $\underline{v} \in V$, $\|T\underline{v}\|_W \leq C \|\underline{v}\|_V$.
- Hint: the same idea is used in problem P1

Supplementary problems: Completeness

- G. Let $\{\underline{v}_n\}_{n=1}^\infty$ be a Cauchy sequence in a normed space. Show that $\{\|\underline{v}_n\|\}_{n=1}^\infty \subset \mathbb{R}_{\geq 0}$ is a Cauchy sequence.
- H. (The completion) Let (X, d) be a metric space.
- (a) Let $\{x_n\}, \{y_n\} \subset X$ be two Cauchy sequences. Show that $\{d(x_n, y_n)\}_{n=1}^\infty \subset \mathbb{R}$ is a Cauchy sequence.
- DEF Let (\tilde{X}, \tilde{d}) denote the set of Cauchy sequences in X with the distance $\tilde{d}(\underline{x}, \underline{y}) = \lim_{n \rightarrow \infty} d(x_n, y_n)$.
- (b) Show that \tilde{d} satisfies all the axioms of a metric except that it can be non-zero for distinct sequences.
- (c) Show that the relation $\underline{x} \sim \underline{y} \iff \tilde{d}(\underline{x}, \underline{y}) = 0$ is an equivalence relation.
- (d) Let $\hat{X} = \tilde{X} / \sim$ be the set of equivalence classes. Show that $\hat{d}: \hat{X} \times \hat{X} \rightarrow \mathbb{R}_{\geq 0}$ descends to a well-defined function $\hat{d}: \hat{X} \times \hat{X} \rightarrow \mathbb{R}_{\geq 0}$ which is a metric.
- (e) Show that (\hat{X}, \hat{d}) is a complete metric space.
- DEF For $x \in X$ let $\iota(x) \in \hat{X}$ be the equivalence class of the constant sequence x .
- (f) Show that $\iota: X \rightarrow \hat{X}$ is an isometric embedding with dense image.
- (g) (Universal property) Show that for any complete metric space (Y, d_Y) and any uniformly continuous $f: X \rightarrow Y$ there is a unique extension $\hat{f}: \hat{X} \rightarrow Y$ such that $\hat{f} \circ \iota = f$.
- (h) Show that triples $(\hat{X}, \hat{d}, \iota)$ satisfying the property of (g) are unique up to a unique isomorphism.

- I. (Complete fields) An *absolute value* on a field F is a map $|\cdot| : F \rightarrow \mathbb{R}_{\geq 0}$ such that (a) $|xy| = |x||y|$ (b) $|x| = 0 \Leftrightarrow x = 0$ (c) $|x + y| \leq |x| + |y|$.

DEF Fix a prime number p . For $x \in \mathbb{Q}^\times$ write $x = \frac{a}{b}p^k$ for some non-zero $a, b \in \mathbb{Z}$ prime to p and $k \in \mathbb{Z}$ and set $|x|_p = p^{-k}$ (also, $|0|_p = 0$).

- (a) Show that $|\cdot|_p$ is an absolute value on \mathbb{Q} satisfying the *ultrametric inequality* $|x + y|_p \leq \max\{|x|_p, |y|_p\}$.
 (b) Let $|\cdot|$ be an absolute value on F . Show that $d(x, y) = |x - y|$ is a metric on F .
 (c) Show that (with respect to the metric of (b)) the absolute value is uniformly continuous $F \rightarrow \mathbb{R}_{\geq 0}$, addition is a uniformly continuous map $F \times F \rightarrow F$, and that for any $r > 0$ multiplication is a uniformly continuous map $B(0, r) \times B(0, r) \rightarrow B(0, r^2)$.
 (d) Let \hat{F} be the completion of F wrt $|\cdot|$. Show that the absolute value and the operations of addition and multiplication extend to maps $|\cdot| : \hat{F} \rightarrow \mathbb{R}_{\geq 0}$, $+, \cdot : \hat{F} \times \hat{F} \rightarrow \hat{F}$ giving it the structure of a ring with an absolute value $|\cdot|$.
 (e) Show that every non-zero element of \hat{F} has an inverse, that is that \hat{F} is a field.

DEF Write \mathbb{Q}_p for the completion of \mathbb{Q} wrt $|\cdot|$.

FACT Closed bounded sets in \mathbb{Q}_p are compact, as in \mathbb{R} .